

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
19 September 2002 (19.09.2002)

PCT

(10) International Publication Number
WO 02/071928 A2

- (51) International Patent Classification: A61B
- (21) International Application Number: PCT/US02/07826
- (22) International Filing Date: 14 March 2002 (14.03.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
- | | | |
|------------|--------------------------------|----|
| 60/276,025 | 14 March 2001 (14.03.2001) | US |
| 60/276,026 | 14 March 2001 (14.03.2001) | US |
| 60/311,732 | 10 August 2001 (10.08.2001) | US |
| 60/323,580 | 19 September 2001 (19.09.2001) | US |
| 60/325,149 | 26 September 2001 (26.09.2001) | US |
| 60/324,967 | 26 September 2001 (26.09.2001) | US |
| 60/325,102 | 26 September 2001 (26.09.2001) | US |

(71) Applicant (for all designated States except US): MILLENNIUM PHARMACEUTICALS, INC. [US/US]; 75 Sidney Street, Cambridge, MA 02139 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): MONAHAN, John, E. [US/US]; 942 West Street, Walpole, MA 02081 (US). GANNAVAPU, Manjula [IN/US]; 10 Windemere Drive, Acton, MA 01720 (US). HOERSCH, Sebastian [US/US]; 127 Brattle Street, Arlington, MA 02474 (US). KAMATKAR, Shubhangi [US/US]; 655 Saw Mill Brook Parkway, Apt. 1, Newton, MA 02459 (US). KOVATIS, Steven, G. [US/US]; 94 Aldrich Road, Wilmington, MA 01887 (US). MEYERS, Rachel, E. [US/US]; 115 Devonshire Road, Newton, MA 02468 (US). MORRISEY, Michael, P. [US/US]; 140 Kenrick Street, Apt. 32, Brighton, MA 02135 (US). OLANDT, Peter, J. [US/US]; 29 Florence Street, Newton, MA 02459 (US). SEN, Ami [US/US]; 66 Dinsmore Avenue, Apt. 507, Framingham,

MA 01702 (US). VIEBY, Petter, Ole [US/US]; 16 Nipmuck Drive, Westborough, MA 01581 (US). MILLS, Gordon, B. [CA/US]; 4124 Amherst Street, Houston, TX 77005 (US). BAST, Robert, C., Jr. [US/US]; 14 Memorial Point Lane, Houston, TX 77024 (US). LI, Karen [US/US]; 4127 Amherst Street, Houston, TX 77005 (US). SCHMANDT, Rosemarie, E. [CA/US]; 7300 Brompton Road, Apt. 5512, Houston, TX 77025 (US). ZHAO, Xomei [US/US]; 6 Wildwood Lane, Burlington, MA 01803 (US). GLATT, Karen [US/US]; 17 Beacon Street, Natick, MA 01760 (US).

(74) Agents: SMITH, DeAnn, F. et al.; Lahive & Cockfield, LLP, 28 State Street, Boston, MA 02109 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NUCLEIC ACID MOLECULES AND PROTEINS FOR THE IDENTIFICATION, ASSESSMENT, PREVENTION, AND THERAPY OF OVARIAN CANCER

(57) Abstract: The invention relates to newly discovered nucleic acid molecules and proteins associated with ovarian cancer. Compositions, kits, and methods for detecting, characterizing, preventing, and treating human ovarian cancers are provided.

- 1 -

NUCLEIC ACID MOLECULES AND PROTEINS FOR THE IDENTIFICATION,
ASSESSMENT, PREVENTION, AND THERAPY OF
OVARIAN CANCER

5

RELATED APPLICATIONS

The present application claims priority from U.S. provisional patent application serial no. 60/276,025, filed on March 14, 2001, which was abandoned on September 25, 2001, and from U.S. provisional patent application serial no. 60/325,149, filed on September 26, 2001. The present application also claims priority from U.S. provisional patent application serial no. 60/276,026, filed on March 14, 2001, which was abandoned on September 25, 2001, and from U.S. provisional patent application serial no. 60/324,967, filed September 26, 2001. The present application additionally claims priority from U.S. provisional patent application serial no. 60/311,732, filed August 10, 2001, which was abandoned on September 25, 2001, and from U.S. provisional patent application serial no. 60/325,102, filed September 26, 2001. The present application also claims priority from U.S. provisional patent application serial no. 60/323,580, filed September 19, 2001. All of the above applications are expressly incorporated by reference.

20

FIELD OF THE INVENTION

The field of the invention is ovarian cancer, including diagnosis, characterization, management, and therapy of ovarian cancer.

BACKGROUND OF THE INVENTION

25

Ovarian cancer is responsible for significant morbidity and mortality in populations around the world. Ovarian cancer is classified, on the basis of clinical and pathological features, in three groups, namely epithelial ovarian cancer (EOC; >90% of ovarian cancer in Western countries), germ cell tumors (*circa* 2-3% of ovarian cancer), and stromal ovarian cancer (*circa* 5% of ovarian cancer; Ozols *et al.*, 1997, *Cancer Principles and Practice of Oncology*, 5th ed., DeVita *et al.*, Eds. pp. 1502). Relative to EOC, germ cell tumors and stromal ovarian cancers are more easily detected and treated

30

- 2 -

at an early stage, translating into higher/better survival rates for patients afflicted with these two types of ovarian cancer.

There are numerous types of ovarian tumors, some of which are benign, and others of which are malignant. Treatment (including non-treatment) options and
5 predictions of patient outcome depend on accurate classification of the ovarian cancer. Ovarian cancers are named according to the type of cells from which the cancer is derived and whether the ovarian cancer is benign or malignant. Recognized histological tumor types include, for example, serous, mucinous, endometrioid, and clear cell tumors. In addition, ovarian cancers are classified according to recognized grade and
10 stage scales.

In grade I, the tumor tissue is well differentiated. In grade II, tumor tissue is moderately well differentiated. In grade III, the tumor tissue is poorly differentiated. This grade correlates with a less favorable prognosis than grades I and II. Stage I is generally confined within the capsule surrounding one (stage IA) or both
15 (stage IB) ovaries, although in some stage I (*i.e.* stage IC) cancers, malignant cells may be detected in ascites, in peritoneal rinse fluid, or on the surface of the ovaries. Stage II involves extension or metastasis of the tumor from one or both ovaries to other pelvic structures. In stage IIA, the tumor extends or has metastasized to the uterus, the fallopian tubes, or both. Stage IIB involves extension of the tumor to the pelvis. Stage
20 IIC is stage IIA or IIB in which malignant cells may be detected in ascites, in peritoneal rinse fluid, or on the surface of the ovaries. In stage III, the tumor comprises at least one malignant extension to the small bowel or the omentum, has formed extrapelvic peritoneal implants of microscopic (stage IIIA) or macroscopic (< 2 centimeter diameter, stage IIIB; > 2 centimeter diameter, stage IIIC) size, or has metastasized to a
25 retroperitoneal or inguinal lymph node (an alternate indicator of stage IIIC). In stage IV, distant (*i.e.* non-peritoneal) metastases of the tumor can be detected.

The durations of the various stages of ovarian cancer are not presently known, but are believed to be at least about a year each (Richart *et al.*, 1969, *Am. J. Obstet. Gynecol.* 105:386). Prognosis declines with increasing stage designation. For
30 example, 5-year survival rates for patients diagnosed with stage I, II, III, and IV ovarian cancer are 80%, 57%, 25%, and 8%, respectively.

- 3 -

Despite being the third most prevalent gynecological cancer, ovarian cancer is the leading cause of death among those afflicted with gynecological cancers. The disproportionate mortality of ovarian cancer is attributable to a substantial absence of symptoms among those afflicted with early-stage ovarian cancer and to difficulty
5 diagnosing ovarian cancer at an early stage. Patients afflicted with ovarian cancer most often present with non-specific complaints, such as abnormal vaginal bleeding, gastrointestinal symptoms, urinary tract symptoms, lower abdominal pain, and generalized abdominal distension. These patients rarely present with paraneoplastic
10 symptoms or with symptoms which clearly indicate their affliction. Presently, less than about 40% of patients afflicted with ovarian cancer present with stage I or stage II. Management of ovarian cancer would be significantly enhanced if the disease could be detected at an earlier stage, when treatments are much more generally efficacious.

Ovarian cancer may be diagnosed, in part, by collecting a routine medical history from a patient and by performing physical examination, x-ray examination, and
15 chemical and hematological studies on the patient. Hematological tests which may be indicative of ovarian cancer in a patient include analyses of serum levels of proteins designated CA125 and DF3 and plasma levels of lysophosphatidic acid (LPA). Palpation of the ovaries and ultrasound techniques (particularly including endovaginal
20 ultrasound and color Doppler flow ultrasound techniques) can aid detection of ovarian tumors and differentiation of ovarian cancer from benign ovarian cysts. However, a definitive diagnosis of ovarian cancer typically requires performing exploratory laparotomy of the patient.

Potential tests for the detection of ovarian cancer (*e.g.*, screening, reflex or monitoring) may be characterized by a number of factors. The "sensitivity" of an
25 assay refers to the probability that the test will yield a positive result in an individual afflicted with ovarian cancer. The "specificity" of an assay refers to the probability that the test will yield a negative result in an individual not afflicted with ovarian cancer. The "positive predictive value" (PPV) of an assay is the ratio of true positive results (*i.e.* positive assay results for patients afflicted with ovarian cancer) to all positive results
30 (*i.e.* positive assay results for patients afflicted with ovarian cancer + positive assay results for patients not afflicted with ovarian cancer). It has been estimated that in order for an assay to be an appropriate population-wide screening tool for ovarian cancer the

assay must have a PPV of at least about 10% (Rosenthal *et al.*, 1998, *Sem. Oncol.* 25:315-325). It would thus be desirable for a screening assay for detecting ovarian cancer in patients to have a high sensitivity and a high PPV. Monitoring and reflex tests would also require appropriate specifications.

- 5 Owing to the cost, limited sensitivity, and limited specificity of known methods of detecting ovarian cancer, screening is not presently performed for the general population. In addition, the need to perform laparotomy in order to diagnose ovarian cancer in patients who screen positive for indications of ovarian cancer limits the desirability of population-wide screening, such that a PPV even greater than 10%
10 would be desirable.

- Prior use of serum CA125 level as a diagnostic marker for ovarian cancer indicated that this method exhibited insufficient specificity for use as a general screening method. Use of a refined algorithm for interpreting CA125 levels in serial retrospective samples obtained from patients improved the specificity of the method
15 without shifting detection of ovarian cancer to an earlier stage (Skakes, 1995, *Cancer* 76:2004). Screening for LPA to detect gynecological cancers including ovarian cancer exhibited a sensitivity of about 96% and a specificity of about 89%. However, CA125-based screening methods and LPA-based screening methods are hampered by the presence of CA125 and LPA, respectively, in the serum of patients afflicted with
20 conditions other than ovarian cancer. For example, serum CA125 levels are known to be associated with menstruation, pregnancy, gastrointestinal and hepatic conditions such as colitis and cirrhosis, pericarditis, renal disease, and various non-ovarian malignancies. Serum LPA is known, for example, to be affected by the presence of non-ovarian gynecological malignancies. A screening method having a greater specificity for
25 ovarian cancer than the current screening methods for CA125 and LPA could provide a population-wide screening for early stage ovarian cancer.

- Presently greater than about 60% of ovarian cancers diagnosed in patients are stage III or stage IV cancers. Treatment at these stages is largely limited to cytoreductive surgery (when feasible) and chemotherapy, both of which aim to slow the
30 spread and development of metastasized tumor. Substantially all late stage ovarian cancer patients currently undergo combination chemotherapy as primary treatment, usually a combination of a platinum compound and a taxane. Median survival for

- 5 -

responding patients is about one year. Combination chemotherapy involving agents such as doxorubicin, cyclophosphamide, cisplatin, hexamethylmelamine, paclitaxel, and methotrexate may improve survival rates in these groups, relative to single-agent therapies. Various recently-developed chemotherapeutic agents and treatment regimens have also demonstrated usefulness for treatment of advanced ovarian cancer. For example, use of the topoisomerase I inhibitor topotecan, use of amifostine to minimize chemotherapeutic side effects, and use of intraperitoneal chemotherapy for patients having peritoneally implanted tumors have demonstrated at least limited utility. Presently, however, the 5-year survival rate for patients afflicted with stage III ovarian cancer is 25%, and the survival rate for patients afflicted with stage IV ovarian cancer is 8%.

In summary, the earlier ovarian cancer is detected, the aggressiveness of therapeutic intervention and the side effects associated with therapeutic intervention are minimized. More importantly, the earlier the cancer is detected, the survival rate and quality of life of ovarian cancer patients is enhanced. Thus, a pressing need exists for methods of detecting ovarian cancer as early as possible. There also exists a need for methods of detecting recurrence of ovarian cancer as well as methods for predicting and monitoring the efficacy of treatment. There further exists a need for new therapeutic methods for treating ovarian cancer. The present invention satisfies these needs.

20

SUMMARY OF THE INVENTION

The invention relates to cancer markers (hereinafter "markers" or "markers of the inventions"), which are listed in Tables 1-3. The invention provides nucleic acids and proteins that are encoded by or correspond to the markers (hereinafter "marker nucleic acids" and "marker proteins," respectively). The invention further provides antibodies, antibody derivatives and antibody fragments which bind specifically with such proteins and/or fragments of the proteins.

In one aspect, the invention relates to various diagnostic, monitoring, test and other methods related to ovarian cancer detection and therapy. In one embodiment, the invention provides a diagnostic method of assessing whether a patient has ovarian cancer or has higher than normal risk for developing ovarian cancer, comprising the steps of comparing the level of expression of a marker of the invention in a patient

- 6 -

sample and the normal level of expression of the marker in a control, *e.g.*, a sample from a patient without ovarian cancer. A significantly higher level of expression of the marker in the patient sample as compared to the normal level is an indication that the patient is afflicted with ovarian cancer or has higher than normal risk for developing
5 ovarian cancer.

In a preferred embodiment of the diagnostic method, the marker is over-expressed by at least two-fold in at least about 20% of stage I ovarian cancer patients, stage II ovarian cancer patients, stage III ovarian cancer patients, stage IV ovarian cancer patients, grade I ovarian cancer patients, grade II ovarian cancer patients, grade
10 III ovarian cancer patients, epithelial ovarian cancer patients, stromal ovarian cancer patients, germ cell ovarian cancer patients, malignant ovarian cancer patients, benign ovarian cancer patients, serous neoplasm ovarian cancer patients, mucinous neoplasm ovarian cancer patients, endometrioid neoplasm ovarian cancer patients and/or clear cell neoplasm ovarian cancer patients.

The diagnostic methods of the present invention are particularly useful for patients with an identified pelvic mass or symptoms associated with ovarian cancer. The methods of the present invention can also be of particular use with patients having an enhanced risk of developing ovarian cancer (*e.g.*, patients having a familial history of ovarian cancer, patients identified as having a mutant oncogene, and patients at least
20 about 50 years of age).

In a preferred diagnostic method of assessing whether a patient is afflicted with ovarian cancer (*e.g.*, new detection ("screening"), detection of recurrence, reflex testing), the method comprises comparing:

- a) the level of expression of a marker of the invention in a patient sample,
25 and
- b) the normal level of expression of the marker in a control non-ovarian cancer sample.

A significantly higher level of expression of the marker in the patient sample as compared to the normal level is an indication that the patient is afflicted with ovarian
30 cancer.

- 7 -

The invention also provides diagnostic methods for assessing the efficacy of a therapy for inhibiting ovarian cancer in a patient. Such methods comprise comparing:

- 5 a) expression of a marker of the invention in a first sample obtained from the patient prior to providing at least a portion of the therapy to the patient, and
- b) expression of the marker in a second sample obtained from the patient following provision of the portion of the therapy.

10 A significantly lower level of expression of the marker in the second sample relative to that in the first sample is an indication that the therapy is efficacious for inhibiting ovarian cancer in the patient.

It will be appreciated that in these methods the "therapy" may be any therapy for treating ovarian cancer including, but not limited to, chemotherapy, radiation therapy, surgical removal of tumor tissue, gene therapy and biologic therapy such as the
15 administering of antibodies and chemokines. Thus, the methods of the invention may be used to evaluate a patient before, during and after therapy, for example, to evaluate the reduction in tumor burden.

In a preferred embodiment, the diagnostic methods of the present invention are directed to therapy using a chemical or biologic agent. These methods
20 comprise comparing:

- a) expression of a marker of the invention in a first sample obtained from the patient and maintained in the presence of the chemical or biologic agent, and
- 25 b) expression of the marker in a second sample obtained from the patient and maintained in the absence of the agent.

A significantly lower level of expression of the marker in the first sample relative to that in the second sample is an indication that the agent is efficacious for inhibiting ovarian cancer in the patient. In one embodiment, the first and second samples can be portions of a single sample obtained from the patient or portions of pooled samples obtained
30 from the patient.

- 8 -

The invention additionally provides a monitoring method for assessing the progression of ovarian cancer in a patient, the method comprising:

- a) detecting in a patient sample at a first time point, the expression of a marker of the invention;
- 5 b) repeating step a) at a subsequent time point in time; and
- c) comparing the level of expression detected in steps a) and b), and therefrom monitoring the progression of ovarian cancer in the patient.

A significantly higher level of expression of the marker in the sample at the subsequent time point from that of the sample at the first time point is an indication that the ovarian
10 cancer has progressed, whereas a significantly lower level of expression is an indication that the ovarian cancer has regressed.

The invention further provides a diagnostic method for determining whether ovarian cancer has metastasized or is likely to metastasize in the future, the method comprising comparing:

- 15 a) the level of expression of a marker of the invention in a patient sample, and
- b) the normal level (or non-metastatic level) of expression of the marker in a control sample.

A significantly higher level of expression in the patient sample as compared to the
20 normal level (or non-metastatic level) is an indication that the ovarian cancer has metastasized or is likely to metastasize in the future.

The invention moreover provides a test method for selecting a composition for inhibiting ovarian cancer in a patient. This method comprises the steps of:

- 25 a) obtaining a sample comprising cancer cells from the patient;
- b) separately maintaining aliquots of the sample in the presence of a plurality of test compositions;
- c) comparing expression of a marker of the invention in each of the aliquots; and
- 30 d) selecting one of the test compositions which significantly reduces the level of expression of the marker in the aliquot containing that test

- 9 -

composition, relative to the levels of expression of the marker in the presence of the other test compositions.

The invention additionally provides a test method of assessing the ovarian carcinogenic potential of a compound. This method comprises the steps of:

- 5 a) maintaining separate aliquots of ovarian cells in the presence and absence of the compound; and
- b) comparing expression of a marker of the invention in each of the aliquots.

A significantly higher level of expression of the marker in the aliquot maintained in the presence of the compound, relative to that of the aliquot maintained in the absence of the compound, is an indication that the compound possesses ovarian carcinogenic potential.

In addition, the invention further provides a method of inhibiting ovarian cancer in a patient. This method comprises the steps of:

- a) obtaining a sample comprising cancer cells from the patient;
- 15 b) separately maintaining aliquots of the sample in the presence of a plurality of compositions;
- c) comparing expression of a marker of the invention in each of the aliquots; and
- d) administering to the patient at least one of the compositions which
- 20 significantly lowers the level of expression of the marker in the aliquot containing that composition, relative to the levels of expression of the marker in the presence of the other compositions.

In the aforementioned methods, the samples or patient samples comprise cells obtained from the patient. The cells may be found in an ovarian tissue sample

25 collected, for example, by an ovarian tissue biopsy or histology section. In one embodiment, the patient sample is an ovary-associated body fluid. Such fluids include, for example, blood fluids, lymph, ascites fluids, gynecological fluids, cystic fluids, urine, and fluids collected by peritoneal rinsing. In another embodiment, the sample comprises cells obtained from the patient. In this embodiment, the cells may be found in

30 a fluid selected from the group consisting of a fluid collected by peritoneal rinsing, a fluid collected by uterine rinsing, a uterine fluid, a uterine exudate, a pleural fluid, and an ovarian exudate. In a further embodiment, the patient sample is *in vivo*.

- 10 -

According to the invention, the level of expression of a marker of the invention in a sample can be assessed, for example, by detecting the presence in the sample of:

- 5 • the corresponding marker protein or a fragment of the protein (*e.g.* by using a reagent, such as an antibody, an antibody derivative, an antibody fragment or single-chain antibody, which binds specifically with the protein or protein fragment).
- 10 • the corresponding marker nucleic acid or a fragment of the nucleic acid (*e.g.* by contacting transcribed polynucleotides obtained from the sample with a substrate having affixed thereto one or more nucleic acids having the entire or a segment of the sequence or a complement thereof)
- a metabolite which is produced directly (*i.e.*, catalyzed) or indirectly by the corresponding marker protein.

According to the invention, any of the aforementioned methods may be performed using a plurality (*e.g.* 2, 3, 5, or 10 or more) of ovarian cancer markers, including ovarian cancer markers known in the art. In such methods, the level of expression in the sample of each of a plurality of markers, at least one of which is a marker of the invention, is compared with the normal level of expression of each of the plurality of markers in samples of the same type obtained from control humans not afflicted with ovarian cancer. A significantly altered (*i.e.*, increased or decreased as specified in the above-described methods using a single marker) level of expression in the sample of one or more markers of the invention, or some combination thereof, relative to that marker's corresponding normal levels, is an indication that the patient is afflicted with ovarian cancer. For all of the aforementioned methods, the marker(s) are preferably selected such that the positive predictive value of the method is at least about 10%.

In a further aspect, the invention provides an antibody, an antibody derivative, or an antibody fragment, which binds specifically with a marker protein or a fragment of the protein. The invention also provides methods for making such antibody, antibody derivative, and antibody fragment. Such methods may comprise immunizing a mammal with a protein or peptide comprising the entirety, or a segment of 10 amino acids or more, of a marker protein, wherein the protein or peptide may be obtained from

- 11 -

a cell or by chemical synthesis. The methods of the invention also encompass producing monoclonal and single-chain antibodies, which would further comprise isolating splenocytes from the immunized mammal, fusing the isolated splenocytes with an immortalized cell line to form hybridomas, and screening individual hybridomas for those that produce an antibody that binds specifically with a marker protein or a fragment of the protein.

In another aspect, the invention relates to various diagnostic and test kits. In one embodiment, the invention provides a kit for assessing whether a patient is afflicted with ovarian cancer. The kit comprises a reagent for assessing expression of a marker of the invention. In another embodiment, the invention provides a kit for assessing the suitability of a chemical or biologic agent for inhibiting an ovarian cancer in a patient. Such kit comprises a reagent for assessing expression of a marker of the invention, and may also comprise one or more of such agents. In a further embodiment, the invention provides kits for assessing the presence of ovarian cancer cells or treating ovarian cancers. Such kits comprise an antibody, an antibody derivative, or an antibody fragment, which binds specifically with a marker protein, or a fragment of the protein. Such kits may also comprise a plurality of antibodies, antibody derivatives, or antibody fragments wherein the plurality of such antibody agents binds specifically with a marker protein, or a fragment of the protein.

In an additional embodiment, the invention also provides a kit for assessing the presence of ovarian cancer cells, wherein the kit comprises a nucleic acid probe that binds specifically with a marker nucleic acid or a fragment of the nucleic acid. The kit may also comprise a plurality of probes, wherein each of the probes binds specifically with a marker nucleic acid, or a fragment of the nucleic acid.

In a further aspect, the invention relates to methods for treating a patient afflicted with ovarian cancer or at risk of developing ovarian cancer. Such methods may comprise reducing the expression and/or interfering with the biological function of a marker of the invention. In one embodiment, the method comprises providing to the patient an antisense oligonucleotide or polynucleotide complementary to a marker nucleic acid, or a segment thereof. For example, an antisense polynucleotide may be provided to the patient through the delivery of a vector that expresses an antisense polynucleotide of a marker nucleic acid or a fragment thereof. In another embodiment,

- 12 -

the method comprises providing to the patient an antibody, an antibody derivative, or antibody fragment, which binds specifically with a marker protein or a fragment of the protein. In a preferred embodiment, the antibody, antibody derivative or antibody fragment binds specifically with a protein having the sequence of any of the markers
5 listed in Table 1, or a fragment of such a protein.

It will be appreciated that the methods and kits of the present invention may also include known cancer markers including known ovarian cancer markers. It will further be appreciated that the methods and kits may be used to identify cancers other than ovarian cancer.

10

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 depicts a graph which represents the results of the TaqMan® expression study.

15

DETAILED DESCRIPTION OF THE INVENTION

The invention relates to newly discovered markers, identified in Tables 1-3, that are associated with the cancerous state of ovarian cells. It has been discovered that the higher than normal level of expression of any of these markers or combination of these markers correlates with the presence of ovarian cancer in a patient. Methods
20 are provided for detecting the presence of ovarian cancer in a sample, the absence of ovarian cancer in a sample, the stage of an ovarian cancer, and with other characteristics of ovarian cancer that are relevant to prevention, diagnosis, characterization, and therapy of ovarian cancer in a patient. Methods of treating ovarian cancer are also provided.

Tables 1-3 list the markers of the present invention. In the Tables the
25 markers are identified with a name ("Marker"), the name the gene is commonly known by, if applicable ("Gene Name"), the Sequence Listing identifier of the cDNA sequence of a nucleotide transcript encoded by or corresponding to the marker ("SEQ ID NO (nts)"), the Sequence Listing identifier of the amino acid sequence of a protein encoded by the nucleotide transcript ("SEQ ID NO (AAs)"), and the location of the protein
30 coding sequence within the cDNA sequence ("CDS").

Table 1 lists all of the markers of the invention, which are over-expressed in ovarian cancer cells compared to normal (*i.e.*, non-cancerous) ovarian cells and comprises markers listed in Tables 2 and 3. Table 2 lists newly-identified nucleotide

and amino acid sequences useful as ovarian cancer markers. Table 3 lists newly-identified nucleotide sequences useful as ovarian cancer markers.

In addition to their use in ovarian cancer, it has been found that the markers of the present invention may be used in the diagnosis, characterization, management, and therapy of additional diseases. For example, OV65 (SEQ ID NOS: 305 and 306), M593 (SEQ ID NOS: 307 and 308) and M594 (SEQ ID NOS: 309 and 310), are spondin molecules, and have one or more of the following activities: (1) neural cell adhesion and (2) neurite extension and can thus be used in, for example, the diagnosis and treatment of brain and CNS related disorders. Such brain and CNS related disorders include, but are not limited to, bacterial and viral meningitis, Alzheimers Disease, cerebral toxoplasmosis, Parkinson's disease, multiple sclerosis, brain cancers (*e.g.*, metastatic carcinoma of the brain, glioblastoma, lymphoma, astrocytoma, acoustic neuroma), hydrocephalus, and encephalitis. In another example, OV65, M593 and M594 polypeptides, nucleic acids, and modulators thereof can be used to treat disorders of the brain, such as cerebral edema, hydrocephalus, brain herniations, iatrogenic disease (due to, *e.g.*, infection, toxins, or drugs), inflammations (*e.g.*, bacterial and viral meningitis, encephalitis, and cerebral toxoplasmosis), cerebrovascular diseases (*e.g.*, hypoxia, ischemia, infarction, intracranial hemorrhage, vascular malformations, and hypertensive encephalopathy), and tumors (*e.g.*, neuroglial tumors, neuronal tumors, tumors of pineal cells, meningeal tumors, primary and secondary lymphomas, intracranial tumors, and medulloblastoma), and to treat injury or trauma to the brain.

OV25 (SEQ ID NOS: 360 and 361), an HE4 protein, has one or more of the following activities: (1) sperm maturation and (2) inhibition of extracellular proteases and can thus be used in, for example, the treatment and diagnosis of diseases and disorders relating to spermatogenesis. For example, OV25 polypeptides, nucleic acids, and modulators thereof can be used to treat testicular disorders, such as unilateral testicular enlargement (*e.g.*, nontuberculous, granulomatous orchitis); inflammatory diseases resulting in testicular dysfunction (*e.g.*, gonorrhea and mumps); cryptorchidism; sperm cell disorders (*e.g.*, immotile cilia syndrome and germinal cell aplasia); acquired testicular defects (*e.g.*, viral orchitis); and tumors (*e.g.*, germ cell tumors, interstitial cell tumors, androblastoma, testicular lymphoma and adenomatoid tumors).

OV52 (SEQ ID NOS: 190 and 191), a Pump-1 proteinase, has been found to have one or more of the following activities: (1) breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and remodeling, as well as in (2) disease processes, such as arthritis, and metastasis. Hence, 5 OV52 nucleic acids, proteins, and modulators thereof can be used to modulate disorders associated with adhesion and migration of cells, *e.g.*, platelet aggregation disorders (*e.g.*, Glanzmann's thromboasthenia, which is a bleeding disorder characterized by failure of platelet aggregation in response to cell stimuli), inflammatory disorders (*e.g.*, leukocyte adhesion deficiency, which is a disorder associated with impaired migration of 10 neutrophils to sites of extravascular inflammation), connective tissue disorders, arthritis, disorders associated with abnormal tissue migration during embryo development, and tumor metastasis.

M604 (SEQ ID NOS: 48 and 49), OV10 (SEQ ID NOS: 50 and 51), and M360 (SEQ ID NOS: 52 and 53), are Claudin molecules which have one or more of the 15 following activities: (1) it elicits fluid accumulation in the intestinal tract by altering the membrane permeability of intestinal epithelial cells and (2) thus acts as the causative agent of diarrhea. The polypeptides, nucleic acids, and modulators thereof can be used to treat colonic disorders, such as congenital anomalies (*e.g.*, megacolon and imperforate anus), idiopathic disorders (*e.g.*, diverticular disease and melanosis coli), vascular 20 lesions (*e.g.*, ischemic colitis, hemorrhoids, angiodysplasia), inflammatory diseases (*e.g.*, colitis (*e.g.*, idiopathic ulcerative colitis, pseudomembranous colitis), and lymphopathia venereum), Crohn's disease, and tumors (*e.g.*, hyperplastic polyps, adenomatous polyps, bronchogenic cancer, colonic carcinoma, squamous cell carcinoma, adenoacanthomas, sarcomas, lymphomas, argentaffinomas, carcinoids, and 25 melanocarcinomas).

OV48 (SEQ ID NOS: 226 and 227), OV49 (SEQ ID NOS: 228 and 229) and OV50 (SEQ ID NOS: 230 and 231), markers for an osteopontin protein, have one or more of the following activities: (1) they act as a vessel extracellular matrix protein involved in calcification and (2) atherosclerosis. Hence, OV48, OV49 and OV50 30 nucleic acids, proteins, and modulators thereof can be used to treat heart disorders, *e.g.*, ischemic heart disease, atherosclerosis, hypertension, angina pectoris, Hypertrophic Cardiomyopathy, and congenital heart disease. They can also be used to treat

cardiovascular disorders, such as ischemic heart disease (*e.g.*, angina pectoris, myocardial infarction, and chronic ischemic heart disease), hypertensive heart disease, pulmonary heart disease, valvular heart disease (*e.g.*, rheumatic fever and rheumatic heart disease, endocarditis, mitral valve prolapse, and aortic valve stenosis), congenital heart disease (*e.g.*, valvular and vascular obstructive lesions, atrial or ventricular septal defect, and patent ductus arteriosus), or myocardial disease (*e.g.*, myocarditis, congestive cardiomyopathy, and hypertrophic cardiomyopathy).

OV37 (SEQ ID NOS: 176 and 177), a lipocalin marker, is known to be a component of the neutrophil gelatinase complex. OV37 nucleic acids, proteins, and modulators thereof can be used to modulate the proliferation, differentiation, and/or function of leukocytes. Thus, OV37 nucleic acids, proteins, and modulators thereof can be used to treat bone marrow, blood, and hematopoietic associated diseases and disorders, *e.g.*, acute myeloid leukemia, hemophilia, leukemia, anemia (*e.g.*, sickle cell anemia), and thalassemia. OV37 polypeptides, nucleic acids, and modulators thereof can be used to treat leukocytic disorders, such as leukopenias (*e.g.*, neutropenia, monocytopenia, lymphopenia, and granulocytopenia), leukocytosis (*e.g.*, granulocytosis, lymphocytosis, eosinophilia, monocytosis, acute and chronic lymphadenitis), malignant lymphomas (*e.g.*, Non-Hodgkin's lymphomas, Hodgkin's lymphomas, leukemias, agnogenic myeloid metaplasia, multiple myeloma, plasmacytoma, Waldenstrom's macroglobulinemia, heavy-chain disease, monoclonal gammopathy, histiocytoses, eosinophilic granuloma, and angioimmunoblastic lymphadenopathy).

OV2 (SEQ ID NOS: 285 and 286), is known to be a protease inhibitor, which is associated with emphysema and liver disease. Hence OV2 polypeptides, nucleic acids, and modulators thereof can be used to diagnose and treat pulmonary (lung) disorders, such as atelectasis, cystic fibrosis, rheumatoid lung disease, pulmonary congestion or edema, chronic obstructive airway disease (*e.g.*, emphysema, chronic bronchitis, bronchial asthma, and bronchiectasis), diffuse interstitial diseases (*e.g.*, sarcoidosis, pneumoconiosis, hypersensitivity pneumonitis, bronchiolitis, Goodpasture's syndrome, idiopathic pulmonary fibrosis, idiopathic pulmonary hemosiderosis, pulmonary alveolar proteinosis, desquamative interstitial pneumonitis, chronic interstitial pneumonia, fibrosing alveolitis, hamman-rich syndrome, pulmonary eosinophilia, diffuse interstitial fibrosis, Wegener's granulomatosis, lymphomatoid

granulomatosis, and lipid pneumonia), or tumors (*e.g.*, bronchogenic carcinoma, bronchioloalveolar carcinoma, bronchial carcinoid, hamartoma, and mesenchymal tumors). In another example, OV2 polypeptides, nucleic acids, and modulators thereof can be used to diagnose and treat hepatic (liver) disorders, such as jaundice, hepatic failure, hereditary hyperbilirubinemias (*e.g.*, Gilbert's syndrome, Crigler-Najjar syndromes and Dubin-Johnson and Rotor's syndromes), hepatic circulatory disorders (*e.g.*, hepatic vein thrombosis and portal vein obstruction and thrombosis), hepatitis (*e.g.*, chronic active hepatitis, acute viral hepatitis, and toxic and drug-induced hepatitis), cirrhosis (*e.g.*, alcoholic cirrhosis, biliary cirrhosis, and hemochromatosis), or malignant tumors (*e.g.*, primary carcinoma, hepatoma, hepatoblastoma, liver cysts, and angiosarcoma).

OV32 (SEQ ID NOS: 166 and 167) and OV33 (SEQ ID NOS: 168 and 169), kallikrein markers, are useful in detection of primary mammary carcinomas, as well as primary ovarian cancers. Hence, OV32 and OV33 polypeptides, nucleic acids, and modulators thereof can be used to diagnose and treat ovarian disorders, such as ovarian endometriosis, non-neoplastic cysts (*e.g.*, follicular and luteal cysts and polycystic ovaries) and tumors (*e.g.*, carcinomas, tumors of surface epithelium, germ cell tumors, ovarian fibroma, sex cord-stromal tumors, and ovarian cancers (*e.g.*, metastatic carcinomas, and ovarian teratoma)).

OV68 (SEQ ID NOS: 192 and 193), OV69 (SEQ ID NOS: 194 and 195), OV70 (SEQ ID NOS: 196 and 197), OV71 (SEQ ID NOS: 198 and 199), OV72 (SEQ ID NOS: 200 and 201), OV41 (SEQ ID NOS: 202 and 203), OV42 (SEQ ID NOS: 204 and 205), OV43 (SEQ ID NOS: 206 and 205), OV44 (SEQ ID NOS: 207 and 208) and OV83 (SEQ ID NOS: 209 and 210), are all mesothelin markers, and have been found to play a role in cellular adhesion. The nucleic acids, proteins, and modulators thereof can be used to diagnose, treat and modulate disorders associated with adhesion and migration of cells, *e.g.*, platelet aggregation disorders (*e.g.*, Glanzmann's thrombasthenia, which is a bleeding disorder characterized by failure of platelet aggregation in response to cell stimuli), inflammatory disorders (*e.g.*, leukocyte adhesion deficiency, which is a disorder associated with impaired migration of neutrophils to sites of extravascular inflammation), disorders associated with abnormal tissue migration during embryo development, and tumor metastasis.

- 17 -

OV17 (SEQ ID NOS: 110 and 111), OV18 (SEQ ID NOS: 112 and 111), OV19 (SEQ ID NOS: 113 and 111), OV20 (SEQ ID NOS: 114 and 111), OV21 (SEQ ID NOS: 115 and 111) and OV22 (SEQ ID NOS: 116 and 117) are folate receptors, which are known to be markers of ovarian cancer. The nucleic acids, proteins, and modulators thereof can be used to diagnose, treat and modulate ovarian disorders (*e.g.*, ovarian cyst, ovarian fibroma, ovarian endometriosis, ovarian teratoma). Although these markers have been previously associated with ovarian cancer, the expression of such markers has not yet been identified in combination with the expression of other markers including those of the present invention. Such combination of markers will provide improved methods of diagnosing, characterizing, managing and treating ovarian cancer.

OV66 (SEQ ID NOS: 54 and 55), OV7 (SEQ ID NOS: 56 and 57), OV8 (SEQ ID NOS: 58 and 59) and OV81 (SEQ ID NOS: 60 and 61) are ceruloplasmin markers, known to encode a plasma metalloprotein that binds copper in the plasma. The nucleic acids, proteins, and modulators thereof can be used to diagnose, treat and modulate disorders in blood haemostasis and diseases caused by such an imbalance *e.g.*, (1) cardiovascular diseases or disorders, such as ischemic heart disease (*e.g.*, angina pectoris, myocardial infarction, and chronic ischemic heart disease), hypertensive heart disease, pulmonary heart disease, valvular heart disease (*e.g.*, rheumatic fever and rheumatic heart disease, endocarditis, mitral valve prolapse, and aortic valve stenosis), congenital heart disease (*e.g.*, valvular and vascular obstructive lesions, atrial or ventricular septal defect, and patent ductus arteriosus), or myocardial disease (*e.g.*, myocarditis, congestive cardiomyopathy, and hypertrophic cardiomyopathy); (2) neuronal diseases such as Alzheimers Disease, cerebral toxoplasmosis, Parkinson's disease, multiple sclerosis, brain cancers (*e.g.*, metastatic carcinoma of the brain, glioblastoma, lymphoma, astrocytoma, acoustic neuroma), hydrocephalus, and encephalitis; and (3) Wilson's Disease.

TABLE 1

Marker	Gene Name	SEQ ID NO (nts)	SEQ ID NO (AAs)	CDS
OV1	ABCB1: ATP-binding cassette, sub-family B (MDR/TAP), member 1	1	2	425..4264
M430	ADPRT: ADP-ribosyltransferase	3	4	160..3204
M571	ANXA2: annexin A2, variant 1	5	6	134..1153
M572	ANXA2: annexin A2, variant 2	7	8	50..1069
M573	ANXA4: annexin A4	9	10	74..1039
OV3	AQP5: aquaporin 5	11	12	619..1316
M352	ARHGAP8: Rho GTPase activating protein 8, variant 1	13	14	142..1536
M353	ARHGAP8: Rho GTPase activating protein 8, variant 2	15	16	1..2043
M354	ARHGAP8: Rho GTPase activating protein 8, variant 3	17	18	1..2256
M608	ARHGAP8: Rho GTPase activating protein 8, variant 4	17	19	1..2157
M355	ARHGAP8: Rho GTPase activating protein 8, variant 5	20	21	<1..1314
M356	ARHGAP8: Rho GTPase activating protein 8, variant 6	22	23	1..1902
M357	ARHGAP8: Rho GTPase activating protein 8, variant 7	24	25	<1..1281
M358	ARHGAP8: Rho GTPase activating protein 8, variant 8	26	27	1..1386
M359	ARHGAP8: Rho GTPase activating protein 8, variant 9	28	29	<1..1059
OV5	BICD1: Bicaudal D homolog 1 (Drosophila)	30	31	82..3009
M431	BTG2: BTG family, member 2	32	33	72..548
M432	CADPS: Ca2+-dependent activator protein for secretion	34	35	240..4412
M609	CDH1: cadherin 1, type 1, E-cadherin (epithelial)	36	37	125..2773
M433	CDH6: cadherin 6, type 2, K-cadherin	38	39	327..2699
M434	CDKN2A: cyclin-dependent kinase inhibitor 2A	40	41	41..511
OV9	CGN: cingulin	42	43	152..3763
OV6	CHI3L1: cartilage glycoprotein-39	44	45	127..1278
M435	CKMT1: creatine kinase, mitochondrial 1 (ubiquitous)	46	47	164..1417
M604	CLDN10: claudin 10	48	49	36..772
OV10	CLDN16: claudin 16	50	51	69..986
M360	CLDN4: claudin 4	52	53	183..812
OV66	CP: ceruloplasmin (ferroxidase), variant 1	54	55	1..3210
OV7	CP: ceruloplasmin (ferroxidase), variant 2	56	57	<1..2561
OV8	CP: ceruloplasmin (ferroxidase), variant 3	58	59	1..3198
OV81	CP: ceruloplasmin (ferroxidase), variant 4	60	61	76..3348
M103	CRABP2: cellular retinoic acid-binding protein 2	62	63	138..554

OV40	DD96: Epithelial protein up-regulated in carcinoma, membrane associated protein 17	64	65	202..546
OV4	DEC2: basic helix-loop-helix protein	66	67	135..1583
M575	dehydrogenase	68	69	339..1364
M436	DLX5: distal-less homeo box 5	70	71	204..1073
OV12	EAB1: Eab1 protein	72	73	<1..1305
OV13	ESX protein	74	75	96..1211
OV67	EVI-1: Evi-1 protein, variant 1	76	77	250..2406
OV14	EVI-1: Evi-1 protein, variant 2	78	79	250..3405
OV15	EVI-1: Evi-1 protein, variant 3	80	81	250..2433
OV16	EVI-1: Evi-1 protein, variant 4	82	83	250..3378
M437	FLJ10546: hypothetical protein FLJ10546	84	85	28..1815
OV28	FLJ12799: hypothetical protein FLJ12799	86	87	39..797
M576	FLJ13710: hypothetical protein FLJ13710	88	89	96..1712
M438	FLJ13782: hypothetical protein FLJ13782	90	91	13..1890
OV29	FLJ20150: hypothetical protein FLJ20150	92	93	78..983
M439	FLJ20327: hypothetical protein FLJ20327	94	95	306..2186
M440	FLJ20758: hypothetical protein FLJ20758, variant 1	96	97	<2..1270
M441	FLJ20758: hypothetical protein FLJ20758, variant 2	98	99	<2..2095
M442	FLJ20758: hypothetical protein FLJ20758, variant 3	100	101	465..1307
M443	FLJ2252: likely ortholog of mouse SRY-box containing gene 17	102	103	205..1449
M444	FLJ22316: hypothetical protein FLJ22316	104	105	508..1206
M400	FLJ22418: hypothetical protein FLJ22418	106	107	71..919
M445	FLJ23499: hypothetical protein FLJ23499	108	109	21..473
OV17	FOLR1: folate receptor 1 (alpha), variant 1	110	111	139..912
OV18	FOLR1: folate receptor 1 (alpha), variant 2	112	111	211..984
OV19	FOLR1: folate receptor 1 (alpha), variant 3	113	111	46..819
OV20	FOLR1: folate receptor 1 (alpha), variant 4	114	111	437..1210
OV21	FOLR1: folate receptor 1 (alpha), variant 5	115	111	11..784
OV22	FOLR3: folate receptor 3 (gamma)	116	117	57..788
OV23	GPR39: G protein-coupled receptor 39	118	119	1..1352
M446	GPRC5B: G protein-coupled receptor, family C, group 5, member B	120	121	109..1320
OV24	G-protein coupled receptor	122	123	274..1236
M447	GRB7: growth factor receptor-bound protein 7	124	125	220..1818
OV11	HAIK1: type I intermediate filament cytokeratin	126	127	61..1329
M448	HOXB7: homeo box B7	128	129	100..753
M138	HSECP1: secretory protein, variant 1	130	131	27..863
M449	HSECP1: secretory protein, variant 2	132	133	136..768
M450	HSECP1: secretory protein, variant 3	134	135	202..933
M451	HSNFRK: HSNFRK protein	136	137	642..2939
OV26	hypothetical protein (1)	138	139	<1..1140
OV27	hypothetical protein (2)	140	141	242..1483
OV31	IFI30: interferon, gamma-inducible protein 30	142	143	41..952
OV58	IGF2: somatomedin A	144	145	553..1095

M452	IMP-2: IGF-II mRNA-binding protein 2	146	147	436..2106
M453	INDO: indoleamine-pyrrole 2, 3 dioxygenase	148	149	23..1234
OV73	IPT: tRNA isopentenylpyrophosphate transferase, variant 1	150	151	15..1418
M610	IPT: tRNA isopentenylpyrophosphate transferase, variant 2	152	153	15..1418
M454	ITGA3: Integrin, alpha 3	154	155	74..3274
OV30	ITGB8: Integrin, beta 8	156	157	681..2990
OV34	KIAA0762: KIAA0762 protein	158	159	<1..1875
M455	KIAA0869: KIAA0869 protein	160	161	<1..2668
OV35	KIAA1154: KIAA1154 protein	162	163	<1..677
OV36	KIAA1456: KIAA1456 protein	164	165	<366..1631
OV32	KLK10: kallikrein 10	166	167	82..912
OV33	KLK6: kallikrein 6	168	169	246..980
M456	KRT7: keratin 7, variant 1	170	171	57..1466
M611	KRT7: keratin 7, variant 2	172	173	54..1463
OV53	LC27: Putative integral membrane transporter	174	175	204..1055
OV37	LCN2: Lipocalin 2 (oncogene 24p3)	176	177	1..597
M457	LEFTB: left-right determination, factor B	178	179	71..1171
M559	LPHB: lipophilin B (uteroglobin family member), prostatein-like	180	181	64..336
OV38	LYST-interacting protein LIP6	182	183	11..586
OV39	MEIS1: MEIS1 protein	184	185	66..1238
M458	MGB2: mammaglobin 2	186	187	65..352
M459	MGC3184: similar to sialyltransferase 7 ((alpha-N-acetylneuraminy) 2, 3- betagalactosyl-1, 3)-N-acetyl galactosaminide alpha-2, 6-sialyltransferase) E	188	189	176..1186
OV52	MMP7: Matrix metalloproteinase 7 (matrilysin, uterine)	190	191	28..831
OV68	MSLN: mesothelin, variant 1	192	193	88..2196
OV69	MSLN: mesothelin, variant 2	194	195	88..1980
OV70	MSLN: mesothelin, variant 3	196	197	88..1950
OV71	MSLN: mesothelin, variant 4	198	199	88..2172
OV72	MSLN: mesothelin, variant 5	200	201	88..1926
OV41	MSLN: mesothelin, variant 6	202	203	<1..>1195
OV42	MSLN: mesothelin, variant 7	204	205	85..1953
OV43	MSLN: mesothelin, variant 8	206	205	88..1956
OV44	MSLN: mesothelin, variant 9	207	208	89..1975
OV83	MSLN: mesothelin, variant 10	209	210	295..2187
OV45	MUC1: mucin 1	211	212	58..1605
M460	MUC16: mucin 16, variant 1	213	214	<1..5352
M461	MUC16: mucin 16, variant 2	215	216	25..3471
M612	MUC16: mucin 16, variant 3	215	217	<1..5673
M462	MYO2: myomesin (M-protein)	218	219	49..4446
M463	NaPi-lib: sodium dependent phosphate transporter isoform	220	221	36..2105
M464	NME5: protein expressed in non-metastatic cells 5	222	223	15..653

OV47	NUFIP1: nuclear fragile X mental retardation protein interacting protein 1	224	225	1..1488
OV48	OPN-a: Secreted phosphoprotein-1 (osteopontin, bone sialoprotein)	226	227	1..942
OV49	OPN-b: Secreted phosphoprotein-1 (osteopontin, bone sialoprotein)	228	229	88..990
OV50	OPN-c: Secreted phosphoprotein-1 (osteopontin, bone sialoprotein)	230	231	1..861
M578	PAEP: progesterone-associated endometrial protein, variant 1	232	233	36..578
M579	PAEP: progesterone-associated endometrial protein, variant 2	234	233	36..578
M580	PAEP: progesterone-associated endometrial protein, variant 3	235	233	36..578
M581	PAEP: progesterone-associated endometrial protein, variant 4	236	233	36..578
M583	PAEP: progesterone-associated endometrial protein, variant 5	237	238	45..305
M582	PAEP: progesterone-associated endometrial protein, variant 6	239	240	45..521
M613	PAEP: progesterone-associated endometrial protein, variant 7	239	241	45..521
M465	PAX8: paired box gene 8, isoform 8A	242	243	11..1363
M466	PAX8: paired box gene 8, isoform 8B, variant 1	244	245	11..1174
M614	PAX8: paired box gene 8, isoform 8B, variant 2	244	246	11..1174
M467	PAX8: paired box gene 8, isoform 8C	247	248	161..1357
M468	PAX8: paired box gene 8, isoform 8D	249	250	161..1126
M469	PAX8: paired box gene 8, isoform 8E	251	252	161..1024
M470	PRAME: preferentially expressed antigen in melanoma	253	254	236..1765
M615	PRKCI: protein kinase C, iota	255	256	205..1968
M605	PRP4: serine/threonine-protein kinase PRP4 homolog, variant 1	257	258	<1..3133
M606	PRP4: serine/threonine-protein kinase PRP4 homolog, variant 2	259	258	<1..3133
M607	PRP4: serine/threonine-protein kinase PRP4 homolog, variant 3	260	258	<1..3133
OV80	PRSS8: prostasin	261	262	229..1260
OV51	PTGS1: prostaglandin-endoperoxide synthase 1	263	264	6..1805
M312	PTK9: protein tyrosine kinase 9	265	266	61..1113
OV54	pyruvate dehydrogenase complex component E2	267	268	49..>358
OV55	S100A1: S100 calcium-binding protein A1	269	270	114..398
M471	S100A11: S100 calcium-binding protein A11 (calgizzarin)	271	272	121..438
M68	S100A2: S100 calcium-binding protein A2	273	274	41..334
M585	S100A6: S100 calcium-binding protein A6 (calcylin)	275	276	103..375

OV57	SCNN1A: sodium channel, nonvoltage-gated 1 alpha, variant 1	277	278	100..2109
OV85	SCNN1A: sodium channel, nonvoltage-gated 1 alpha, variant 2	279	280	96..2105
M472	secreted protein (HETKL27)	281	282	88..618
M473	SEMA3A: sema domain, immunoglobulin domain (Ig), short basic domain, secreted, (semaphorin) 3A	283	284	16..2331
OV2	SERPINA1: alpha-1 antitrypsin	285	286	35..1291
M474	Similar to hypothetical protein, MGC: 7199	287	288	173..1053
M586	Similar to proteasome (prosome, macropain) subunit, alpha type, 3	289	290	45..791
M587	Similar to zinc finger protein 136	291	292	139..1524
M475	SLPI: secretory leukocyte protease inhibitor (antileukoproteinase), variant 1	293	294	271..447
M185	SLPI: secretory leukocyte protease inhibitor (antileukoproteinase), variant 2	295	296	19..417
OV60	SNCG: synuclein, gamma	297	298	49..432
OV59	SORL1: sortilin-related receptor	299	300	198..6842
OV56	SPINT2: serine protease inhibitor, Kunitz type, 2, variant 1	301	302	301..1059
OV84	SPINT2: serine protease inhibitor, Kunitz type, 2, variant 2	303	304	332..919
OV65	SPON1: VSGP/F-spondin, variant 1	305	306	25..2448
M593	SPON1: VSGP/F-spondin, variant 2	307	308	180..2984
M594	SPON1: VSGP/F-spondin, variant 3	309	310	180..2687
OV82	ST14: matriptase	311	312	209..2557
M476	TACSTD2: tumor-associated calcium signal transducer 2	313	314	616..1587
M588	TFPI2: tissue factor pathway inhibitor 2	315	316	57..764
OV86	TMPRSS4: transmembrane protease, serine 4	317	318	310..1623
OV74	TPH: tryptophan hydroxylase, variant 1	319	320	1..1335
OV75	TPH: tryptophan hydroxylase, variant 2	321	322	1..1401
M327	TSPAN-1: Tetraspan NET-1 protein, variant 1	323	324	124..900
M328	TSPAN-1: Tetraspan NET-1 protein, variant 2	325	326	1..726
OV46	TTID: myotilin	327	328	281..1777
M589	UCH2: Ubiquitin carboxyl-terminal hydrolases family 2	329	330	551..2940
OV63	unnamed gene (1)	331	332	71..919
OV64	unnamed gene (2)	333	334	28..804
OV76	unnamed gene (3)	335	336	69..773
OV77	unnamed gene (4)	337	338	223..1284
OV78	unnamed gene (5), variant 1	339	340	84..2450
M616	unnamed gene (5), variant 2	341	342	84..2450
OV79	unnamed gene (6)	343	344	69..392
OV87	unnamed gene (7)	345	346	509..2428
OV88	unnamed gene (8)	347	348	71..919
M477	unnamed gene (9), variant 1	349	350	246..992
M617	unnamed gene (9), variant 2	349	351	246..992
M478	unnamed gene (9), variant 3	352	353	246..1004

M479	unnamed gene (9), variant 4	354	355	246..1049
M590	unnamed gene (10), variant 1	356	357	21..404
M591	unnamed gene (10), variant 2	358	357	21..404
M592	unnamed gene (10), variant 3	359	357	21..404
OV25	WFDC2: Epididymis-specific, whey-acidic protein type, four-disulfide core; putative ovarian carcinoma marker	360	361	28..405
M480	XRCC5, KU80: ATP-dependant DNA helicase II	362	363	34..2232

TABLE 2

Marker	Gene Name	SEQ ID NO (nts)	SEQ ID NO (AAs)	CDS
M354	ARHGAP8: Rho GTPase activating protein 8, variant 3	17	18	1..2256
M608	ARHGAP8: Rho GTPase activating protein 8, variant 4	17	19	1..2157
M355	ARHGAP8: Rho GTPase activating protein 8, variant 5	20	21	<1..1314
M356	ARHGAP8: Rho GTPase activating protein 8, variant 6	22	23	1..1902
M357	ARHGAP8: Rho GTPase activating protein 8, variant 7	24	25	<1..1281
M358	ARHGAP8: Rho GTPase activating protein 8, variant 8	26	27	1..1386
M359	ARHGAP8: Rho GTPase activating protein 8, variant 9	28	29	<1..1059
OV68	CP: ceruloplasmin (ferroxidase), variant 1	54	55	1..3210
OV81	CP: ceruloplasmin (ferroxidase), variant 4	60	61	76..3348
M575	dehydrogenase	68	69	339..1364
OV67	EVI-1: Evi-1 protein, variant 1	76	77	250..2406
M440	FLJ20758: hypothetical protein FLJ20758, variant 1	96	97	<2..1270
M441	FLJ20758: hypothetical protein FLJ20758, variant 2	98	99	<2..2095
M449	HSECP1: secretory protein, variant 2	132	133	136..768
M450	HSECP1: secretory protein, variant 3	134	135	202..933
OV73	IPT: tRNA isopentenylpyrophosphate transferase, variant 1	150	151	15..1418
M610	IPT: tRNA isopentenylpyrophosphate transferase, variant 2	152	153	15..1418
M611	KRT7: keratin 7, variant 2	172	173	54..1463
OV68	MSLN: mesothelin, variant 1	192	193	88..2196
OV69	MSLN: mesothelin, variant 2	194	195	88..1980
OV70	MSLN: mesothelin, variant 3	196	197	88..1950
OV71	MSLN: mesothelin, variant 4	198	199	88..2172
OV72	MSLN: mesothelin, variant 5	200	201	88..1926
OV83	MSLN: mesothelin, variant 10	209	210	295..2187
M460	MUC16: mucin 16, variant 1	213	214	<1..5352
M583	PAEP: progesterone-associated endometrial protein, variant 5	237	238	45..305

M613	PAEP: progestagen-associated endometrial protein, variant 7	239	241	45..521
M614	PAX8: paired box gene 8, isoform 8B, variant 2	244	246	11..1174
M605	PRP4: serine/threonine-protein kinase PRP4 homolog, variant 1	257	258	<1..3133
M606	PRP4: serine/threonine-protein kinase PRP4 homolog, variant 2	259	258	<1..3133
M607	PRP4: serine/threonine-protein kinase PRP4 homolog, variant 3	260	258	<1..3133
OV85	SCNN1A: sodium channel, nonvoltage-gated 1 alpha, variant 2	279	280	96..2105
M475	SLPI: secretory leukocyte protease inhibitor (antileukoproteinase), variant 1	293	294	271..447
OV84	SPINT2: serine protease inhibitor, Kunitz type, 2, variant 2	303	304	332..919
M593	SPON1: VSGP/F-spondin, variant 2	307	308	180..2984
M594	SPON1: VSGP/F-spondin, variant 3	309	310	180..2687
OV82	ST14: matriptase	311	312	209..2557
OV86	TMPRSS4: transmembrane protease, serine 4	317	318	310..1623
OV74	TPH: tryptophan hydroxylase, variant 1	319	320	1..1335
OV75	TPH: tryptophan hydroxylase, variant 2	321	322	1..1401
M327	TSPAN-1: Tetraspan NET-1 protein, variant 1	323	324	124..900
M589	UCH2: Ubiquitin carboxyl-terminal hydrolases family 2	329	330	551..2940
OV76	unnamed gene (3)	335	336	69..773
OV77	unnamed gene (4)	337	338	223..1284
OV78	unnamed gene (5), variant 1	339	340	84..2450
M616	unnamed gene (5), variant 2	341	342	84..2450
OV79	unnamed gene (6)	343	344	69..392
OV87	unnamed gene (7)	345	346	509..2428
OV88	unnamed gene (8)	347	348	71..919
M477	unnamed gene (9), variant 1	349	350	246..992
M617	unnamed gene (9), variant 2	349	351	246..992
M478	unnamed gene (9), variant 3	352	353	246..1004
M479	unnamed gene (9), variant 4	354	355	246..1049

TABLE 3

Marker	Gene Name	SEQ ID NO (nts)	SEQ ID NO (AAs)	CDS
M604	CLDN10: claudin 10	48	49	36..772
OV14	EVI-1: Evi-1 protein, variant 2	78	79	250..3405
OV15	EVI-1: Evi-1 protein, variant 3	80	81	250..2433
OV16	EVI-1: Evi-1 protein, variant 4	82	83	250..3378
M576	FLJ13710: hypothetical protein FLJ13710	88	89	96..1712
M444	FLJ22316: hypothetical protein FLJ22316	104	105	508..1206
OV30	ITGB8: integrin, beta 8	156	157	681..2990
OV43	MSLN: mesothelin, variant 8	206	205	88..1956

M581	PAEP: progestagen-associated endometrial protein, variant 4	236	233	36..578
M582	PAEP: progestagen-associated endometrial protein, variant 6	239	240	45..521
M466	PAX8: paired box gene 8, isoform 8B, variant 1	244	245	11..1174
M467	PAX8: paired box gene 8, isoform 8C	247	248	161..1357
M468	PAX8: paired box gene 8, isoform 8D	249	250	161..1126
M469	PAX8: paired box gene 8, isoform 8E	251	252	161..1024
OV2	SERPINA1: alpha-1 antitrypsin	285	286	35..1291
M474	Similar to hypothetical protein, MGC: 7199	287	288	173..1053
M590	unnamed gene (10), variant 1	356	357	21..404
M591	unnamed gene (10), variant 2	358	357	21..404
M592	unnamed gene (10), variant 3	359	357	21..404

Definitions

- As used herein, each of the following terms has the meaning associated with it in this section.

The articles "a" and "an" are used herein to refer to one or to more than one (*i.e.* to at least one) of the grammatical object of the article. By way of example, "an element" means one element or more than one element.

- A "marker" is a gene whose altered level of expression in a tissue or cell from its expression level in normal or healthy tissue or cell is associated with a disease state, such as cancer. A "marker nucleic acid" is a nucleic acid (*e.g.*, mRNA, cDNA) encoded by or corresponding to a marker of the invention. Such marker nucleic acids can be DNA (*e.g.*, cDNA) comprising the sequences listed in Table 1 or the complement of such sequences. The marker nucleic acids also can be RNA comprising the sequences listed in Table 1 or the complement of such sequence, wherein all thymidine residues are replaced with uridine residues. A "marker protein" is a protein encoded by or corresponding to a marker of the invention. A marker protein comprises the sequence of any of the sequences listed in Table 1. The terms "protein" and "polypeptide" are used interchangeably.

- The term "probe" refers to any molecule which is capable of selectively binding to a specifically intended target molecule, for example, a nucleotide transcript or protein encoded by or corresponding to a marker. Probes can be either synthesized by one skilled in the art, or derived from appropriate biological preparations. For purposes of detection of the target molecule, probes may be specifically designed to be

labeled, as described herein. Examples of molecules that can be utilized as probes include, but are not limited to, RNA, DNA, proteins, antibodies, and organic molecules.

An "ovary-associated" body fluid is a fluid which, when in the body of a patient, contacts or passes through ovarian cells or into which cells or proteins shed from ovarian cells *e.g.* ovarian epithelium, are capable of passing. Exemplary ovary-associated body fluids include blood fluids, lymph, ascites, gynecological fluids, cystic fluid, urine, and fluids collected by peritoneal rinsing.

The "normal" level of expression of a marker is the level of expression of the marker in ovarian cells of a human subject or patient not afflicted with ovarian cancer

An "over-expression" or "significantly higher level of expression" of a marker refers to an expression level in a test sample that is greater than the standard error of the assay employed to assess expression, and is preferably at least twice, and more preferably three, four, five or ten times the expression level of the marker in a control sample (*e.g.*, sample from a healthy subjects not having the marker associated disease) and preferably, the average expression level of the marker in several control samples.

As used herein, the term "promoter/regulatory sequence" means a nucleic acid sequence which is required for expression of a gene product operably linked to the promoter/regulatory sequence. In some instances, this sequence may be the core promoter sequence and in other instances, this sequence may also include an enhancer sequence and other regulatory elements which are required for expression of the gene product. The promoter/regulatory sequence may, for example, be one which expresses the gene product in a tissue-specific manner.

A "constitutive" promoter is a nucleotide sequence which, when operably linked with a polynucleotide which encodes or specifies a gene product, causes the gene product to be produced in a living human cell under most or all physiological conditions of the cell.

An "inducible" promoter is a nucleotide sequence which, when operably linked with a polynucleotide which encodes or specifies a gene product, causes the gene product to be produced in a living human cell substantially only when an inducer which corresponds to the promoter is present in the cell.

- 27 -

A "tissue-specific" promoter is a nucleotide sequence which, when operably linked with a polynucleotide which encodes or specifies a gene product, causes the gene product to be produced in a living human cell substantially only if the cell is a cell of the tissue type corresponding to the promoter.

- 5 A "transcribed polynucleotide" or "nucleotide transcript" is a polynucleotide (*e.g.* an mRNA, hnRNA, a cDNA, or an analog of such RNA or cDNA) which is complementary to or homologous with all or a portion of a mature mRNA made by transcription of a marker of the invention and normal post-transcriptional processing (*e.g.* splicing), if any, of the RNA transcript, and reverse transcription of the
10 RNA transcript.

- "Complementary" refers to the broad concept of sequence complementarity between regions of two nucleic acid strands or between two regions of the same nucleic acid strand. It is known that an adenine residue of a first nucleic acid region is capable of forming specific hydrogen bonds ("base pairing") with a residue of
15 a second nucleic acid region which is antiparallel to the first region if the residue is thymine or uracil. Similarly, it is known that a cytosine residue of a first nucleic acid strand is capable of base pairing with a residue of a second nucleic acid strand which is antiparallel to the first strand if the residue is guanine. A first region of a nucleic acid is complementary to a second region of the same or a different nucleic acid if, when the
20 two regions are arranged in an antiparallel fashion, at least one nucleotide residue of the first region is capable of base pairing with a residue of the second region. Preferably, the first region comprises a first portion and the second region comprises a second portion, whereby, when the first and second portions are arranged in an antiparallel fashion, at least about 50%, and preferably at least about 75%, at least about 90%, or at
25 least about 95% of the nucleotide residues of the first portion are capable of base pairing with nucleotide residues in the second portion. More preferably, all nucleotide residues of the first portion are capable of base pairing with nucleotide residues in the second portion.

- "Homologous" as used herein, refers to nucleotide sequence similarity
30 between two regions of the same nucleic acid strand or between regions of two different nucleic acid strands. When a nucleotide residue position in both regions is occupied by the same nucleotide residue, then the regions are homologous at that position. A first

- 28 -

region is homologous to a second region if at least one nucleotide residue position of each region is occupied by the same residue. Homology between two regions is expressed in terms of the proportion of nucleotide residue positions of the two regions that are occupied by the same nucleotide residue. By way of example, a region having the nucleotide sequence 5'-ATTGCC-3' and a region having the nucleotide sequence 5'-TATGGC-3' share 50% homology. Preferably, the first region comprises a first portion and the second region comprises a second portion, whereby, at least about 50%, and preferably at least about 75%, at least about 90%, or at least about 95% of the nucleotide residue positions of each of the portions are occupied by the same nucleotide residue. More preferably, all nucleotide residue positions of each of the portions are occupied by the same nucleotide residue.

A molecule is "fixed" or "affixed" to a substrate if it is covalently or non-covalently associated with the substrate such the substrate can be rinsed with a fluid (e.g. standard saline citrate, pH 7.4) without a substantial fraction of the molecule dissociating from the substrate.

As used herein, a "naturally-occurring" nucleic acid molecule refers to an RNA or DNA molecule having a nucleotide sequence that occurs in an organism found in nature.

A cancer is "inhibited" if at least one symptom of the cancer is alleviated, terminated, slowed, or prevented. As used herein, ovarian cancer is also "inhibited" if recurrence or metastasis of the cancer is reduced, slowed, delayed, or prevented.

A kit is any manufacture (e.g. a package or container) comprising at least one reagent, e.g. a probe, for specifically detecting the expression of a marker of the invention. The kit may be promoted, distributed, or sold as a unit for performing the methods of the present invention.

"Proteins of the invention" encompass marker proteins and their fragments; variant marker proteins and their fragments; peptides and polypeptides comprising an at least 15 amino acid segment of a marker or variant marker protein; and fusion proteins comprising a marker or variant marker protein, or an at least 15 amino acid segment of a marker or variant marker protein.

- 29 -

Unless otherwise specified herewithin, the terms “antibody” and “antibodies” broadly encompass naturally-occurring forms of antibodies (e.g., IgG, IgA, IgM, IgE) and recombinant antibodies such as single-chain antibodies, chimeric and humanized antibodies and multi-specific antibodies, as well as fragments and derivatives of all of the foregoing, which fragments and derivatives have at least an antigenic binding site. Antibody derivatives may comprise a protein or chemical moiety conjugated to an antibody moiety.

Description

The present invention is based, in part, on newly identified markers which are over-expressed in ovarian cancer cells as compared to their expression in normal (*i.e.* non-cancerous) ovarian cells. The enhanced expression of one or more of these markers in ovarian cells is herein correlated with the cancerous state of the tissue. The invention provides compositions, kits, and methods for assessing the cancerous state of ovarian cells (*e.g.* cells obtained from a human, cultured human cells, archived or preserved human cells and *in vivo* cells) as well as treating patients afflicted with ovarian cancer.

The compositions, kits, and methods of the invention have the following uses, among others:

- 1) assessing whether a patient is afflicted with ovarian cancer;
- 2) assessing the stage of ovarian cancer in a human patient;
- 3) assessing the grade of ovarian cancer in a patient;
- 4) assessing the benign or malignant nature of ovarian cancer in a patient;
- 5) assessing the metastatic potential of ovarian cancer in a patient;
- 6) assessing the histological type of neoplasm (*e.g.* serous, mucinous, endometrioid, or clear cell neoplasm) associated with ovarian cancer in a patient;
- 7) making antibodies, antibody fragments or antibody derivatives that are useful for treating ovarian cancer and/or assessing whether a patient is afflicted with ovarian cancer;

- 30 -

- 8) assessing the presence of ovarian cancer cells;
- 9) assessing the efficacy of one or more test compounds for inhibiting ovarian cancer in a patient;
- 10) assessing the efficacy of a therapy for inhibiting ovarian cancer in a patient;
- 11) monitoring the progression of ovarian cancer in a patient;
- 12) selecting a composition or therapy for inhibiting ovarian cancer in a patient;
- 13) treating a patient afflicted with ovarian cancer;
- 14) inhibiting ovarian cancer in a patient;
- 15) assessing the ovarian carcinogenic potential of a test compound; and
- 16) preventing the onset of ovarian cancer in a patient at risk for developing ovarian cancer.

The invention thus includes a method of assessing whether a patient is afflicted with ovarian cancer which includes assessing whether the patient has pre-metastasized ovarian cancer. This method comprises comparing the level of expression of a marker of the invention (listed in Table 1) in a patient sample and the normal level of expression of the marker in a control, *e.g.*, a non-ovarian cancer sample. A significantly higher level of expression of the marker in the patient sample as compared to the normal level is an indication that the patient is afflicted with ovarian cancer.

Gene delivery vehicles, host cells and compositions (all described herein) containing nucleic acids comprising the entirety, or a segment of 15 or more nucleotides, of any of the sequences listed in Tables 1-3 or the complement of such sequences, and polypeptides comprising the entirety, or a segment of 10 or more amino acids, of any of the sequences listed in Tables 1-3 are also provided by this invention.

As described herein, ovarian cancer in patients is associated with an increased level of expression of one or more markers of the invention. While, as discussed above, some of these changes in expression level result from occurrence of the ovarian cancer, others of these changes induce, maintain, and promote the cancerous state of ovarian cancer cells. Thus, ovarian cancer characterized by an increase in the level of expression of one or more markers of the invention can be inhibited by reducing

- 31 -

and/or interfering with the expression of the markers and/or function of the proteins encoded by those markers.

Expression of a marker of the invention can be inhibited in a number of ways generally known in the art. For example, an antisense oligonucleotide can be provided to the ovarian cancer cells in order to inhibit transcription, translation, or both, of the marker(s). Alternately, a polynucleotide encoding an antibody, an antibody derivative, or an antibody fragment which specifically binds a marker protein, and operably linked with an appropriate promoter/regulator region, can be provided to the cell in order to generate intracellular antibodies which will inhibit the function or activity of the protein. The expression and/or function of a marker may also be inhibited by treating the ovarian cancer cell with an antibody, antibody derivative or antibody fragment that specifically binds a marker protein. Using the methods described herein, a variety of molecules, particularly including molecules sufficiently small that they are able to cross the cell membrane, can be screened in order to identify molecules which inhibit expression of a marker or inhibit the function of a marker protein. The compound so identified can be provided to the patient in order to inhibit ovarian cancer cells of the patient.

Any marker or combination of markers of the invention, as well as any known markers in combination with the markers of the invention, may be used in the compositions, kits, and methods of the present invention. In general, it is preferable to use markers for which the difference between the level of expression of the marker in ovarian cancer cells and the level of expression of the same marker in normal ovarian cells is as great as possible. Although this difference can be as small as the limit of detection of the method for assessing expression of the marker, it is preferred that the difference be at least greater than the standard error of the assessment method, and preferably a difference of at least 2-, 3-, 4-, 5-, 6-, 7-, 8-, 9-, 10-, 15-, 20-, 25-, 100-, 500-, 1000-fold or greater than the level of expression of the same marker in normal ovarian tissue.

It is recognized that certain marker proteins are secreted from ovarian cells (*i.e.* one or both of normal and cancerous cells) to the extracellular space surrounding the cells. These markers are preferably used in certain embodiments of the compositions, kits, and methods of the invention, owing to the fact that the such marker

- 32 -

proteins can be detected in an ovary-associated body fluid sample, which may be more easily collected from a human patient than a tissue biopsy sample. In addition, preferred *in vivo* techniques for detection of a marker protein include introducing into a subject a labeled antibody directed against the protein. For example, the antibody can be labeled
5 with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques.

It is a simple matter for the skilled artisan to determine whether any particular marker protein is a secreted protein. In order to make this determination, the marker protein is expressed in, for example, a mammalian cell, preferably a human
10 ovarian cell line, extracellular fluid is collected, and the presence or absence of the protein in the extracellular fluid is assessed (*e.g.* using a labeled antibody which binds specifically with the protein).

The following is an example of a method which can be used to detect secretion of a protein. About 8×10^5 293T cells are incubated at 37°C in wells
15 containing growth medium (Dulbecco's modified Eagle's medium {DMEM} supplemented with 10% fetal bovine serum) under a 5% (v/v) CO₂, 95% air atmosphere to about 60-70% confluence. The cells are then transfected using a standard transfection mixture comprising 2 micrograms of DNA comprising an expression vector encoding the protein and 10 microliters of LipofectAMINE™ (GIBCO/BRL Catalog no. 18342-
20 012) per well. The transfection mixture is maintained for about 5 hours, and then replaced with fresh growth medium and maintained in an air atmosphere. Each well is gently rinsed twice with DMEM which does not contain methionine or cysteine (DMEM-MC; ICN Catalog no. 16-424- 54). About 1 milliliter of DMEM-MC and about 50 microcuries of Trans-³⁵S™ reagent (ICN Catalog no. 51006) are added to each
25 well. The wells are maintained under the 5% CO₂ atmosphere described above and incubated at 37°C for a selected period. Following incubation, 150 microliters of conditioned medium is removed and centrifuged to remove floating cells and debris. The presence of the protein in the supernatant is an indication that the protein is secreted.

Examples of ovary-associated body fluids include blood fluids (*e.g.* whole blood, blood serum, blood having platelets removed therefrom, etc.), lymph, ascitic fluids, gynecological fluids (*e.g.* ovarian, fallopian, and uterine secretions, menses, vaginal douching fluids, fluids used to rinse ovarian cell samples, etc.), cystic
5 fluid, urine, and fluids collected by peritoneal rinsing (*e.g.* fluids applied and collected during laparoscopy or fluids instilled into and withdrawn from the peritoneal cavity of a human patient). In these embodiments, the level of expression of the marker can be assessed by assessing the amount (*e.g.* absolute amount or concentration) of the marker protein in an ovary-associated body fluid obtained from a patient. The fluid can, of
10 course, be subjected to a variety of well-known post-collection preparative and storage techniques (*e.g.* storage, freezing, ultrafiltration, concentration, evaporation, centrifugation, etc.) prior to assessing the amount of the marker in the fluid.

Many ovary-associated body fluids (*i.e.* usually excluding urine) can have ovarian cells, *e.g.* ovarian epithelium, therein, particularly when the ovarian cells
15 are cancerous, and, more particularly, when the ovarian cancer is metastasizing. Cell-containing fluids which can contain ovarian cancer cells include, but are not limited to, peritoneal ascites, fluids collected by peritoneal rinsing, fluids collected by uterine rinsing, uterine fluids such as uterine exudate and menses, pleural fluid, and ovarian exudates. Thus, the compositions, kits, and methods of the invention can be used to
20 detect expression of marker proteins having at least one portion which is displayed on the surface of cells which express it. It is a simple matter for the skilled artisan to determine whether a marker protein, or a portion thereof, is exposed on the cell surface. For example, immunological methods may be used to detect such proteins on whole cells, or well known computer-based sequence analysis methods (*e.g.* the SIGNALP
25 program; Nielsen *et al.*, 1997, *Protein Engineering* 10:1-6) may be used to predict the presence of at least one extracellular domain (*i.e.* including both secreted proteins and proteins having at least one cell-surface domain). Expression of a marker protein having at least one portion which is displayed on the surface of a cell which expresses it may be detected without necessarily lysing the cell (*e.g.* using a labeled antibody which binds
30 specifically with a cell-surface domain of the protein).

Expression of a marker of the invention may be assessed by any of a wide variety of well known methods for detecting expression of a transcribed nucleic acid or protein. Non-limiting examples of such methods include immunological methods for detection of secreted, cell-surface, cytoplasmic, or nuclear proteins, protein
5 purification methods, protein function or activity assays, nucleic acid hybridization methods, nucleic acid reverse transcription methods, and nucleic acid amplification methods.

In a preferred embodiment, expression of a marker is assessed using an antibody (*e.g.* a radio-labeled, chromophore-labeled, fluorophore-labeled, or enzyme-labeled antibody), an antibody derivative (*e.g.* an antibody conjugated with a substrate
10 or with the protein or ligand of a protein-ligand pair {*e.g.* biotin-streptavidin}), or an antibody fragment (*e.g.* a single-chain antibody, an isolated antibody hypervariable domain, etc.) or derivative which binds specifically with a marker protein or fragment thereof, including a marker protein which has undergone all or a portion of its normal
15 post-translational modification.

In another preferred embodiment, expression of a marker is assessed by preparing mRNA/cDNA (*i.e.* a transcribed polynucleotide) from cells in a patient sample, and by hybridizing the mRNA/cDNA with a reference polynucleotide which is a complement of a marker nucleic acid, or a fragment thereof. cDNA can, optionally, be
20 amplified using any of a variety of polymerase chain reaction methods prior to hybridization with the reference polynucleotide; preferably, it is not amplified. Expression of one or more markers can likewise be detected using quantitative PCR to assess the level of expression of the marker(s). Alternatively, any of the many known methods of detecting mutations or variants (*e.g.* single nucleotide polymorphisms,
25 deletions, etc.) of a marker of the invention may be used to detect occurrence of a marker in a patient.

In a related embodiment, a mixture of transcribed polynucleotides obtained from the sample is contacted with a substrate having fixed thereto a polynucleotide complementary to or homologous with at least a portion (*e.g.* at least 7,
30 10, 15, 20, 25, 30, 40, 50, 100, 500, or more nucleotide residues) of a marker nucleic acid. If polynucleotides complementary to or homologous with several marker nucleic acids are differentially detectable on the substrate (*e.g.* detectable using different

chromophores or fluorophores, or fixed to different selected positions), then the levels of expression of a plurality of markers can be assessed simultaneously using a single substrate (e.g. a "gene chip" microarray of polynucleotides fixed at selected positions). When a method of assessing marker expression is used which involves hybridization of one nucleic acid with another, it is preferred that the hybridization be performed under stringent hybridization conditions.

Because the compositions, kits, and methods of the invention rely on detection of a difference in expression levels of one or more markers of the invention, it is preferable that the level of expression of the marker is significantly greater than the minimum detection limit of the method used to assess expression in at least one of normal ovarian cells and cancerous ovarian cells.

It is understood that by routine screening of additional patient samples using one or more of the markers of the invention, it will be realized that certain of the markers are over-expressed in cancers of various types, including specific ovarian cancers, as well as other cancers such as breast cancer, cervical cancer, etc. For example, it will be confirmed that some of the markers of the invention are over-expressed in most (i.e. 50% or more) or substantially all (i.e. 80% or more) of ovarian cancer. Furthermore, it will be confirmed that certain of the markers of the invention are associated with ovarian cancer of various stages (i.e. stage I, II, III, and IV ovarian cancers, as well as subclassifications IA, IB, IC, IIA, IIB, IIC, IIIA, IIIB, and IIIC, using the FIGO Stage Grouping system for primary carcinoma of the ovary; 1987, *Am. J. Obstet. Gynecol.* 156:236), of various histologic subtypes (e.g. serous, mucinous, endometrioid, and clear cell subtypes, as well as subclassifications and alternate classifications adenocarcinoma, papillary adenocarcinoma, papillary cystadenocarcinoma, surface papillary carcinoma, malignant adenofibroma, cystadenofibroma, adenocarcinoma, cystadenocarcinoma, adenoacanthoma, endometrioid stromal sarcoma, mesodermal (Müllerian) mixed tumor, mesonephroid tumor, malignant carcinoma, Brenner tumor, mixed epithelial tumor, and undifferentiated carcinoma, using the WHO/FIGO system for classification of malignant ovarian tumors; Scully, *Atlas of Tumor Pathology*, 3d series, Washington DC), and various grades (i.e. grade I {well differentiated}, grade II {moderately well differentiated}, and grade III {poorly differentiated from surrounding normal tissue}).

- 36 -

In addition, as a greater number of patient samples are assessed for expression of the markers of the invention and the outcomes of the individual patients from whom the samples were obtained are correlated, it will also be confirmed that increased expression of certain of the markers of the invention are strongly correlated with malignant cancers and that increased expression of other markers of the invention are strongly correlated with benign tumors. The compositions, kits, and methods of the invention are thus useful for characterizing one or more of the stage, grade, histological type, and benign/malignant nature of ovarian cancer in patients. In addition, these compositions, kits, and methods can be used to detect and differentiate epithelial, stromal, and germ cell ovarian cancers.

When the compositions, kits, and methods of the invention are used for characterizing one or more of the stage, grade, histological type, and benign/malignant nature of ovarian cancer in a patient, it is preferred that the marker or panel of markers of the invention is selected such that a positive result is obtained in at least about 20%, and preferably at least about 40%, 60%, or 80%, and more preferably in substantially all patients afflicted with an ovarian cancer of the corresponding stage, grade, histological type, or benign/malignant nature. Preferably, the marker or panel of markers of the invention is selected such that a PPV of greater than about 10% is obtained for the general population (more preferably coupled with an assay specificity greater than 99.5%).

When a plurality of markers of the invention are used in the compositions, kits, and methods of the invention, the level of expression of each marker in a patient sample can be compared with the normal level of expression of each of the plurality of markers in non-cancerous samples of the same type, either in a single reaction mixture (*i.e.* using reagents, such as different fluorescent probes, for each marker) or in individual reaction mixtures corresponding to one or more of the markers. In one embodiment, a significantly increased level of expression of more than one of the plurality of markers in the sample, relative to the corresponding normal levels, is an indication that the patient is afflicted with ovarian cancer. When a plurality of markers is used, it is preferred that 2, 3, 4, 5, 8, 10, 12, 15, 20, 30, or 50 or more individual markers be used, wherein fewer markers are preferred.

In order to maximize the sensitivity of the compositions, kits, and methods of the invention (*i.e.* by interference attributable to cells of non-ovarian origin in a patient sample), it is preferable that the marker of the invention used therein be a marker which has a restricted tissue distribution, *e.g.*, normally not expressed in a non-epithelial tissue, and more preferably a marker which is normally not expressed in a non-ovarian tissue.

Only a small number of markers are known to be associated with ovarian cancers (*e.g.* *AKT2*, *Ki-RAS*, *ERBB2*, *c-MYC*, *RB1*, and *TP53*; Lynch, *supra*). These markers are not, of course, included among the markers of the invention, although they may be used together with one or more markers of the invention in a panel of markers, for example. It is well known that certain types of genes, such as oncogenes, tumor suppressor genes, growth factor-like genes, protease-like genes, and protein kinase-like genes are often involved with development of cancers of various types. Thus, among the markers of the invention, use of those which correspond to proteins which resemble proteins encoded by known oncogenes and tumor suppressor genes, and those which correspond to proteins which resemble growth factors, proteases, and protein kinases are preferred.

It is recognized that the compositions, kits, and methods of the invention will be of particular utility to patients having an enhanced risk of developing ovarian cancer and their medical advisors. Patients recognized as having an enhanced risk of developing ovarian cancer include, for example, patients having a familial history of ovarian cancer, patients identified as having a mutant oncogene (*i.e.* at least one allele), and patients of advancing age (*i.e.* women older than about 50 or 60 years).

The level of expression of a marker in normal (*i.e.* non-cancerous) human ovarian tissue can be assessed in a variety of ways. In one embodiment, this normal level of expression is assessed by assessing the level of expression of the marker in a portion of ovarian cells which appears to be non-cancerous and by comparing this normal level of expression with the level of expression in a portion of the ovarian cells which is suspected of being cancerous. For example, when laparoscopy or other medical procedure, reveals the presence of a lump on one portion of a patient's ovary, but not on another portion of the same ovary or on the other ovary, the normal level of expression of a marker may be assessed using one or both or the non-affected ovary and

- 38 -

a non-affected portion of the affected ovary, and this normal level of expression may be compared with the level of expression of the same marker in an affected portion (*i.e.* the lump) of the affected ovary. Alternately, and particularly as further information becomes available as a result of routine performance of the methods described herein, population-average values for normal expression of the markers of the invention may be used. In other embodiments, the 'normal' level of expression of a marker may be determined by assessing expression of the marker in a patient sample obtained from a non-cancer-afflicted patient, from a patient sample obtained from a patient before the suspected onset of ovarian cancer in the patient, from archived patient samples, and the like.

The invention includes compositions, kits, and methods for assessing the presence of ovarian cancer cells in a sample (*e.g.* an archived tissue sample or a sample obtained from a patient). These compositions, kits, and methods are substantially the same as those described above, except that, where necessary, the compositions, kits, and methods are adapted for use with samples other than patient samples. For example, when the sample to be used is a paraffinized, archived human tissue sample, it can be necessary to adjust the ratio of compounds in the compositions of the invention, in the kits of the invention, or the methods used to assess levels of marker expression in the sample. Such methods are well known in the art and within the skill of the ordinary artisan.

The invention includes a kit for assessing the presence of ovarian cancer cells (*e.g.* in a sample such as a patient sample). The kit comprises a plurality of reagents, each of which is capable of binding specifically with a marker nucleic acid or protein. Suitable reagents for binding with a marker protein include antibodies, antibody derivatives, antibody fragments, and the like. Suitable reagents for binding with a marker nucleic acid (*e.g.* a genomic DNA, an mRNA, a spliced mRNA, a cDNA, or the like) include complementary nucleic acids. For example, the nucleic acid reagents may include oligonucleotides (labeled or non-labeled) fixed to a substrate, labeled oligonucleotides not bound with a substrate, pairs of PCR primers, molecular beacon probes, and the like.

- 39 -

The kit of the invention may optionally comprise additional components useful for performing the methods of the invention. By way of example, the kit may comprise fluids (*e.g.* SSC buffer) suitable for annealing complementary nucleic acids or for binding an antibody with a protein with which it specifically binds, one or more
5 sample compartments, an instructional material which describes performance of a method of the invention, a sample of normal ovarian cells, a sample of ovarian cancer cells, and the like.

The invention also includes a method of making an isolated hybridoma which produces an antibody useful for assessing whether patient is afflicted with an
10 ovarian cancer. In this method, a protein or peptide comprising the entirety or a segment of a marker protein is synthesized or isolated (*e.g.* by purification from a cell in which it is expressed or by transcription and translation of a nucleic acid encoding the protein or peptide *in vivo* or *in vitro* using known methods). A vertebrate, preferably a mammal such as a mouse, rat, rabbit, or sheep, is immunized using the protein or peptide. The
15 vertebrate may optionally (and preferably) be immunized at least one additional time with the protein or peptide, so that the vertebrate exhibits a robust immune response to the protein or peptide. Splenocytes are isolated from the immunized vertebrate and fused with an immortalized cell line to form hybridomas, using any of a variety of methods well known in the art. Hybridomas formed in this manner are then screened
20 using standard methods to identify one or more hybridomas which produce an antibody which specifically binds with the marker protein or a fragment thereof. The invention also includes hybridomas made by this method and antibodies made using such hybridomas.

The invention also includes a method of assessing the efficacy of a test
25 compound for inhibiting ovarian cancer cells. As described above, differences in the level of expression of the markers of the invention correlate with the cancerous state of ovarian cells. Although it is recognized that changes in the levels of expression of certain of the markers of the invention likely result from the cancerous state of ovarian cells, it is likewise recognized that changes in the levels of expression of other of the
30 markers of the invention induce, maintain, and promote the cancerous state of those cells. Thus, compounds which inhibit an ovarian cancer in a patient will cause the level of expression of one or more of the markers of the invention to change to a level nearer

the normal level of expression for that marker (*i.e.* the level of expression for the marker in non-cancerous ovarian cells).

This method thus comprises comparing expression of a marker in a first ovarian cell sample and maintained in the presence of the test compound and expression
5 of the marker in a second ovarian cell sample and maintained in the absence of the test compound. A significantly reduced expression of a marker of the invention in the presence of the test compound is an indication that the test compound inhibits ovarian cancer. The ovarian cell samples may, for example, be aliquots of a single sample of
10 normal ovarian cells obtained from a patient, pooled samples of normal ovarian cells obtained from a patient, cells of a normal ovarian cell line, aliquots of a single sample of ovarian cancer cells obtained from a patient, pooled samples of ovarian cancer cells
obtained from a patient, cells of an ovarian cancer cell line, or the like. In one embodiment, the samples are ovarian cancer cells obtained from a patient and a plurality of compounds known to be effective for inhibiting various ovarian cancers are tested in
15 order to identify the compound which is likely to best inhibit the ovarian cancer in the patient.

This method may likewise be used to assess the efficacy of a therapy for inhibiting ovarian cancer in a patient. In this method, the level of expression of one or more markers of the invention in a pair of samples (one subjected to the therapy, the
20 other not subjected to the therapy) is assessed. As with the method of assessing the efficacy of test compounds, if the therapy induces a significantly lower level of expression of a marker of the invention then the therapy is efficacious for inhibiting ovarian cancer. As above, if samples from a selected patient are used in this method, then alternative therapies can be assessed *in vitro* in order to select a therapy most likely
25 to be efficacious for inhibiting ovarian cancer in the patient.

As described above, the cancerous state of human ovarian cells is correlated with changes in the levels of expression of the markers of the invention. The invention includes a method for assessing the human ovarian cell carcinogenic potential of a test compound. This method comprises maintaining separate aliquots of human
30 ovarian cells in the presence and absence of the test compound. Expression of a marker of the invention in each of the aliquots is compared. A significantly higher level of expression of a marker of the invention in the aliquot maintained in the presence of the

test compound (relative to the aliquot maintained in the absence of the test compound) is an indication that the test compound possesses human ovarian cell carcinogenic potential. The relative carcinogenic potentials of various test compounds can be assessed by comparing the degree of enhancement or inhibition of the level of
5 expression of the relevant markers, by comparing the number of markers for which the level of expression is enhanced or inhibited, or by comparing both.

Various aspects of the invention are described in further detail in the following subsections.

10 I. Isolated Nucleic Acid Molecules

One aspect of the invention pertains to isolated nucleic acid molecules, including nucleic acids which encode a marker protein or a portion thereof. Isolated nucleic acids of the invention also include nucleic acid molecules sufficient for use as hybridization probes to identify marker nucleic acid molecules, and fragments of marker
15 nucleic acid molecules, *e.g.*, those suitable for use as PCR primers for the amplification or mutation of marker nucleic acid molecules. As used herein, the term "nucleic acid molecule" is intended to include DNA molecules (*e.g.*, cDNA or genomic DNA) and RNA molecules (*e.g.*, mRNA) and analogs of the DNA or RNA generated using nucleotide analogs. The nucleic acid molecule can be single-stranded or double-
20 stranded, but preferably is double-stranded DNA.

An "isolated" nucleic acid molecule is one which is separated from other nucleic acid molecules which are present in the natural source of the nucleic acid molecule. Preferably, an "isolated" nucleic acid molecule is free of sequences (preferably protein-encoding sequences) which naturally flank the nucleic acid (*i.e.*,
25 sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. For example, in various embodiments, the isolated nucleic acid molecule can contain less than about 5 kB, 4 kB, 3 kB, 2 kB, 1 kB, 0.5 kB or 0.1 kB of nucleotide sequences which naturally flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived. Moreover,
30 an "isolated" nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques,

- 42 -

or substantially free of chemical precursors or other chemicals when chemically synthesized.

5 A nucleic acid molecule of the present invention can be isolated using standard molecular biology techniques and the sequence information in the database records described herein. Using all or a portion of such nucleic acid sequences, nucleic acid molecules of the invention can be isolated using standard hybridization and cloning techniques (e.g., as described in Sambrook *et al.*, ed., *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989).

10 A nucleic acid molecule of the invention can be amplified using cDNA, mRNA, or genomic DNA as a template and appropriate oligonucleotide primers according to standard PCR amplification techniques. The nucleic acid so amplified can be cloned into an appropriate vector and characterized by DNA sequence analysis. Furthermore, nucleotides corresponding to all or a portion of a nucleic acid molecule of the invention can be prepared by standard synthetic techniques, e.g., using an automated DNA synthesizer.

In another preferred embodiment, an isolated nucleic acid molecule of the invention comprises a nucleic acid molecule which has a nucleotide sequence complementary to the nucleotide sequence of a marker nucleic acid or to the nucleotide sequence of a nucleic acid encoding a marker protein. A nucleic acid molecule which is complementary to a given nucleotide sequence is one which is sufficiently complementary to the given nucleotide sequence that it can hybridize to the given nucleotide sequence thereby forming a stable duplex.

Moreover, a nucleic acid molecule of the invention can comprise only a portion of a nucleic acid sequence, wherein the full length nucleic acid sequence comprises a marker nucleic acid or which encodes a marker protein. Such nucleic acids can be used, for example, as a probe or primer. The probe/primer typically is used as one or more substantially purified oligonucleotides. The oligonucleotide typically comprises a region of nucleotide sequence that hybridizes under stringent conditions to at least about 7, preferably about 15, more preferably about 25, 50, 75, 100, 125, 150, 175, 200, 250, 300, 350, or 400 or more consecutive nucleotides of a nucleic acid of the invention.

Probes based on the sequence of a nucleic acid molecule of the invention can be used to detect transcripts or genomic sequences corresponding to one or more markers of the invention. The probe comprises a label group attached thereto, *e.g.*, a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor. Such probes
5 can be used as part of a diagnostic test kit for identifying cells or tissues which mis-express the protein, such as by measuring levels of a nucleic acid molecule encoding the protein in a sample of cells from a subject, *e.g.*, detecting mRNA levels or determining whether a gene encoding the protein has been mutated or deleted.

The invention further encompasses nucleic acid molecules that differ, due
10 to degeneracy of the genetic code, from the nucleotide sequence of nucleic acids encoding a marker protein and thus encode the same protein.

It will be appreciated by those skilled in the art that DNA sequence polymorphisms that lead to changes in the amino acid sequence can exist within a population (*e.g.*, the human population). Such genetic polymorphisms can exist among
15 individuals within a population due to natural allelic variation. An allele is one of a group of genes which occur alternatively at a given genetic locus. In addition, it will be appreciated that DNA polymorphisms that affect RNA expression levels can also exist that may affect the overall expression level of that gene (*e.g.*, by affecting regulation or degradation).

20 As used herein, the phrase "allelic variant" refers to a nucleotide sequence which occurs at a given locus or to a polypeptide encoded by the nucleotide sequence.

As used herein, the terms "gene" and "recombinant gene" refer to nucleic acid molecules comprising an open reading frame encoding a polypeptide corresponding
25 to a marker of the invention. Such natural allelic variations can typically result in 1-5% variance in the nucleotide sequence of a given gene. Alternative alleles can be identified by sequencing the gene of interest in a number of different individuals. This can be readily carried out by using hybridization probes to identify the same genetic locus in a variety of individuals. Any and all such nucleotide variations and resulting amino acid
30 polymorphisms or variations that are the result of natural allelic variation and that do not alter the functional activity are intended to be within the scope of the invention.

- 44 -

In another embodiment, an isolated nucleic acid molecule of the invention is at least 7, 15, 20, 25, 30, 40, 60, 80, 100, 150, 200, 250, 300, 350, 400, 450, 550, 650, 700, 800, 900, 1000, 1200, 1400, 1600, 1800, 2000, 2200, 2400, 2600, 2800, 3000, 3500, 4000, 4500, or more nucleotides in length and hybridizes under stringent
5 conditions to a marker nucleic acid or to a nucleic acid encoding a marker protein. As used herein, the term "hybridizes under stringent conditions" is intended to describe conditions for hybridization and washing under which nucleotide sequences at least 60% (65%, 70%, preferably 75%) identical to each other typically remain hybridized to each other. Such stringent conditions are known to those skilled in the art and can be found
10 in sections 6.3.1-6.3.6 of *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989). A preferred, non-limiting example of stringent hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2X SSC, 0.1% SDS at 50-65°C.

In addition to naturally-occurring allelic variants of a nucleic acid
15 molecule of the invention that can exist in the population, the skilled artisan will further appreciate that sequence changes can be introduced by mutation thereby leading to changes in the amino acid sequence of the encoded protein, without altering the biological activity of the protein encoded thereby. For example, one can make nucleotide substitutions leading to amino acid substitutions at "non-essential" amino
20 acid residues. A "non-essential" amino acid residue is a residue that can be altered from the wild-type sequence without altering the biological activity, whereas an "essential" amino acid residue is required for biological activity. For example, amino acid residues that are not conserved or only semi-conserved among homologs of various species may be non-essential for activity and thus would be likely targets for alteration.
25 Alternatively, amino acid residues that are conserved among the homologs of various species (e.g., murine and human) may be essential for activity and thus would not be likely targets for alteration.

Accordingly, another aspect of the invention pertains to nucleic acid molecules encoding a variant marker protein that contain changes in amino acid residues
30 that are not essential for activity. Such variant marker proteins differ in amino acid sequence from the naturally-occurring marker proteins, yet retain biological activity. In one embodiment, such a variant marker protein has an amino acid sequence that is at

- 45 -

least about 40% identical, 50%, 60%, 70%, 80%, 90%, 95%, or 98% identical to the amino acid sequence of a marker protein.

- An isolated nucleic acid molecule encoding a variant marker protein can be created by introducing one or more nucleotide substitutions, additions or deletions
- 5 into the nucleotide sequence of marker nucleic acids, such that one or more amino acid residue substitutions, additions, or deletions are introduced into the encoded protein. Mutations can be introduced by standard techniques, such as site-directed mutagenesis and PCR-mediated mutagenesis. Preferably, conservative amino acid substitutions are made at one or more predicted non-essential amino acid residues. A "conservative
- 10 amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (*e.g.*, lysine, arginine, histidine), acidic side chains (*e.g.*, aspartic acid, glutamic acid), uncharged polar side chains (*e.g.*, glycine, asparagine, glutamine,
- 15 serine, threonine, tyrosine, cysteine), non-polar side chains (*e.g.*, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (*e.g.*, threonine, valine, isoleucine) and aromatic side chains (*e.g.*, tyrosine, phenylalanine, tryptophan, histidine). Alternatively, mutations can be introduced randomly along all or part of the coding sequence, such as by saturation mutagenesis,
- 20 and the resultant mutants can be screened for biological activity to identify mutants that retain activity. Following mutagenesis, the encoded protein can be expressed recombinantly and the activity of the protein can be determined.

- The present invention encompasses antisense nucleic acid molecules, *i.e.*, molecules which are complementary to a sense nucleic acid of the invention, *e.g.*,
- 25 complementary to the coding strand of a double-stranded marker cDNA molecule or complementary to a marker mRNA sequence. Accordingly, an antisense nucleic acid of the invention can hydrogen bond to (*i.e.* anneal with) a sense nucleic acid of the invention. The antisense nucleic acid can be complementary to an entire coding strand, or to only a portion thereof, *e.g.*, all or part of the protein coding region (or open reading
- 30 frame). An antisense nucleic acid molecule can also be antisense to all or part of a non-coding region of the coding strand of a nucleotide sequence encoding a marker protein.

The non-coding regions ("5' and 3' untranslated regions") are the 5' and 3' sequences which flank the coding region and are not translated into amino acids.

An antisense oligonucleotide can be, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 45, or 50 or more nucleotides in length. An antisense nucleic acid of the invention can be constructed using chemical synthesis and enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid (e.g., an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, e.g., phosphorothioate derivatives and acridine substituted nucleotides can be used. Examples of modified nucleotides which can be used to generate the antisense nucleic acid include 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine. Alternatively, the antisense nucleic acid can be produced biologically using an expression vector into which a nucleic acid has been sub-cloned in an antisense orientation (i.e., RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest, described further in the following subsection).

The antisense nucleic acid molecules of the invention are typically administered to a subject or generated *in situ* such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding a marker protein to thereby inhibit expression of the marker, e.g., by inhibiting transcription and/or translation. The

- 47 -

hybridization can be by conventional nucleotide complementarity to form a stable duplex, or, for example, in the case of an antisense nucleic acid molecule which binds to DNA duplexes, through specific interactions in the major groove of the double helix. Examples of a route of administration of antisense nucleic acid molecules of the invention includes direct injection at a tissue site or infusion of the antisense nucleic acid into an ovary-associated body fluid. Alternatively, antisense nucleic acid molecules can be modified to target selected cells and then administered systemically. For example, for systemic administration, antisense molecules can be modified such that they specifically bind to receptors or antigens expressed on a selected cell surface, *e.g.*, by linking the antisense nucleic acid molecules to peptides or antibodies which bind to cell surface receptors or antigens. The antisense nucleic acid molecules can also be delivered to cells using the vectors described herein. To achieve sufficient intracellular concentrations of the antisense molecules, vector constructs in which the antisense nucleic acid molecule is placed under the control of a strong pol II or pol III promoter are preferred.

An antisense nucleic acid molecule of the invention can be an α -anomeric nucleic acid molecule. An α -anomeric nucleic acid molecule forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual α -units, the strands run parallel to each other (Gaultier *et al.*, 1987, *Nucleic Acids Res.* 15:6625-6641). The antisense nucleic acid molecule can also comprise a 2'-*o*-methylribonucleotide (Inoue *et al.*, 1987, *Nucleic Acids Res.* 15:6131-6148) or a chimeric RNA-DNA analogue (Inoue *et al.*, 1987, *FEBS Lett.* 215:327-330).

The invention also encompasses ribozymes. Ribozymes are catalytic RNA molecules with ribonuclease activity which are capable of cleaving a single-stranded nucleic acid, such as an mRNA, to which they have a complementary region. Thus, ribozymes (*e.g.*, hammerhead ribozymes as described in Haselhoff and Gerlach, 1988, *Nature* 334:585-591) can be used to catalytically cleave mRNA transcripts to thereby inhibit translation of the protein encoded by the mRNA. A ribozyme having specificity for a nucleic acid molecule encoding a marker protein can be designed based upon the nucleotide sequence of a cDNA corresponding to the marker. For example, a derivative of a *Tetrahymena* L-19 IVS RNA can be constructed in which the nucleotide sequence of the active site is complementary to the nucleotide sequence to be cleaved

(see Cech *et al.* U.S. Patent No. 4,987,071; and Cech *et al.* U.S. Patent No. 5,116,742). Alternatively, an mRNA encoding a polypeptide of the invention can be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules (see, *e.g.*, Bartel and Szostak, 1993, *Science* 261:1411-1418).

- 5 The invention also encompasses nucleic acid molecules which form triple helical structures. For example, expression of a marker of the invention can be inhibited by targeting nucleotide sequences complementary to the regulatory region of the gene encoding the marker nucleic acid or protein (*e.g.*, the promoter and/or enhancer) to form triple helical structures that prevent transcription of the gene in target cells. See
- 10 generally Helene (1991) *Anticancer Drug Des.* 6(6):569-84; Helene (1992) *Ann. N.Y. Acad. Sci.* 660:27-36; and Maher (1992) *Bioassays* 14(12):807-15.

- In various embodiments, the nucleic acid molecules of the invention can be modified at the base moiety, sugar moiety or phosphate backbone to improve, *e.g.*, the stability, hybridization, or solubility of the molecule. For example, the deoxyribose
- 15 phosphate backbone of the nucleic acids can be modified to generate peptide nucleic acids (see Hyrup *et al.*, 1996, *Bioorganic & Medicinal Chemistry* 4(1): 5-23). As used herein, the terms "peptide nucleic acids" or "PNAs" refer to nucleic acid mimics, *e.g.*, DNA mimics, in which the deoxyribose phosphate backbone is replaced by a pseudopeptide backbone and only the four natural nucleobases are retained. The neutral
- 20 backbone of PNAs has been shown to allow for specific hybridization to DNA and RNA under conditions of low ionic strength. The synthesis of PNA oligomers can be performed using standard solid phase peptide synthesis protocols as described in Hyrup *et al.* (1996), *supra*; Perry-O'Keefe *et al.* (1996) *Proc. Natl. Acad. Sci. USA* 93:14670-675.

- 25 PNAs can be used in therapeutic and diagnostic applications. For example, PNAs can be used as antisense or antigene agents for sequence-specific modulation of gene expression by, *e.g.*, inducing transcription or translation arrest or inhibiting replication. PNAs can also be used, *e.g.*, in the analysis of single base pair mutations in a gene by, *e.g.*, PNA directed PCR clamping; as artificial restriction
- 30 enzymes when used in combination with other enzymes, *e.g.*, S1 nucleases (Hyrup (1996), *supra*; or as probes or primers for DNA sequence and hybridization (Hyrup, 1996, *supra*; Perry-O'Keefe *et al.*, 1996, *Proc. Natl. Acad. Sci. USA* 93:14670-675).

In another embodiment, PNAs can be modified, *e.g.*, to enhance their stability or cellular uptake, by attaching lipophilic or other helper groups to PNA, by the formation of PNA-DNA chimeras, or by the use of liposomes or other techniques of drug delivery known in the art. For example, PNA-DNA chimeras can be generated

5 which can combine the advantageous properties of PNA and DNA. Such chimeras allow DNA recognition enzymes, *e.g.*, RNase H and DNA polymerases, to interact with the DNA portion while the PNA portion would provide high binding affinity and specificity. PNA-DNA chimeras can be linked using linkers of appropriate lengths selected in terms of base stacking, number of bonds between the nucleobases, and

10 orientation (Hyrup, 1996, *supra*). The synthesis of PNA-DNA chimeras can be performed as described in Hyrup (1996, *supra*, and Finn *et al.* (1996) *Nucleic Acids Res.* 24(17):3357-63. For example, a DNA chain can be synthesized on a solid support using standard phosphoramidite coupling chemistry and modified nucleoside analogs. Compounds such as 5'-(4-methoxytrityl)amino-5'-deoxy-thymidine phosphoramidite can

15 be used as a link between the PNA and the 5' end of DNA (Mag *et al.*, 1989, *Nucleic Acids Res.* 17:5973-88). PNA monomers are then coupled in a step-wise manner to produce a chimeric molecule with a 5' PNA segment and a 3' DNA segment (Finn *et al.*, 1996, *Nucleic Acids Res.* 24(17):3357-63). Alternatively, chimeric molecules can be synthesized with a 5' DNA segment and a 3' PNA segment (Peterser *et al.*, 1975,

20 *Bioorganic Med. Chem. Lett.* 5:1119-11124).

In other embodiments, the oligonucleotide can include other appended groups such as peptides (*e.g.*, for targeting host cell receptors *in vivo*), or agents facilitating transport across the cell membrane (see, *e.g.*, Letsinger *et al.*, 1989, *Proc. Natl. Acad. Sci. USA* 86:6553-6556; Lemaitre *et al.*, 1987, *Proc. Natl. Acad. Sci. USA*

25 84:648-652; PCT Publication No. WO 88/09810) or the blood-brain barrier (see, *e.g.*, PCT Publication No. WO 89/10134). In addition, oligonucleotides can be modified with hybridization-triggered cleavage agents (see, *e.g.*, Krol *et al.*, 1988, *Bio/Techniques* 6:958-976) or intercalating agents (see, *e.g.*, Zon, 1988, *Pharm. Res.* 5:539-549). To this end, the oligonucleotide can be conjugated to another molecule, *e.g.*, a peptide,

30 hybridization triggered cross-linking agent, transport agent, hybridization-triggered cleavage agent, etc.

- 50 -

The invention also includes molecular beacon nucleic acids having at least one region which is complementary to a nucleic acid of the invention, such that the molecular beacon is useful for quantitating the presence of the nucleic acid of the invention in a sample. A "molecular beacon" nucleic acid is a nucleic acid comprising a pair of complementary regions and having a fluorophore and a fluorescent quencher associated therewith. The fluorophore and quencher are associated with different portions of the nucleic acid in such an orientation that when the complementary regions are annealed with one another, fluorescence of the fluorophore is quenched by the quencher. When the complementary regions of the nucleic acid are not annealed with one another, fluorescence of the fluorophore is quenched to a lesser degree. Molecular beacon nucleic acids are described, for example, in U.S. Patent 5,876,930.

II. Isolated Proteins and Antibodies

One aspect of the invention pertains to isolated marker proteins and biologically active portions thereof, as well as polypeptide fragments suitable for use as immunogens to raise antibodies directed against a marker protein or a fragment thereof. In one embodiment, the native marker protein can be isolated from cells or tissue sources by an appropriate purification scheme using standard protein purification techniques. In another embodiment, a protein or peptide comprising the whole or a segment of the marker protein is produced by recombinant DNA techniques. Alternative to recombinant expression, such protein or peptide can be synthesized chemically using standard peptide synthesis techniques.

An "isolated" or "purified" protein or biologically active portion thereof is substantially free of cellular material or other contaminating proteins from the cell or tissue source from which the protein is derived, or substantially free of chemical precursors or other chemicals when chemically synthesized. The language "substantially free of cellular material" includes preparations of protein in which the protein is separated from cellular components of the cells from which it is isolated or recombinantly produced. Thus, protein that is substantially free of cellular material includes preparations of protein having less than about 30%, 20%, 10%, or 5% (by dry weight) of heterologous protein (also referred to herein as a "contaminating protein"). When the protein or biologically active portion thereof is recombinantly produced, it is

also preferably substantially free of culture medium, *i.e.*, culture medium represents less than about 20%, 10%, or 5% of the volume of the protein preparation. When the protein is produced by chemical synthesis, it is preferably substantially free of chemical precursors or other chemicals, *i.e.*, it is separated from chemical precursors or other chemicals which are involved in the synthesis of the protein. Accordingly such preparations of the protein have less than about 30%, 20%, 10%, 5% (by dry weight) of chemical precursors or compounds other than the polypeptide of interest.

Biologically active portions of a marker protein include polypeptides comprising amino acid sequences sufficiently identical to or derived from the amino acid sequence of the marker protein, which include fewer amino acids than the full length protein, and exhibit at least one activity of the corresponding full-length protein. Typically, biologically active portions comprise a domain or motif with at least one activity of the corresponding full-length protein. A biologically active portion of a marker protein of the invention can be a polypeptide which is, for example, 10, 25, 50, 100 or more amino acids in length. Moreover, other biologically active portions, in which other regions of the marker protein are deleted, can be prepared by recombinant techniques and evaluated for one or more of the functional activities of the native form of the marker protein.

Preferred marker proteins are encoded by nucleotide sequences comprising the sequences listed in Tables 1-3. Other useful proteins are substantially identical (*e.g.*, at least about 40%, preferably 50%, 60%, 70%, 80%, 90%, 95%, or 99%) to one of these sequences and retain the functional activity of the corresponding naturally-occurring marker protein yet differ in amino acid sequence due to natural allelic variation or mutagenesis.

To determine the percent identity of two amino acid sequences or of two nucleic acids, the sequences are aligned for optimal comparison purposes (*e.g.*, gaps can be introduced in the sequence of a first amino acid or nucleic acid sequence for optimal alignment with a second amino or nucleic acid sequence). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences

- 52 -

is a function of the number of identical positions shared by the sequences (*i.e.*, % identity = # of identical positions/total # of positions (*e.g.*, overlapping positions) $\times 100$). In one embodiment the two sequences are the same length.

- The determination of percent identity between two sequences can be accomplished using a mathematical algorithm. A preferred, non-limiting example of a mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin and Altschul (1990) *Proc. Natl. Acad. Sci. USA* 87:2264-2268, modified as in Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90:5873-5877. Such an algorithm is incorporated into the BLASTN and BLASTX programs of Altschul, *et al.* (1990) *J. Mol. Biol.* 215:403-410. BLAST nucleotide searches can be performed with the BLASTN program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to a nucleic acid molecules of the invention. BLAST protein searches can be performed with the BLASTP program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to a protein molecules of the invention. To obtain gapped alignments for comparison purposes, a newer version of the BLAST algorithm called Gapped BLAST can be utilized as described in Altschul *et al.* (1997) *Nucleic Acids Res.* 25:3389-3402, which is able to perform gapped local alignments for the programs BLASTN, BLASTP and BLASTX. Alternatively, PSI-Blast can be used to perform an iterated search which detects distant relationships between molecules. When utilizing BLAST, Gapped BLAST, and PSI-Blast programs, the default parameters of the respective programs (*e.g.*, BLASTX and BLASTN) can be used. Another preferred, non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller, (1988) *CABIOS* 4:11-17. Such an algorithm is incorporated into the ALIGN program (version 2.0) which is part of the GCG sequence alignment software package. When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used. Yet another useful algorithm for identifying regions of local sequence similarity and alignment is the FASTA algorithm as described in Pearson and Lipman (1988) *Proc. Natl. Acad. Sci. USA* 85:2444-2448. When using the FASTA algorithm for comparing nucleotide or amino acid sequences, a PAM120 weight residue table can, for example, be used with a *k*-tuple value of 2.

- 53 -

The percent identity between two sequences can be determined using techniques similar to those described above, with or without allowing gaps. In calculating percent identity, only exact matches are counted.

The invention also provides chimeric or fusion proteins comprising a marker protein or a segment thereof. As used herein, a "chimeric protein" or "fusion protein" comprises all or part (preferably a biologically active part) of a marker protein operably linked to a heterologous polypeptide (*i.e.*, a polypeptide other than the marker protein). Within the fusion protein, the term "operably linked" is intended to indicate that the marker protein or segment thereof and the heterologous polypeptide are fused in-frame to each other. The heterologous polypeptide can be fused to the amino-terminus or the carboxyl-terminus of the marker protein or segment.

One useful fusion protein is a GST fusion protein in which a marker protein or segment is fused to the carboxyl terminus of GST sequences. Such fusion proteins can facilitate the purification of a recombinant polypeptide of the invention.

In another embodiment, the fusion protein contains a heterologous signal sequence at its amino terminus. For example, the native signal sequence of a marker protein can be removed and replaced with a signal sequence from another protein. For example, the gp67 secretory sequence of the baculovirus envelope protein can be used as a heterologous signal sequence (Ausubel *et al.*, ed., *Current Protocols in Molecular Biology*, John Wiley & Sons, NY, 1992). Other examples of eukaryotic heterologous signal sequences include the secretory sequences of melittin and human placental alkaline phosphatase (Stratagene; La Jolla, California). In yet another example, useful prokaryotic heterologous signal sequences include the phoA secretory signal (Sambrook *et al.*, *supra*) and the protein A secretory signal (Pharmacia Biotech; Piscataway, New Jersey).

In yet another embodiment, the fusion protein is an immunoglobulin fusion protein in which all or part of a marker protein is fused to sequences derived from a member of the immunoglobulin protein family. The immunoglobulin fusion proteins of the invention can be incorporated into pharmaceutical compositions and administered to a subject to inhibit an interaction between a ligand (soluble or membrane-bound) and a protein on the surface of a cell (receptor), to thereby suppress signal transduction *in vivo*. The immunoglobulin fusion protein can be used to affect the bioavailability of a

cognate ligand of a marker protein. Inhibition of ligand/receptor interaction can be useful therapeutically, both for treating proliferative and differentiative disorders and for modulating (*e.g.* promoting or inhibiting) cell survival. Moreover, the immunoglobulin fusion proteins of the invention can be used as immunogens to produce antibodies
5 directed against a marker protein in a subject, to purify ligands and in screening assays to identify molecules which inhibit the interaction of the marker protein with ligands.

- Chimeric and fusion proteins of the invention can be produced by standard recombinant DNA techniques. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers.
- 10 Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed and re-amplified to generate a chimeric gene sequence (see, *e.g.*, Ausubel *et al.*, *supra*). Moreover, many expression vectors are commercially available that already encode a fusion moiety (*e.g.*, a GST polypeptide).
- 15 A nucleic acid encoding a polypeptide of the invention can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the polypeptide of the invention.

- A signal sequence can be used to facilitate secretion and isolation of marker proteins. Signal sequences are typically characterized by a core of hydrophobic
20 amino acids which are generally cleaved from the mature protein during secretion in one or more cleavage events. Such signal peptides contain processing sites that allow cleavage of the signal sequence from the mature proteins as they pass through the secretory pathway. Thus, the invention pertains to marker proteins, fusion proteins or segments thereof having a signal sequence, as well as to such proteins from which the
- 25 signal sequence has been proteolytically cleaved (*i.e.*, the cleavage products). In one embodiment, a nucleic acid sequence encoding a signal sequence can be operably linked in an expression vector to a protein of interest, such as a marker protein or a segment thereof. The signal sequence directs secretion of the protein, such as from a eukaryotic host into which the expression vector is transformed, and the signal sequence is
- 30 subsequently or concurrently cleaved. The protein can then be readily purified from the extracellular medium by art recognized methods. Alternatively, the signal sequence can

be linked to the protein of interest using a sequence which facilitates purification, such as with a GST domain.

The present invention also pertains to variants of the marker proteins. Such variants have an altered amino acid sequence which can function as either agonists (mimetics) or as antagonists. Variants can be generated by mutagenesis, *e.g.*, discrete point mutation or truncation. An agonist can retain substantially the same, or a subset, of the biological activities of the naturally occurring form of the protein. An antagonist of a protein can inhibit one or more of the activities of the naturally occurring form of the protein by, for example, competitively binding to a downstream or upstream member of a cellular signaling cascade which includes the protein of interest. Thus, specific biological effects can be elicited by treatment with a variant of limited function. Treatment of a subject with a variant having a subset of the biological activities of the naturally occurring form of the protein can have fewer side effects in a subject relative to treatment with the naturally occurring form of the protein.

15 Variants of a marker protein which function as either agonists (mimetics) or as antagonists can be identified by screening combinatorial libraries of mutants, *e.g.*, truncation mutants, of the protein of the invention for agonist or antagonist activity. In one embodiment, a variegated library of variants is generated by combinatorial mutagenesis at the nucleic acid level and is encoded by a variegated gene library. A
20 variegated library of variants can be produced by, for example, enzymatically ligating a mixture of synthetic oligonucleotides into gene sequences such that a degenerate set of potential protein sequences is expressible as individual polypeptides, or alternatively, as a set of larger fusion proteins (*e.g.*, for phage display). There are a variety of methods which can be used to produce libraries of potential variants of the marker proteins from
25 a degenerate oligonucleotide sequence. Methods for synthesizing degenerate oligonucleotides are known in the art (see, *e.g.*, Narang, 1983, *Tetrahedron* 39:3; Itakura *et al.*, 1984, *Annu. Rev. Biochem.* 53:323; Itakura *et al.*, 1984, *Science* 198:1056; Ike *et al.*, 1983 *Nucleic Acid Res.* 11:477).

In addition, libraries of segments of a marker protein can be used to
30 generate a variegated population of polypeptides for screening and subsequent selection of variant marker proteins or segments thereof. For example, a library of coding sequence fragments can be generated by treating a double stranded PCR fragment of the

- 56 -

coding sequence of interest with a nuclease under conditions wherein nicking occurs only about once per molecule, denaturing the double stranded DNA, renaturing the DNA to form double stranded DNA which can include sense/antisense pairs from different nicked products, removing single stranded portions from reformed duplexes by treatment with S1 nuclease, and ligating the resulting fragment library into an expression vector. By this method, an expression library can be derived which encodes amino terminal and internal fragments of various sizes of the protein of interest.

Several techniques are known in the art for screening gene products of combinatorial libraries made by point mutations or truncation, and for screening cDNA libraries for gene products having a selected property. The most widely used techniques, which are amenable to high through-put analysis, for screening large gene libraries typically include cloning the gene library into replicable expression vectors, transforming appropriate cells with the resulting library of vectors, and expressing the combinatorial genes under conditions in which detection of a desired activity facilitates isolation of the vector encoding the gene whose product was detected. Recursive ensemble mutagenesis (REM), a technique which enhances the frequency of functional mutants in the libraries, can be used in combination with the screening assays to identify variants of a protein of the invention (Arkin and Yourvan, 1992, *Proc. Natl. Acad. Sci. USA* 89:7811-7815; Delgrave *et al.*, 1993, *Protein Engineering* 6(3):327- 331).

Another aspect of the invention pertains to antibodies directed against a protein of the invention. In preferred embodiments, the antibodies specifically bind a marker protein or a fragment thereof. The terms "antibody" and "antibodies" as used interchangeably herein refer to immunoglobulin molecules as well as fragments and derivatives thereof that comprise an immunologically active portion of an immunoglobulin molecule, (*i.e.*, such a portion contains an antigen binding site which specifically binds an antigen, such as a marker protein, *e.g.*, an epitope of a marker protein). An antibody which specifically binds to a protein of the invention is an antibody which binds the protein, but does not substantially bind other molecules in a sample, *e.g.*, a biological sample, which naturally contains the protein. Examples of an immunologically active portion of an immunoglobulin molecule include, but are not limited to, single-chain antibodies (scAb), F(ab) and F(ab')₂ fragments.

- 57 -

An isolated protein of the invention or a fragment thereof can be used as an immunogen to generate antibodies. The full-length protein can be used or, alternatively, the invention provides antigenic peptide fragments for use as immunogens. The antigenic peptide of a protein of the invention comprises at least 8 (preferably 10, 15, 20, or 30 or more) amino acid residues of the amino acid sequence of one of the proteins of the invention, and encompasses at least one epitope of the protein such that an antibody raised against the peptide forms a specific immune complex with the protein. Preferred epitopes encompassed by the antigenic peptide are regions that are located on the surface of the protein, *e.g.*, hydrophilic regions. Hydrophobicity sequence analysis, hydrophilicity sequence analysis, or similar analyses can be used to identify hydrophilic regions. In preferred embodiments, an isolated marker protein or fragment thereof is used as an immunogen.

An immunogen typically is used to prepare antibodies by immunizing a suitable (*i.e.* immunocompetent) subject such as a rabbit, goat, mouse, or other mammal or vertebrate. An appropriate immunogenic preparation can contain, for example, recombinantly-expressed or chemically-synthesized protein or peptide. The preparation can further include an adjuvant, such as Freund's complete or incomplete adjuvant, or a similar immunostimulatory agent. Preferred immunogen compositions are those that contain no other human proteins such as, for example, immunogen compositions made using a non-human host cell for recombinant expression of a protein of the invention. In such a manner, the resulting antibody compositions have reduced or no binding of human proteins other than a protein of the invention.

The invention provides polyclonal and monoclonal antibodies. The term "monoclonal antibody" or "monoclonal antibody composition", as used herein, refers to a population of antibody molecules that contain only one species of an antigen binding site capable of immunoreacting with a particular epitope. Preferred polyclonal and monoclonal antibody compositions are ones that have been selected for antibodies directed against a protein of the invention. Particularly preferred polyclonal and monoclonal antibody preparations are ones that contain only antibodies directed against a marker protein or fragment thereof.

- 58 -

Polyclonal antibodies can be prepared by immunizing a suitable subject with a protein of the invention as an immunogen. The antibody titer in the immunized subject can be monitored over time by standard techniques, such as with an enzyme linked immunosorbent assay (ELISA) using immobilized polypeptide. At an appropriate time after immunization, *e.g.*, when the specific antibody titers are highest, antibody-producing cells can be obtained from the subject and used to prepare monoclonal antibodies (mAb) by standard techniques, such as the hybridoma technique originally described by Kohler and Milstein (1975) *Nature* 256:495-497, the human B cell hybridoma technique (see Kozbor *et al.*, 1983, *Immunol. Today* 4:72), the EBV-hybridoma technique (see Cole *et al.*, pp. 77-96 In *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., 1985) or trioma techniques. The technology for producing hybridomas is well known (see generally *Current Protocols in Immunology*, Coligan *et al.* ed., John Wiley & Sons, New York, 1994). Hybridoma cells producing a monoclonal antibody of the invention are detected by screening the hybridoma culture supernatants for antibodies that bind the polypeptide of interest, *e.g.*, using a standard ELISA assay.

Alternative to preparing monoclonal antibody-secreting hybridomas, a monoclonal antibody directed against a protein of the invention can be identified and isolated by screening a recombinant combinatorial immunoglobulin library (*e.g.*, an antibody phage display library) with the polypeptide of interest. Kits for generating and screening phage display libraries are commercially available (*e.g.*, the Pharmacia *Recombinant Phage Antibody System*, Catalog No. 27-9400-01; and the Stratagene *SurfZAP Phage Display Kit*, Catalog No. 240612). Additionally, examples of methods and reagents particularly amenable for use in generating and screening antibody display library can be found in, for example, U.S. Patent No. 5,223,409; PCT Publication No. WO 92/18619; PCT Publication No. WO 91/17271; PCT Publication No. WO 92/20791; PCT Publication No. WO 92/15679; PCT Publication No. WO 93/01288; PCT Publication No. WO 92/01047; PCT Publication No. WO 92/09690; PCT Publication No. WO 90/02809; Fuchs *et al.* (1991) *Bio/Technology* 9:1370-1372; Hay *et al.* (1992) *Hum. Antibod. Hybridomas* 3:81-85; Huse *et al.* (1989) *Science* 246:1275-1281; Griffiths *et al.* (1993) *EMBO J.* 12:725-734.

The invention also provides recombinant antibodies that specifically bind a protein of the invention. In preferred embodiments, the recombinant antibodies specifically binds a marker protein or fragment thereof. Recombinant antibodies include, but are not limited to, chimeric and humanized monoclonal antibodies, comprising both human and non-human portions, single-chain antibodies and multi-specific antibodies. A chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region. (See, e.g., Cabilly et al., U.S. Patent No. 4,816,567; and Boss et al., U.S. Patent No. 4,816,397, which are incorporated herein by reference in their entirety.) Single-chain antibodies have an antigen binding site and consist of single polypeptides. They can be produced by techniques known in the art, for example using methods described in Ladner et al. U.S. Pat. No. 4,946,778 (which is incorporated herein by reference in its entirety); Bird et al., (1988) *Science* 242:423-426; Whitlow et al., (1991) *Methods in Enzymology* 2:1-9; Whitlow et al., (1991) *Methods in Enzymology* 2:97-105; and Huston et al., (1991) *Methods in Enzymology Molecular Design and Modeling: Concepts and Applications* 203:46-88. Multi-specific antibodies are antibody molecules having at least two antigen-binding sites that specifically bind different antigens. Such molecules can be produced by techniques known in the art, for example using methods described in Segal, U.S. Patent No. 4,676,980 (the disclosure of which is incorporated herein by reference in its entirety); Holliger et al., (1993) *Proc. Natl. Acad. Sci. USA* 90:6444-6448; Whitlow et al., (1994) *Protein Eng.* 7:1017-1026 and U.S. Pat. No. 6,121,424.

Humanized antibodies are antibody molecules from non-human species having one or more complementarity determining regions (CDRs) from the non-human species and a framework region from a human immunoglobulin molecule. (See, e.g., Queen, U.S. Patent No. 5,585,089, which is incorporated herein by reference in its entirety.) Humanized monoclonal antibodies can be produced by recombinant DNA techniques known in the art, for example using methods described in PCT Publication No. WO 87/02671; European Patent Application 184,187; European Patent Application 171,496; European Patent Application 173,494; PCT Publication No. WO 86/01533; U.S. Patent No. 4,816,567; European Patent Application 125,023; Better et al. (1988) *Science* 240:1041-1043; Liu et al. (1987) *Proc. Natl. Acad. Sci. USA* 84:3439-3443; Liu

- 60 -

- et al.* (1987) *J. Immunol.* 139:3521- 3526; Sun *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:214-218; Nishimura *et al.* (1987) *Cancer Res.* 47:999-1005; Wood *et al.* (1985) *Nature* 314:446-449; and Shaw *et al.* (1988) *J. Natl. Cancer Inst.* 80:1553-1559; Morrison (1985) *Science* 229:1202-1207; Oi *et al.* (1986) *Bio/Techniques* 4:214; U.S. Patent 5,225,539; Jones *et al.* (1986) *Nature* 321:552-525; Verhoeven *et al.* (1988) *Science* 239:1534; and Beidler *et al.* (1988) *J. Immunol.* 141:4053-4060.

More particularly, humanized antibodies can be produced, for example, using transgenic mice which are incapable of expressing endogenous immunoglobulin heavy and light chains genes, but which can express human heavy and light chain genes.

10 The transgenic mice are immunized in the normal fashion with a selected antigen, *e.g.*, all or a portion of a polypeptide corresponding to a marker of the invention. Monoclonal antibodies directed against the antigen can be obtained using conventional hybridoma technology. The human immunoglobulin transgenes harbored by the transgenic mice rearrange during B cell differentiation, and subsequently undergo class

15 switching and somatic mutation. Thus, using such a technique, it is possible to produce therapeutically useful IgG, IgA and IgE antibodies. For an overview of this technology for producing human antibodies, see Lonberg and Huszar (1995) *Int. Rev. Immunol.* 13:65-93). For a detailed discussion of this technology for producing human antibodies and human monoclonal antibodies and protocols for producing such antibodies, see, *e.g.*,

20 U.S. Patent 5,625,126; U.S. Patent 5,633,425; U.S. Patent 5,569,825; U.S. Patent 5,661,016; and U.S. Patent 5,545,806. In addition, companies such as Abgenix, Inc. (Freemont, CA), can be engaged to provide human antibodies directed against a selected antigen using technology similar to that described above.

Completely human antibodies which recognize a selected epitope can be

25 generated using a technique referred to as "guided selection." In this approach a selected non-human monoclonal antibody, *e.g.*, a murine antibody, is used to guide the selection of a completely human antibody recognizing the same epitope (Jespersen *et al.*, 1994, *Bio/technology* 12:899-903).

The antibodies of the invention can be isolated after production (*e.g.*,

30 from the blood or serum of the subject) or synthesis and further purified by well-known techniques. For example, IgG antibodies can be purified using protein A chromatography. Antibodies specific for a protein of the invention can be selected or

- 61 -

(*e.g.*, partially purified) or purified by, *e.g.*, affinity chromatography. For example, a recombinantly expressed and purified (or partially purified) protein of the invention is produced as described herein, and covalently or non-covalently coupled to a solid support such as, for example, a chromatography column. The column can then be used to affinity purify antibodies specific for the proteins of the invention from a sample containing antibodies directed against a large number of different epitopes, thereby generating a substantially purified antibody composition, *i.e.*, one that is substantially free of contaminating antibodies. By a substantially purified antibody composition is meant, in this context, that the antibody sample contains at most only 30% (by dry weight) of contaminating antibodies directed against epitopes other than those of the desired protein of the invention, and preferably at most 20%, yet more preferably at most 10%, and most preferably at most 5% (by dry weight) of the sample is contaminating antibodies. A purified antibody composition means that at least 99% of the antibodies in the composition are directed against the desired protein of the invention.

In a preferred embodiment, the substantially purified antibodies of the invention may specifically bind to a signal peptide, a secreted sequence, an extracellular domain, a transmembrane or a cytoplasmic domain or cytoplasmic membrane of a protein of the invention. In a particularly preferred embodiment, the substantially purified antibodies of the invention specifically bind to a secreted sequence or an extracellular domain of the amino acid sequences of a protein of the invention. In a more preferred embodiment, the substantially purified antibodies of the invention specifically bind to a secreted sequence or an extracellular domain of the amino acid sequences of a marker protein.

An antibody directed against a protein of the invention can be used to isolate the protein by standard techniques, such as affinity chromatography or immunoprecipitation. Moreover, such an antibody can be used to detect the marker protein or fragment thereof (*e.g.*, in a cellular lysate or cell supernatant) in order to evaluate the level and pattern of expression of the marker. The antibodies can also be used diagnostically to monitor protein levels in tissues or body fluids (*e.g.* in an ovary-associated body fluid) as part of a clinical testing procedure, *e.g.*, to, for example, determine the efficacy of a given treatment regimen. Detection can be facilitated by the

- use of an antibody derivative, which comprises an antibody of the invention coupled to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish
- 5 peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol;
- 10 examples of bioluminescent materials include luciferase, luciferin, and aquorin, and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S or ^3H .

- Antibodies of the invention may also be used as therapeutic agents in treating cancers. In a preferred embodiment, completely human antibodies of the invention are used for therapeutic treatment of human cancer patients, particularly those
- 15 having an ovarian cancer. In another preferred embodiment, antibodies that bind specifically to a marker protein or fragment thereof are used for therapeutic treatment. Further, such therapeutic antibody may be an antibody derivative or immunotoxin comprising an antibody conjugated to a therapeutic moiety such as a cytotoxin, a therapeutic agent or a radioactive metal ion. A cytotoxin or cytotoxic agent includes any
- 20 agent that is detrimental to cells. Examples include taxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, teniposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof.
- 25 Therapeutic agents include, but are not limited to, antimetabolites (*e.g.*, methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (*e.g.*, mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BSNU) and lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis-dichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines
- 30 (*e.g.*, daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (*e.g.*, dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC)), and anti-mitotic agents (*e.g.*, vincristine and vinblastine).

- 63 -

The conjugated antibodies of the invention can be used for modifying a given biological response, for the drug moiety is not to be construed as limited to classical chemical therapeutic agents. For example, the drug moiety may be a protein or polypeptide possessing a desired biological activity. Such proteins may include, for example, a toxin such as ribosome-inhibiting protein (see Better et al., U.S. Patent No. 6,146,631, the disclosure of which is incorporated herein in its entirety), abrin, ricin A, pseudomonas exotoxin, or diphtheria toxin; a protein such as tumor necrosis factor, .alpha.-interferon, .beta.-interferon, nerve growth factor, platelet derived growth factor, tissue plasminogen activator; or, biological response modifiers such as, for example, lymphokines, interleukin-1 ("IL-1"), interleukin-2 ("IL-2"), interleukin-6 ("IL-6"), granulocyte macrophase colony stimulating factor ("GM-CSF"), granulocyte colony stimulating factor ("G-CSF"), or other growth factors.

Techniques for conjugating such therapeutic moiety to antibodies are well known, see, *e.g.*, Arnon et al., "Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy", in *Monoclonal Antibodies And Cancer Therapy*, Reisfeld et al. (eds.), pp. 243-56 (Alan R. Liss, Inc. 1985); Hellstrom et al., "Antibodies For Drug Delivery", in *Controlled Drug Delivery* (2nd Ed.), Robinson et al. (eds.), pp. 623-53 (Marcel Dekker, Inc. 1987); Thorpe, "Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review", in *Monoclonal Antibodies '84: Biological And Clinical Applications*, Pinchera et al. (eds.), pp. 475-506 (1985); "Analysis, Results, And Future Prospective Of The Therapeutic Use Of Radiolabeled Antibody In Cancer Therapy", in *Monoclonal Antibodies For Cancer Detection And Therapy*, Baldwin et al. (eds.), pp. 303-16 (Academic Press 1985), and Thorpe et al., "The Preparation And Cytotoxic Properties Of Antibody-Toxin Conjugates", *Immunol. Rev.*, 62:119-58 (1982).

Accordingly, in one aspect, the invention provides substantially purified antibodies, antibody fragments and derivatives, all of which specifically bind to a protein of the invention and preferably, a marker protein. In various embodiments, the substantially purified antibodies of the invention, or fragments or derivatives thereof, can be human, non-human, chimeric and/or humanized antibodies. In another aspect, the invention provides non-human antibodies, antibody fragments and derivatives, all of which specifically bind to a protein of the invention and preferably, a marker protein. Such non-human antibodies can be goat, mouse, sheep, horse, chicken, rabbit, or rat

antibodies. Alternatively, the non-human antibodies of the invention can be chimeric and/or humanized antibodies. In addition, the non-human antibodies of the invention can be polyclonal antibodies or monoclonal antibodies. In still a further aspect, the invention provides monoclonal antibodies, antibody fragments and derivatives, all of which specifically bind to a protein of the invention and preferably, a marker protein. The monoclonal antibodies can be human, humanized, chimeric and/or non-human antibodies.

The invention also provides a kit containing an antibody of the invention conjugated to a detectable substance, and instructions for use. Still another aspect of the invention is a pharmaceutical composition comprising an antibody of the invention and a pharmaceutically acceptable carrier. In preferred embodiments, the pharmaceutical composition contains an antibody of the invention, a therapeutic moiety, and a pharmaceutically acceptable carrier.

15 III. Recombinant Expression Vectors and Host Cells

Another aspect of the invention pertains to vectors, preferably expression vectors, containing a nucleic acid encoding a marker protein (or a portion of such a protein). As used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments can be ligated. Another type of vector is a viral vector, wherein additional DNA segments can be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (*e.g.*, bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (*e.g.*, non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors, namely expression vectors, are capable of directing the expression of genes to which they are operably linked. In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids (vectors). However, the invention is intended to include such other forms of expression vectors, such as viral vectors (*e.g.*, replication defective

- 65 -

retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

The recombinant expression vectors of the invention comprise a nucleic acid of the invention in a form suitable for expression of the nucleic acid in a host cell.

- 5 This means that the recombinant expression vectors include one or more regulatory sequences, selected on the basis of the host cells to be used for expression, which is operably linked to the nucleic acid sequence to be expressed. Within a recombinant expression vector, "operably linked" is intended to mean that the nucleotide sequence of interest is linked to the regulatory sequence(s) in a manner which allows for expression
- 10 of the nucleotide sequence (*e.g.*, in an *in vitro* transcription/translation system or in a host cell when the vector is introduced into the host cell). The term "regulatory sequence" is intended to include promoters, enhancers and other expression control elements (*e.g.*, polyadenylation signals). Such regulatory sequences are described, for example, in Goeddel, *Methods in Enzymology: Gene Expression Technology* vol.185,
- 15 Academic Press, San Diego, CA (1991). Regulatory sequences include those which direct constitutive expression of a nucleotide sequence in many types of host cell and those which direct expression of the nucleotide sequence only in certain host cells (*e.g.*, tissue-specific regulatory sequences). It will be appreciated by those skilled in the art that the design of the expression vector can depend on such factors as the choice of the
- 20 host cell to be transformed, the level of expression of protein desired, and the like. The expression vectors of the invention can be introduced into host cells to thereby produce proteins or peptides, including fusion proteins or peptides, encoded by nucleic acids as described herein.

- The recombinant expression vectors of the invention can be designed for
- 25 expression of a marker protein or a segment thereof in prokaryotic (*e.g.*, *E. coli*) or eukaryotic cells (*e.g.*, insect cells {using baculovirus expression vectors}, yeast cells or mammalian cells). Suitable host cells are discussed further in Goeddel, *supra*. Alternatively, the recombinant expression vector can be transcribed and translated *in vitro*, for example using T7 promoter regulatory sequences and T7 polymerase.

- 30 Expression of proteins in prokaryotes is most often carried out in *E. coli* with vectors containing constitutive or inducible promoters directing the expression of either fusion or non-fusion proteins. Fusion vectors add a number of amino acids to a

- protein encoded therein, usually to the amino terminus of the recombinant protein. Such fusion vectors typically serve three purposes: 1) to increase expression of recombinant protein; 2) to increase the solubility of the recombinant protein; and 3) to aid in the purification of the recombinant protein by acting as a ligand in affinity purification.
- 5 Often, in fusion expression vectors, a proteolytic cleavage site is introduced at the junction of the fusion moiety and the recombinant protein to enable separation of the recombinant protein from the fusion moiety subsequent to purification of the fusion protein. Such enzymes, and their cognate recognition sequences, include Factor Xa, thrombin and enterokinase. Typical fusion expression vectors include pGEX
- 10 (Pharmacia Biotech Inc; Smith and Johnson, 1988, *Gene* 67:31-40), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein.

- Examples of suitable inducible non-fusion *E. coli* expression vectors
- 15 include pTrc (Amann *et al.*, 1988, *Gene* 69:301-315) and pET 11d (Studier *et al.*, p. 60-89, In *Gene Expression Technology: Methods in Enzymology* vol.185, Academic Press, San Diego, CA, 1991). Target gene expression from the pTrc vector relies on host RNA polymerase transcription from a hybrid trp-lac fusion promoter. Target gene expression from the pET 11d vector relies on transcription from a T7 gn10-lac fusion promoter
- 20 mediated by a co-expressed viral RNA polymerase (T7 gn1). This viral polymerase is supplied by host strains BL21(DE3) or HMS174(DE3) from a resident prophage harboring a T7 gn1 gene under the transcriptional control of the lacUV 5 promoter.

- One strategy to maximize recombinant protein expression in *E. coli* is to express the protein in a host bacteria with an impaired capacity to proteolytically cleave the recombinant protein (Gottesman, p. 119-128, In *Gene Expression Technology: Methods in Enzymology* vol. 185, Academic Press, San Diego, CA, 1990. Another strategy is to alter the nucleic acid sequence of the nucleic acid to be inserted into an expression vector so that the individual codons for each amino acid are those preferentially utilized in *E. coli* (Wada *et al.*, 1992, *Nucleic Acids Res.* 20:2111-2118).
- 30 Such alteration of nucleic acid sequences of the invention can be carried out by standard DNA synthesis techniques.

In another embodiment, the expression vector is a yeast expression vector. Examples of vectors for expression in yeast *S. cerevisiae* include pYepSec1 (Baldari *et al.*, 1987, *EMBO J.* 6:229-234), pMFa (Kurjan and Herskowitz, 1982, *Cell* 30:933-943), pJRY88 (Schultz *et al.*, 1987, *Gene* 54:113-123), pYES2 (Invitrogen Corporation, San Diego, CA), and pPicZ (Invitrogen Corp, San Diego, CA).

Alternatively, the expression vector is a baculovirus expression vector. Baculovirus vectors available for expression of proteins in cultured insect cells (*e.g.*, Sf 9 cells) include the pAc series (Smith *et al.*, 1983, *Mol. Cell Biol.* 3:2156-2165) and the pVL series (Lucklow and Summers, 1989, *Virology* 170:31-39).

In yet another embodiment, a nucleic acid of the invention is expressed in mammalian cells using a mammalian expression vector. Examples of mammalian expression vectors include pCDM8 (Seed, 1987, *Nature* 329:840) and pMT2PC (Kaufman *et al.*, 1987, *EMBO J.* 6:187-195). When used in mammalian cells, the expression vector's control functions are often provided by viral regulatory elements. For example, commonly used promoters are derived from polyoma, Adenovirus 2, cytomegalovirus and Simian Virus 40. For other suitable expression systems for both prokaryotic and eukaryotic cells see chapters 16 and 17 of Sambrook *et al.*, *supra*.

In another embodiment, the recombinant mammalian expression vector is capable of directing expression of the nucleic acid preferentially in a particular cell type (*e.g.*, tissue-specific regulatory elements are used to express the nucleic acid). Tissue-specific regulatory elements are known in the art. Non-limiting examples of suitable tissue-specific promoters include the albumin promoter (liver-specific; Pinkert *et al.*, 1987, *Genes Dev.* 1:268-277), lymphoid-specific promoters (Calame and Eaton, 1988, *Adv. Immunol.* 43:235-275), in particular promoters of T cell receptors (Winoto and Baltimore, 1989, *EMBO J.* 8:729-733) and immunoglobulins (Banerji *et al.*, 1983, *Cell* 33:729-740; Queen and Baltimore, 1983, *Cell* 33:741-748), neuron-specific promoters (*e.g.*, the neurofilament promoter; Byrne and Ruddle, 1989, *Proc. Natl. Acad. Sci. USA* 86:5473-5477), pancreas-specific promoters (Edlund *et al.*, 1985, *Science* 230:912-916), and mammary gland-specific promoters (*e.g.*, milk whey promoter; U.S. Patent No. 4,873,316 and European Application Publication No. 264,166). Developmentally-regulated promoters are also encompassed, for example the murine *hox* promoters

(Kessel and Gruss, 1990, *Science* 249:374-379) and the α -fetoprotein promoter (Camper and Tilghman, 1989, *Genes Dev.* 3:537-546).

The invention further provides a recombinant expression vector comprising a DNA molecule of the invention cloned into the expression vector in an antisense orientation. That is, the DNA molecule is operably linked to a regulatory sequence in a manner which allows for expression (by transcription of the DNA molecule) of an RNA molecule which is antisense to the mRNA encoding a polypeptide of the invention. Regulatory sequences operably linked to a nucleic acid cloned in the antisense orientation can be chosen which direct the continuous expression of the antisense RNA molecule in a variety of cell types, for instance viral promoters and/or enhancers, or regulatory sequences can be chosen which direct constitutive, tissue-specific or cell type specific expression of antisense RNA. The antisense expression vector can be in the form of a recombinant plasmid, phagemid, or attenuated virus in which antisense nucleic acids are produced under the control of a high efficiency regulatory region, the activity of which can be determined by the cell type into which the vector is introduced. For a discussion of the regulation of gene expression using antisense genes see Weintraub *et al.*, 1986, *Trends in Genetics*, Vol. 1(1).

Another aspect of the invention pertains to host cells into which a recombinant expression vector of the invention has been introduced. The terms "host cell" and "recombinant host cell" are used interchangeably herein. It is understood that such terms refer not only to the particular subject cell but to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

A host cell can be any prokaryotic (*e.g.*, *E. coli*) or eukaryotic cell (*e.g.*, insect cells, yeast or mammalian cells).

Vector DNA can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. As used herein, the terms "transformation" and "transfection" are intended to refer to a variety of art-recognized techniques for introducing foreign nucleic acid into a host cell, including calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection,

- 69 -

lipofection, or electroporation. Suitable methods for transforming or transfecting host cells can be found in Sambrook, *et al.* (*supra*), and other laboratory manuals.

For stable transfection of mammalian cells, it is known that, depending upon the expression vector and transfection technique used, only a small fraction of cells may integrate the foreign DNA into their genome. In order to identify and select these integrants, a gene that encodes a selectable marker (*e.g.*, for resistance to antibiotics) is generally introduced into the host cells along with the gene of interest. Preferred selectable markers include those which confer resistance to drugs, such as G418, hygromycin and methotrexate. Cells stably transfected with the introduced nucleic acid can be identified by drug selection (*e.g.*, cells that have incorporated the selectable marker gene will survive, while the other cells die).

A host cell of the invention, such as a prokaryotic or eukaryotic host cell in culture, can be used to produce a marker protein or a segment thereof. Accordingly, the invention further provides methods for producing a marker protein or a segment thereof using the host cells of the invention. In one embodiment, the method comprises culturing the host cell of the invention (into which a recombinant expression vector encoding a marker protein or a segment thereof has been introduced) in a suitable medium such that the is produced. In another embodiment, the method further comprises isolating the a marker protein or a segment thereof from the medium or the host cell.

The host cells of the invention can also be used to produce nonhuman transgenic animals. For example, in one embodiment, a host cell of the invention is a fertilized oocyte or an embryonic stem cell into which a sequences encoding a marker protein or a segment thereof have been introduced. Such host cells can then be used to create non-human transgenic animals in which exogenous sequences encoding a marker protein of the invention have been introduced into their genome or homologous recombinant animals in which endogenous gene(s) encoding a marker protein have been altered. Such animals are useful for studying the function and/or activity of the marker protein and for identifying and/or evaluating modulators of marker protein. As used herein, a "transgenic animal" is a non-human animal, preferably a mammal, more preferably a rodent such as a rat or mouse, in which one or more of the cells of the animal includes a transgene. Other examples of transgenic animals include non-human

- 70 -

primates, sheep, dogs, cows, goats, chickens, amphibians, etc. A transgene is exogenous DNA which is integrated into the genome of a cell from which a transgenic animal develops and which remains in the genome of the mature animal, thereby directing the expression of an encoded gene product in one or more cell types or tissues of the transgenic animal. As used herein, an "homologous recombinant animal" is a non-human animal, preferably a mammal, more preferably a mouse, in which an endogenous gene has been altered by homologous recombination between the endogenous gene and an exogenous DNA molecule introduced into a cell of the animal, *e.g.*, an embryonic cell of the animal, prior to development of the animal.

A transgenic animal of the invention can be created by introducing a nucleic acid encoding a marker protein into the male pronuclei of a fertilized oocyte, *e.g.*, by microinjection, retroviral infection, and allowing the oocyte to develop in a pseudopregnant female foster animal. Intronic sequences and polyadenylation signals can also be included in the transgene to increase the efficiency of expression of the transgene. A tissue-specific regulatory sequence(s) can be operably linked to the transgene to direct expression of the polypeptide of the invention to particular cells. Methods for generating transgenic animals via embryo manipulation and microinjection, particularly animals such as mice, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866 and 4,870,009, U.S. Patent No. 4,873,191 and in Hogan, *Manipulating the Mouse Embryo*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1986. Similar methods are used for production of other transgenic animals. A transgenic founder animal can be identified based upon the presence of the transgene in its genome and/or expression of mRNA encoding the transgene in tissues or cells of the animals. A transgenic founder animal can then be used to breed additional animals carrying the transgene. Moreover, transgenic animals carrying the transgene can further be bred to other transgenic animals carrying other transgenes.

To create an homologous recombinant animal, a vector is prepared which contains at least a portion of a gene encoding a marker protein into which a deletion, addition or substitution has been introduced to thereby alter, *e.g.*, functionally disrupt, the gene. In a preferred embodiment, the vector is designed such that, upon homologous recombination, the endogenous gene is functionally disrupted (*i.e.*, no longer encodes a

functional protein; also referred to as a "knock out" vector). Alternatively, the vector can be designed such that, upon homologous recombination, the endogenous gene is mutated or otherwise altered but still encodes functional protein (e.g., the upstream regulatory region can be altered to thereby alter the expression of the endogenous protein). In the homologous recombination vector, the altered portion of the gene is flanked at its 5' and 3' ends by additional nucleic acid of the gene to allow for homologous recombination to occur between the exogenous gene carried by the vector and an endogenous gene in an embryonic stem cell. The additional flanking nucleic acid sequences are of sufficient length for successful homologous recombination with the endogenous gene. Typically, several kilobases of flanking DNA (both at the 5' and 3' ends) are included in the vector (see, e.g., Thomas and Capecchi, 1987, *Cell* 51:503 for a description of homologous recombination vectors). The vector is introduced into an embryonic stem cell line (e.g., by electroporation) and cells in which the introduced gene has homologously recombined with the endogenous gene are selected (see, e.g., Li et al., 1992, *Cell* 69:915). The selected cells are then injected into a blastocyst of an animal (e.g., a mouse) to form aggregation chimeras (see, e.g., Bradley, *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, Robertson, Ed., IRL, Oxford, 1987, pp. 113-152). A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term. Progeny harboring the homologously recombined DNA in their germ cells can be used to breed animals in which all cells of the animal contain the homologously recombined DNA by germline transmission of the transgene. Methods for constructing homologous recombination vectors and homologous recombinant animals are described further in Bradley (1991) *Current Opinion in Bio/Technology* 2:823-829 and in PCT Publication NOS. WO 90/11354, WO 91/01140, WO 92/0968, and WO 93/04169.

In another embodiment, transgenic non-human animals can be produced which contain selected systems which allow for regulated expression of the transgene. One example of such a system is the *cre/loxP* recombinase system of bacteriophage P1. For a description of the *cre/loxP* recombinase system, see, e.g., Lakso et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:6232-6236. Another example of a recombinase system is the FLP recombinase system of *Saccharomyces cerevisiae* (O'Gorman et al., 1991, *Science* 251:1351-1355). If a *cre/loxP* recombinase system is used to regulate expression of the

- 72 -

transgene, animals containing transgenes encoding both the *Cre* recombinase and a selected protein are required. Such animals can be provided through the construction of "double" transgenic animals, *e.g.*, by mating two transgenic animals, one containing a transgene encoding a selected protein and the other containing a transgene encoding a recombinase.

Clones of the non-human transgenic animals described herein can also be produced according to the methods described in Wilmot *et al.* (1997) *Nature* 385:810-813 and PCT Publication NOS. WO 97/07668 and WO 97/07669.

10 IV. Pharmaceutical Compositions

The nucleic acid molecules, polypeptides, and antibodies (also referred to herein as "active compounds") of the invention can be incorporated into pharmaceutical compositions suitable for administration. Such compositions typically comprise the nucleic acid molecule, protein, or antibody and a pharmaceutically acceptable carrier.

15 As used herein the language "pharmaceutically acceptable carrier" is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

The invention includes methods for preparing pharmaceutical compositions for modulating the expression or activity of a marker nucleic acid or protein. Such methods comprise formulating a pharmaceutically acceptable carrier with an agent which modulates expression or activity of a marker nucleic acid or protein. Such compositions can further include additional active agents. Thus, the invention further includes methods for preparing a pharmaceutical composition by formulating a pharmaceutically acceptable carrier with an agent which modulates expression or activity of a marker nucleic acid or protein and one or more additional active compounds.

The invention also provides methods (also referred to herein as "screening assays") for identifying modulators, *i.e.*, candidate or test compounds or agents (*e.g.*, peptides, peptidomimetics, peptoids, small molecules or other drugs) which (a) bind to the marker, or (b) have a modulatory (*e.g.*, stimulatory or inhibitory) effect on the activity of the marker or, more specifically, (c) have a modulatory effect on the interactions of the marker with one or more of its natural substrates (*e.g.*, peptide, protein, hormone, co-factor, or nucleic acid), or (d) have a modulatory effect on the expression of the marker. Such assays typically comprise a reaction between the marker and one or more assay components. The other components may be either the test compound itself, or a combination of test compound and a natural binding partner of the marker.

The test compounds of the present invention may be obtained from any available source, including systematic libraries of natural and/or synthetic compounds. Test compounds may also be obtained by any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; peptoid libraries (libraries of molecules having the functionalities of peptides, but with a novel, non-peptide backbone which are resistant to enzymatic degradation but which nevertheless remain bioactive; see, *e.g.*, Zuckermann *et al.*, 1994, *J. Med. Chem.* 37:2678-85); spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the 'one-bead one-compound' library method; and synthetic library methods using affinity chromatography selection. The biological library and peptoid library approaches are limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds (Lam, 1997, *Anticancer Drug Des.* 12:145).

Examples of methods for the synthesis of molecular libraries can be found in the art, for example in: DeWitt *et al.* (1993) *Proc. Natl. Acad. Sci. U.S.A.* 90:6909; Erb *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:11422; Zuckermann *et al.* (1994). *J. Med. Chem.* 37:2678; Cho *et al.* (1993) *Science* 261:1303; Carrell *et al.* (1994) *Angew. Chem. Int. Ed. Engl.* 33:2059; Carell *et al.* (1994) *Angew. Chem. Int. Ed. Engl.* 33:2061; and in Gallop *et al.* (1994) *J. Med. Chem.* 37:1233.

Libraries of compounds may be presented in solution (*e.g.*, Houghten, 1992, *Biotechniques* 13:412-421), or on beads (Lam, 1991, *Nature* 354:82-84), chips (Fodor, 1993, *Nature* 364:555-556), bacteria and/or spores, (Ladner, USP 5,223,409), plasmids (Cull *et al.*, 1992, *Proc Natl Acad Sci USA* 89:1865-1869) or on phage (Scott and Smith, 1990, *Science* 249:386-390; Devlin, 1990, *Science* 249:404-406; Cwirla *et al.*, 1990, *Proc. Natl. Acad. Sci.* 87:6378-6382; Felici, 1991, *J. Mol. Biol.* 222:301-310; Ladner, *supra.*).

In one embodiment, the invention provides assays for screening candidate or test compounds which are substrates of a protein encoded by or corresponding to a marker or biologically active portion thereof. In another embodiment, the invention provides assays for screening candidate or test compounds which bind to a protein encoded by or corresponding to a marker or biologically active portion thereof. Determining the ability of the test compound to directly bind to a protein can be accomplished, for example, by coupling the compound with a radioisotope or enzymatic label such that binding of the compound to the marker can be determined by detecting the labeled marker compound in a complex. For example, compounds (*e.g.*, marker substrates) can be labeled with ^{125}I , ^{35}S , ^{14}C , or ^3H , either directly or indirectly, and the radioisotope detected by direct counting of radioemission or by scintillation counting. Alternatively, assay components can be enzymatically labeled with, for example, horseradish peroxidase, alkaline phosphatase, or luciferase, and the enzymatic label detected by determination of conversion of an appropriate substrate to product.

In another embodiment, the invention provides assays for screening candidate or test compounds which modulate the expression of a marker or the activity of a protein encoded by or corresponding to a marker, or a biologically active portion thereof. In all likelihood, the protein encoded by or corresponding to the marker can, *in vivo*, interact with one or more molecules, such as but not limited to, peptides, proteins, hormones, cofactors and nucleic acids. For the purposes of this discussion, such cellular and extracellular molecules are referred to herein as "binding partners" or marker "substrate".

One necessary embodiment of the invention in order to facilitate such screening is the use of a protein encoded by or corresponding to marker to identify the protein's natural *in vivo* binding partners. There are many ways to accomplish this which are known to one skilled in the art. One example is the use of the marker protein as "bait protein" in a two-hybrid assay or three-hybrid assay (see, *e.g.*, U.S. Patent No. 5,283,317; Zervos *et al*, 1993, *Cell* 72:223-232; Madura *et al*, 1993, *J. Biol. Chem.* 268:12046-12054; Bartel *et al*, 1993, *Biotechniques* 14:920-924; Iwabuchi *et al*, 1993 *Oncogene* 8:1693-1696; Brent WO94/10300) in order to identify other proteins which bind to or interact with the marker (binding partners) and, therefore, are possibly involved in the natural function of the marker. Such marker binding partners are also likely to be involved in the propagation of signals by the marker protein or downstream elements of a marker protein-mediated signaling pathway. Alternatively, such marker protein binding partners may also be found to be inhibitors of the marker protein.

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. In one construct, the gene that encodes a marker protein fused to a gene encoding the DNA binding domain of a known transcription factor (*e.g.*, GAL-4). In the other construct, a DNA sequence, from a library of DNA sequences, that encodes an unidentified protein ("prey" or "sample") is fused to a gene that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact, *in vivo*, forming a marker-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (*e.g.*, LacZ) which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be readily detected and cell colonies containing the functional transcription factor can be isolated and used to obtain the cloned gene which encodes the protein which interacts with the marker protein.

In a further embodiment, assays may be devised through the use of the invention for the purpose of identifying compounds which modulate (*e.g.*, affect either positively or negatively) interactions between a marker protein and its substrates and/or binding partners. Such compounds can include, but are not limited to, molecules such as antibodies, peptides, hormones, oligonucleotides, nucleic acids, and analogs thereof.

- 76 -

Such compounds may also be obtained from any available source, including systematic libraries of natural and/or synthetic compounds. The preferred assay components for use in this embodiment is an ovarian cancer marker protein identified herein, the known binding partner and/or substrate of same, and the test compound. Test compounds can be
5 supplied from any source.

The basic principle of the assay systems used to identify compounds that interfere with the interaction between the marker protein and its binding partner involves preparing a reaction mixture containing the marker protein and its binding partner under conditions and for a time sufficient to allow the two products to interact
10 and bind, thus forming a complex. In order to test an agent for inhibitory activity, the reaction mixture is prepared in the presence and absence of the test compound. The test compound can be initially included in the reaction mixture, or can be added at a time subsequent to the addition of the marker protein and its binding partner. Control reaction mixtures are incubated without the test compound or with a placebo. The
15 formation of any complexes between the marker protein and its binding partner is then detected. The formation of a complex in the control reaction, but less or no such formation in the reaction mixture containing the test compound, indicates that the compound interferes with the interaction of the marker protein and its binding partner. Conversely, the formation of more complex in the presence of compound than in the
20 control reaction indicates that the compound may enhance interaction of the marker protein and its binding partner.

The assay for compounds that interfere with the interaction of the marker protein with its binding partner may be conducted in a heterogeneous or homogeneous format. Heterogeneous assays involve anchoring either the marker protein or its binding
25 partner onto a solid phase and detecting complexes anchored to the solid phase at the end of the reaction. In homogeneous assays, the entire reaction is carried out in a liquid phase. In either approach, the order of addition of reactants can be varied to obtain different information about the compounds being tested. For example, test compounds that interfere with the interaction between the marker proteins and the binding partners
30 (e.g., by competition) can be identified by conducting the reaction in the presence of the test substance, i.e., by adding the test substance to the reaction mixture prior to or simultaneously with the marker and its interactive binding partner. Alternatively, test

- 77 -

compounds that disrupt preformed complexes, *e.g.*, compounds with higher binding constants that displace one of the components from the complex, can be tested by adding the test compound to the reaction mixture after complexes have been formed. The various formats are briefly described below.

5 In a heterogeneous assay system, either the marker protein or its binding partner is anchored onto a solid surface or matrix, while the other corresponding non-anchored component may be labeled, either directly or indirectly. In practice, microtitre plates are often utilized for this approach. The anchored species can be immobilized by a number of methods, either non-covalent or covalent, that are typically well known to
10 one who practices the art. Non-covalent attachment can often be accomplished simply by coating the solid surface with a solution of the marker protein or its binding partner and drying. Alternatively, an immobilized antibody specific for the assay component to be anchored can be used for this purpose. Such surfaces can often be prepared in
advance and stored.

15 In related embodiments, a fusion protein can be provided which adds a domain that allows one or both of the assay components to be anchored to a matrix. For example, glutathione-S-transferase/marker fusion proteins or glutathione-S-transferase/binding partner can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtiter plates, which are then
20 combined with the test compound or the test compound and either the non-adsorbed marker or its binding partner, and the mixture incubated under conditions conducive to complex formation (*e.g.*, physiological conditions). Following incubation, the beads or microtiter plate wells are washed to remove any unbound assay components, the immobilized complex assessed either directly or indirectly, for example, as described
25 above. Alternatively, the complexes can be dissociated from the matrix, and the level of marker binding or activity determined using standard techniques.

Other techniques for immobilizing proteins on matrices can also be used in the screening assays of the invention. For example, either a marker protein or a marker protein binding partner can be immobilized utilizing conjugation of biotin and
30 streptavidin. Biotinylated marker protein or target molecules can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques known in the art (*e.g.*, biotinylation kit, Pierce Chemicals, Rockford, IL), and immobilized in the wells of

streptavidin-coated 96 well plates (Pierce Chemical). In certain embodiments, the protein-immobilized surfaces can be prepared in advance and stored.

- In order to conduct the assay, the corresponding partner of the immobilized assay component is exposed to the coated surface with or without the test compound. After the reaction is complete, unreacted assay components are removed (e.g., by washing) and any complexes formed will remain immobilized on the solid surface. The detection of complexes anchored on the solid surface can be accomplished in a number of ways. Where the non-immobilized component is pre-labeled, the detection of label immobilized on the surface indicates that complexes were formed.
- Where the non-immobilized component is not pre-labeled, an indirect label can be used to detect complexes anchored on the surface; e.g., using a labeled antibody specific for the initially non-immobilized species (the antibody, in turn, can be directly labeled or indirectly labeled with, e.g., a labeled anti-Ig antibody). Depending upon the order of addition of reaction components, test compounds which modulate (inhibit or enhance) complex formation or which disrupt preformed complexes can be detected.

- In an alternate embodiment of the invention, a homogeneous assay may be used. This is typically a reaction, analogous to those mentioned above, which is conducted in a liquid phase in the presence or absence of the test compound. The formed complexes are then separated from unreacted components, and the amount of complex formed is determined. As mentioned for heterogeneous assay systems, the order of addition of reactants to the liquid phase can yield information about which test compounds modulate (inhibit or enhance) complex formation and which disrupt preformed complexes.

- In such a homogeneous assay, the reaction products may be separated from unreacted assay components by any of a number of standard techniques, including but not limited to: differential centrifugation, chromatography, electrophoresis and immunoprecipitation. In differential centrifugation, complexes of molecules may be separated from uncomplexed molecules through a series of centrifugal steps, due to the different sedimentation equilibria of complexes based on their different sizes and densities (see, for example, Rivas, G., and Minton, A.P., *Trends Biochem Sci* 1993 Aug;18(8):284-7). Standard chromatographic techniques may also be utilized to separate complexed molecules from uncomplexed ones. For example, gel filtration

- 79 -

chromatography separates molecules based on size, and through the utilization of an appropriate gel filtration resin in a column format, for example, the relatively larger complex may be separated from the relatively smaller uncomplexed components. Similarly, the relatively different charge properties of the complex as compared to the uncomplexed molecules may be exploited to differentially separate the complex from the remaining individual reactants, for example through the use of ion-exchange chromatography resins. Such resins and chromatographic techniques are well known to one skilled in the art (see, e.g., Heegaard, 1998, *J Mol. Recognit.* 11:141-148; Hage and Tweed, 1997, *J. Chromatogr. B. Biomed. Sci. Appl.*, 699:499-525). Gel electrophoresis may also be employed to separate complexed molecules from unbound species (see, e.g., Ausubel *et al* (eds.), as described in : Current Protocols in Molecular Biology, J. Wiley & Sons, New York. 1999). In this technique, protein or nucleic acid complexes are separated based on size or charge, for example. In order to maintain the binding interaction during the electrophoretic process, nondenaturing gels in the absence of reducing agent are typically preferred, but conditions appropriate to the particular interactants will be well known to one skilled in the art. Immunoprecipitation is another common technique utilized for the isolation of a protein-protein complex from solution (see, e.g., Ausubel *et al* (eds.), In: Current Protocols in Molecular Biology, J. Wiley & Sons, New York. 1999). In this technique, all proteins binding to an antibody specific to one of the binding molecules are precipitated from solution by conjugating the antibody to a polymer bead that may be readily collected by centrifugation. The bound assay components are released from the beads (through a specific proteolysis event or other technique well known in the art which will not disturb the protein-protein interaction in the complex), and a second immunoprecipitation step is performed, this time utilizing antibodies specific for the correspondingly different interacting assay component. In this manner, only formed complexes should remain attached to the beads. Variations in complex formation in both the presence and the absence of a test compound can be compared, thus offering information about the ability of the compound to modulate interactions between the marker protein and its binding partner.

Also within the scope of the present invention are methods for direct detection of interactions between the marker protein and its natural binding partner and/or a test compound in a homogeneous or heterogeneous assay system without

- 80 -

further sample manipulation. For example, the technique of fluorescence energy transfer may be utilized (see, *e.g.*, Lakowicz *et al.*, U.S. Patent No. 5,631,169; Stavrianopoulos *et al.*, U.S. Patent No. 4,868,103). Generally, this technique involves the addition of a fluorophore label on a first 'donor' molecule (*e.g.*, marker or test compound) such that its emitted fluorescent energy will be absorbed by a fluorescent label on a second, 'acceptor' molecule (*e.g.*, marker or test compound), which in turn is able to fluoresce due to the absorbed energy. Alternately, the 'donor' protein molecule may simply utilize the natural fluorescent energy of tryptophan residues. Labels are chosen that emit different wavelengths of light, such that the 'acceptor' molecule label may be differentiated from that of the 'donor'. Since the efficiency of energy transfer between the labels is related to the distance separating the molecules, spatial relationships between the molecules can be assessed. In a situation in which binding occurs between the molecules, the fluorescent emission of the 'acceptor' molecule label in the assay should be maximal. An FET binding event can be conveniently measured through standard fluorometric detection means well known in the art (*e.g.*, using a fluorimeter). A test substance which either enhances or hinders participation of one of the species in the preformed complex will result in the generation of a signal variant to that of background. In this way, test substances that modulate interactions between a marker and its binding partner can be identified in controlled assays.

In another embodiment, modulators of marker expression are identified in a method wherein a cell is contacted with a candidate compound and the expression of marker mRNA or protein in the cell, is determined. The level of expression of marker mRNA or protein in the presence of the candidate compound is compared to the level of expression of marker mRNA or protein in the absence of the candidate compound. The candidate compound can then be identified as a modulator of marker expression based on this comparison. For example, when expression of marker mRNA or protein is greater (statistically significantly greater) in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of marker mRNA or protein expression. Conversely, when expression of marker mRNA or protein is less (statistically significantly less) in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of marker mRNA or protein expression. The level of marker mRNA or protein expression

in the cells can be determined by methods described herein for detecting marker mRNA or protein.

In another aspect, the invention pertains to a combination of two or more of the assays described herein. For example, a modulating agent can be identified using
5 a cell-based or a cell free assay, and the ability of the agent to modulate the activity of a marker protein can be further confirmed *in vivo*, *e.g.*, in a whole animal model for cellular transformation and/or tumorigenesis.

This invention further pertains to novel agents identified by the above-described screening assays. Accordingly, it is within the scope of this invention to
10 further use an agent identified as described herein in an appropriate animal model. For example, an agent identified as described herein (*e.g.*, an marker modulating agent, an antisense marker nucleic acid molecule, an marker-specific antibody, or an marker-binding partner) can be used in an animal model to determine the efficacy, toxicity, or side effects of treatment with such an agent. Alternatively, an agent identified as
15 described herein can be used in an animal model to determine the mechanism of action of such an agent. Furthermore, this invention pertains to uses of novel agents identified by the above-described screening assays for treatments as described herein.

It is understood that appropriate doses of small molecule agents and
protein or polypeptide agents depends upon a number of factors within the knowledge of
20 the ordinarily skilled physician, veterinarian, or researcher. The dose(s) of these agents will vary, for example, depending upon the identity, size, and condition of the subject or sample being treated, further depending upon the route by which the composition is to be administered, if applicable, and the effect which the practitioner desires the agent to have upon the nucleic acid or polypeptide of the invention. Exemplary doses of a small
25 molecule include milligram or microgram amounts per kilogram of subject or sample weight (*e.g.* about 1 microgram per kilogram to about 500 milligrams per kilogram, about 100 micrograms per kilogram to about 5 milligrams per kilogram, or about 1 microgram per kilogram to about 50 micrograms per kilogram). Exemplary doses of a protein or polypeptide include gram, milligram or microgram amounts per kilogram of
30 subject or sample weight (*e.g.* about 1 microgram per kilogram to about 5 grams per kilogram, about 100 micrograms per kilogram to about 500 milligrams per kilogram, or about 1 milligram per kilogram to about 50 milligrams per kilogram). It is furthermore

understood that appropriate doses of one of these agents depend upon the potency of the agent with respect to the expression or activity to be modulated. Such appropriate doses can be determined using the assays described herein. When one or more of these agents is to be administered to an animal (*e.g.* a human) in order to modulate expression or
5 activity of a polypeptide or nucleic acid of the invention, a physician, veterinarian, or researcher can, for example, prescribe a relatively low dose at first, subsequently increasing the dose until an appropriate response is obtained. In addition, it is understood that the specific dose level for any particular animal subject will depend upon a variety of factors including the activity of the specific agent employed, the age,
10 body weight, general health, gender, and diet of the subject, the time of administration, the route of administration, the rate of excretion, any drug combination, and the degree of expression or activity to be modulated.

A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of
15 administration include parenteral, *e.g.*, intravenous, intradermal, subcutaneous, oral (*e.g.*, inhalation), transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other
20 synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediamine-tetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The
25 parenteral preparation can be enclosed in ampules, disposable syringes or multiple dose vials made of glass or plastic.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. For
30 intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL (BASF; Parsippany, NJ) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy

- 83 -

- syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, or sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.
- 15 Sterile injectable solutions can be prepared by incorporating the active compound (e.g., a polypeptide or antibody) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium, and then incorporating the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.
- 25 Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed.
- 30

- 84 -

Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches, and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

For administration by inhalation, the compounds are delivered in the form of an aerosol spray from a pressurized container or dispenser which contains a suitable propellant, *e.g.*, a gas such as carbon dioxide, or a nebulizer.

Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

The compounds can also be prepared in the form of suppositories (*e.g.*, with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes having monoclonal antibodies incorporated therein or thereon) can also be used as pharmaceutically

acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811.

It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit
15 form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active compound
10 and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

For antibodies, the preferred dosage is 0.1 mg/kg to 100 mg/kg of body weight (generally 10 mg/kg to 20 mg/kg). If the antibody is to act in the brain, a dosage of 50 mg/kg to 100 mg/kg is usually appropriate. Generally, partially human antibodies
15 and fully human antibodies have a longer half-life within the human body than other antibodies. Accordingly, lower dosages and less frequent administration is often possible. Modifications such as lipidation can be used to stabilize antibodies and to enhance uptake and tissue penetration (*e.g.*, into the ovarian epithelium). A method for lipidation of antibodies is described by Cruikshank *et al.* (1997) *J. Acquired Immune*
20 *Deficiency Syndromes and Human Retrovirology* 14:193.

The invention also provides vaccine compositions for the prevention and/or treatment of ovarian cancer. The invention provides ovarian cancer vaccine compositions in which a protein of a marker of Table 1, or a combination of proteins of the markers of Table 1, are introduced into a subject in order to stimulate an immune
25 response against the ovarian cancer. The invention also provides ovarian cancer vaccine compositions in which a gene expression construct, which expresses a marker or fragment of a marker identified in Table 1, is introduced into the subject such that a protein or fragment of a protein encoded by a marker of Table 1 is produced by transfected cells in the subject at a higher than normal level and elicits an immune
30 response.

In one embodiment, an ovarian cancer vaccine is provided and employed as an immunotherapeutic agent for the prevention of ovarian cancer. In another embodiment, an ovarian cancer vaccine is provided and employed as an immunotherapeutic agent for the treatment of ovarian cancer.

- 5 By way of example, an ovarian cancer vaccine comprised of the proteins of the markers of Table 1, may be employed for the prevention and/or treatment of ovarian cancer in a subject by administering the vaccine by a variety of routes, *e.g.*, intradermally, subcutaneously, or intramuscularly. In addition, the ovarian cancer vaccine can be administered together with adjuvants and/or immunomodulators to boost
10 the activity of the vaccine and the subject's response. In one embodiment, devices and/or compositions containing the vaccine, suitable for sustained or intermittent release could be, implanted in the body or topically applied thereto for the relatively slow release of such materials into the body. The ovarian cancer vaccine can be introduced along with immunomodulatory compounds, which can alter the type of immune
15 response produced in order to produce a response which will be more effective in eliminating the cancer.

- In another embodiment, an ovarian cancer vaccine comprised of an expression construct of the markers of Table 1, may be introduced by injection into muscle or by coating onto microprojectiles and using a device designed for the purpose
20 to fire the projectiles at high speed into the skin. The cells of the subject will then express the protein(s) or fragments of proteins of the markers of Table 1 and induce an immune response. In addition, the ovarian cancer vaccine may be introduced along with expression constructs for immunomodulatory molecules, such as cytokines, which may increase the immune response or modulate the type of immune response produced in
25 order to produce a response which will be more effective in eliminating the cancer.

- The marker nucleic acid molecules of the present invention can also be inserted into vectors and used as gene therapy vectors. Gene therapy vectors can be delivered to a subject by, for example, intravenous injection, local administration (U.S. Patent 5,328,470), or by stereotactic injection (see, *e.g.*, Chen *et al.*, 1994, *Proc. Natl. Acad. Sci. USA* 91:3054-3057). The pharmaceutical preparation of the gene therapy
30 vector can include the gene therapy vector in an acceptable diluent, or can comprise a slow release matrix in which the gene delivery vehicle is imbedded. Alternatively,

- 87 -

where the complete gene delivery vector can be produced intact from recombinant cells, *e.g.* retroviral vectors, the pharmaceutical preparation can include one or more cells which produce the gene delivery system.

The pharmaceutical compositions can be included in a container, pack, or
5 dispenser together with instructions for administration.

V. Predictive Medicine

The present invention pertains to the field of predictive medicine in which diagnostic assays, prognostic assays, pharmacogenomics, and monitoring clinical
10 trails are used for prognostic (predictive) purposes to thereby treat an individual prophylactically. Accordingly, one aspect of the present invention relates to diagnostic assays for determining the level of expression of one or more marker proteins or nucleic acids, in order to determine whether an individual is at risk of developing ovarian cancer. Such assays can be used for prognostic or predictive purposes to thereby
15 prophylactically treat an individual prior to the onset of the cancer.

Yet another aspect of the invention pertains to monitoring the influence of agents (*e.g.*, drugs or other compounds administered either to inhibit ovarian cancer or to treat or prevent any other disorder {*i.e.* in order to understand any ovarian carcinogenic effects that such treatment may have}) on the expression or activity of a
20 marker of the invention in clinical trials. These and other agents are described in further detail in the following sections.

A. Diagnostic Assays

An exemplary method for detecting the presence or absence of a marker
25 protein or nucleic acid in a biological sample involves obtaining a biological sample (*e.g.* an ovary-associated body fluid) from a test subject and contacting the biological sample with a compound or an agent capable of detecting the polypeptide or nucleic acid (*e.g.*, mRNA, genomic DNA, or cDNA). The detection methods of the invention can thus be used to detect mRNA, protein, cDNA, or genomic DNA, for example, in a
30 biological sample *in vitro* as well as *in vivo*. For example, *in vitro* techniques for detection of mRNA include Northern hybridizations and *in situ* hybridizations. *In vitro* techniques for detection of a marker protein include enzyme linked immunosorbent

assays (ELISAs), Western blots, immunoprecipitations and immunofluorescence. *In vitro* techniques for detection of genomic DNA include Southern hybridizations. Furthermore, *in vivo* techniques for detection of a marker protein include introducing into a subject a labeled antibody directed against the protein or fragment thereof. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques.

A general principle of such diagnostic and prognostic assays involves preparing a sample or reaction mixture that may contain a marker, and a probe, under appropriate conditions and for a time sufficient to allow the marker and probe to interact and bind, thus forming a complex that can be removed and/or detected in the reaction mixture. These assays can be conducted in a variety of ways.

For example, one method to conduct such an assay would involve anchoring the marker or probe onto a solid phase support, also referred to as a substrate, and detecting target marker/probe complexes anchored on the solid phase at the end of the reaction. In one embodiment of such a method, a sample from a subject, which is to be assayed for presence and/or concentration of marker, can be anchored onto a carrier or solid phase support. In another embodiment, the reverse situation is possible, in which the probe can be anchored to a solid phase and a sample from a subject can be allowed to react as an unanchored component of the assay.

There are many established methods for anchoring assay components to a solid phase. These include, without limitation, marker or probe molecules which are immobilized through conjugation of biotin and streptavidin. Such biotinylated assay components can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques known in the art (*e.g.*, biotinylation kit, Pierce Chemicals, Rockford, IL), and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). In certain embodiments, the surfaces with immobilized assay components can be prepared in advance and stored.

Other suitable carriers or solid phase supports for such assays include any material capable of binding the class of molecule to which the marker or probe belongs. Well-known supports or carriers include, but are not limited to, glass, polystyrene, nylon, polypropylene, nylon, polyethylene, dextran, amylases, natural and modified celluloses, polyacrylamides, gabbros, and magnetite.

In order to conduct assays with the above mentioned approaches, the non-immobilized component is added to the solid phase upon which the second component is anchored. After the reaction is complete, uncomplexed components may be removed (*e.g.*, by washing) under conditions such that any complexes formed will remain immobilized upon the solid phase. The detection of marker/probe complexes anchored to the solid phase can be accomplished in a number of methods outlined herein.

In a preferred embodiment, the probe, when it is the unanchored assay component, can be labeled for the purpose of detection and readout of the assay, either directly or indirectly, with detectable labels discussed herein and which are well-known to one skilled in the art.

It is also possible to directly detect marker/probe complex formation without further manipulation or labeling of either component (marker or probe), for example by utilizing the technique of fluorescence energy transfer (see, for example, Lakowicz *et al.*, U.S. Patent No. 5,631,169; Stavrianopoulos, *et al.*, U.S. Patent No. 4,868,103). A fluorophore label on the first, 'donor' molecule is selected such that, upon excitation with incident light of appropriate wavelength, its emitted fluorescent energy will be absorbed by a fluorescent label on a second 'acceptor' molecule, which in turn is able to fluoresce due to the absorbed energy. Alternately, the 'donor' protein molecule may simply utilize the natural fluorescent energy of tryptophan residues. Labels are chosen that emit different wavelengths of light, such that the 'acceptor' molecule label may be differentiated from that of the 'donor'. Since the efficiency of energy transfer between the labels is related to the distance separating the molecules, spatial relationships between the molecules can be assessed. In a situation in which binding occurs between the molecules, the fluorescent emission of the 'acceptor' molecule label in the assay should be maximal. An FET binding event can be conveniently measured through standard fluorometric detection means well known in the art (*e.g.*, using a fluorimeter).

In another embodiment, determination of the ability of a probe to recognize a marker can be accomplished without labeling either assay component (probe or marker) by utilizing a technology such as real-time Biomolecular Interaction Analysis (BIA) (see, *e.g.*, Sjolander, S. and Urbaniczky, C., 1991, *Anal. Chem.* 63:2338-2345

and Szabo *et al.*, 1995, *Curr. Opin. Struct. Biol.* 5:699-705). As used herein, "BIA" or "surface plasmon resonance" is a technology for studying biospecific interactions in real time, without labeling any of the interactants (*e.g.*, BIAcore). Changes in the mass at the binding surface (indicative of a binding event) result in alterations of the refractive index of light near the surface (the optical phenomenon of surface plasmon resonance (SPR)), resulting in a detectable signal which can be used as an indication of real-time reactions between biological molecules.

- Alternatively, in another embodiment, analogous diagnostic and prognostic assays can be conducted with marker and probe as solutes in a liquid phase.
- 10 In such an assay, the complexed marker and probe are separated from uncomplexed components by any of a number of standard techniques, including but not limited to: differential centrifugation, chromatography, electrophoresis and immunoprecipitation. In differential centrifugation, marker/probe complexes may be separated from uncomplexed assay components through a series of centrifugal steps, due to the different
- 15 sedimentation equilibria of complexes based on their different sizes and densities (see, for example, Rivas, G., and Minton, A.P., 1993, *Trends Biochem Sci.* 18(8):284-7). Standard chromatographic techniques may also be utilized to separate complexed molecules from uncomplexed ones. For example, gel filtration chromatography separates molecules based on size, and through the utilization of an appropriate gel
- 20 filtration resin in a column format, for example, the relatively larger complex may be separated from the relatively smaller uncomplexed components. Similarly, the relatively different charge properties of the marker/probe complex as compared to the uncomplexed components may be exploited to differentiate the complex from uncomplexed components, for example through the utilization of ion-exchange
- 25 chromatography resins. Such resins and chromatographic techniques are well known to one skilled in the art (see, *e.g.*, Heegaard, N.H., 1998, *J. Mol. Recognit.* Winter 11(1-6):141-8; Hage, D.S., and Tweed, S.A. *J Chromatogr B Biomed Sci Appl* 1997 Oct 10;699(1-2):499-525). Gel electrophoresis may also be employed to separate complexed assay components from unbound components (see, *e.g.*, Ausubel *et al.*, ed.,
- 30 *Current Protocols in Molecular Biology*, John Wiley & Sons, New York, 1987-1999). In this technique, protein or nucleic acid complexes are separated based on size or charge, for example. In order to maintain the binding interaction during the

electrophoretic process, non-denaturing gel matrix materials and conditions in the absence of reducing agent are typically preferred. Appropriate conditions to the particular assay and components thereof will be well known to one skilled in the art.

In a particular embodiment, the level of marker mRNA can be
5 determined both by *in situ* and by *in vitro* formats in a biological sample using methods known in the art. The term "biological sample" is intended to include tissues, cells, biological fluids and isolates thereof, isolated from a subject, as well as tissues, cells and fluids present within a subject. Many expression detection methods use isolated RNA. For *in vitro* methods, any RNA isolation technique that does not select against the
10 isolation of mRNA can be utilized for the purification of RNA from ovarian cells (see, e.g., Ausubel *et al.*, ed., *Current Protocols in Molecular Biology*, John Wiley & Sons, New York 1987-1999). Additionally, large numbers of tissue samples can readily be processed using techniques well known to those of skill in the art, such as, for example, the single-step RNA isolation process of Chomczynski (1989, U.S. Patent No.
15 4,843,155).

The isolated mRNA can be used in hybridization or amplification assays that include, but are not limited to, Southern or Northern analyses, polymerase chain reaction analyses and probe arrays. One preferred diagnostic method for the detection of mRNA levels involves contacting the isolated mRNA with a nucleic acid molecule
20 (probe) that can hybridize to the mRNA encoded by the gene being detected. The nucleic acid probe can be, for example, a full-length cDNA, or a portion thereof, such as an oligonucleotide of at least 7, 15, 30, 50, 100, 250 or 500 nucleotides in length and sufficient to specifically hybridize under stringent conditions to a mRNA or genomic DNA encoding a marker of the present invention. Other suitable probes for use in the
25 diagnostic assays of the invention are described herein. Hybridization of an mRNA with the probe indicates that the marker in question is being expressed.

In one format, the mRNA is immobilized on a solid surface and contacted with a probe, for example by running the isolated mRNA on an agarose gel and transferring the mRNA from the gel to a membrane, such as nitrocellulose. In an
30 alternative format, the probe(s) are immobilized on a solid surface and the mRNA is contacted with the probe(s), for example, in an Affymetrix gene chip array. A skilled

- 92 -

artisan can readily adapt known mRNA detection methods for use in detecting the level of mRNA encoded by the markers of the present invention.

An alternative method for determining the level of mRNA marker in a sample involves the process of nucleic acid amplification, *e.g.*, by rtPCR (the experimental embodiment set forth in Mullis, 1987, U.S. Patent No. 4,683,202), ligase chain reaction (Barany, 1991, *Proc. Natl. Acad. Sci. USA*, 88:189-193), self sustained sequence replication (Guatelli *et al.*, 1990, *Proc. Natl. Acad. Sci. USA* 87:1874-1878), transcriptional amplification system (Kwoh *et al.*, 1989, *Proc. Natl. Acad. Sci. USA* 86:1173-1177), Q-Beta Replicase (Lizardi *et al.*, 1988, *Bio/Technology* 6:1197), rolling circle replication (Lizardi *et al.*, U.S. Patent No. 5,854,033) or any other nucleic acid amplification method, followed by the detection of the amplified molecules using techniques well known to those of skill in the art. These detection schemes are especially useful for the detection of nucleic acid molecules if such molecules are present in very low numbers. As used herein, amplification primers are defined as being a pair of nucleic acid molecules that can anneal to 5' or 3' regions of a gene (plus and minus strands, respectively, or vice-versa) and contain a short region in between. In general, amplification primers are from about 10 to 30 nucleotides in length and flank a region from about 50 to 200 nucleotides in length. Under appropriate conditions and with appropriate reagents, such primers permit the amplification of a nucleic acid molecule comprising the nucleotide sequence flanked by the primers.

For *in situ* methods, mRNA does not need to be isolated from the ovarian cells prior to detection. In such methods, a cell or tissue sample is prepared/processed using known histological methods. The sample is then immobilized on a support, typically a glass slide, and then contacted with a probe that can hybridize to mRNA that encodes the marker.

As an alternative to making determinations based on the absolute expression level of the marker, determinations may be based on the normalized expression level of the marker. Expression levels are normalized by correcting the absolute expression level of a marker by comparing its expression to the expression of a gene that is not a marker, *e.g.*, a housekeeping gene that is constitutively expressed. Suitable genes for normalization include housekeeping genes such as the actin gene, or epithelial cell-specific genes. This normalization allows the comparison of the

expression level in one sample, *e.g.*, a patient sample, to another sample, *e.g.*, a non-ovarian cancer sample, or between samples from different sources.

Alternatively, the expression level can be provided as a relative expression level. To determine a relative expression level of a marker, the level of
5 expression of the marker is determined for 10 or more samples of normal versus cancer cell isolates, preferably 50 or more samples, prior to the determination of the expression level for the sample in question. The mean expression level of each of the genes assayed in the larger number of samples is determined and this is used as a baseline expression level for the marker. The expression level of the marker determined for the
10 test sample (absolute level of expression) is then divided by the mean expression value obtained for that marker. This provides a relative expression level.

Preferably, the samples used in the baseline determination will be from ovarian cancer or from non-ovarian cancer cells of ovarian tissue. The choice of the cell source is dependent on the use of the relative expression level. Using expression found
15 in normal tissues as a mean expression score aids in validating whether the marker assayed is ovarian specific (versus normal cells). In addition, as more data is accumulated, the mean expression value can be revised, providing improved relative expression values based on accumulated data. Expression data from ovarian cells provides a means for grading the severity of the ovarian cancer state.

20 In another embodiment of the present invention, a marker protein is detected. A preferred agent for detecting marker protein of the invention is an antibody capable of binding to such a protein or a fragment thereof, preferably an antibody with a detectable label. Antibodies can be polyclonal, or more preferably, monoclonal. An intact antibody, or a fragment or derivatives thereof (*e.g.*, Fab or F(ab')₂) can be used.
25 The term "labeled", with regard to the probe or antibody, is intended to encompass direct labeling of the probe or antibody by coupling (*i.e.*, physically linking) a detectable substance to the probe or antibody, as well as indirect labeling of the probe or antibody by reactivity with another reagent that is directly labeled. Examples of indirect labeling include detection of a primary antibody using a fluorescently labeled secondary antibody
30 and end-labeling of a DNA probe with biotin such that it can be detected with fluorescently labeled streptavidin.

Proteins from ovarian cells can be isolated using techniques that are well known to those of skill in the art. The protein isolation methods employed can, for example, be such as those described in Harlow and Lane (Harlow and Lane, 1988, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York).

A variety of formats can be employed to determine whether a sample contains a protein that binds to a given antibody. Examples of such formats include, but are not limited to, enzyme immunoassay (EIA), radioimmunoassay (RIA), Western blot analysis and enzyme linked immunoabsorbant assay (ELISA). A skilled artisan can readily adapt known protein/antibody detection methods for use in determining whether ovarian cells express a marker of the present invention.

In one format, antibodies, or antibody fragments or derivatives, can be used in methods such as Western blots or immunofluorescence techniques to detect the expressed proteins. In such uses, it is generally preferable to immobilize either the antibody or proteins on a solid support. Suitable solid phase supports or carriers include any support capable of binding an antigen or an antibody. Well-known supports or carriers include glass, polystyrene, polypropylene, polyethylene, dextran, nylon, amylases, natural and modified celluloses, polyacrylamides, gabbros, and magnetite.

One skilled in the art will know many other suitable carriers for binding antibody or antigen, and will be able to adapt such support for use with the present invention. For example, protein isolated from ovarian cells can be run on a polyacrylamide gel electrophoresis and immobilized onto a solid phase support such as nitrocellulose. The support can then be washed with suitable buffers followed by treatment with the detectably labeled antibody. The solid phase support can then be washed with the buffer a second time to remove unbound antibody. The amount of bound label on the solid support can then be detected by conventional means.

The invention also encompasses kits for detecting the presence of a marker protein or nucleic acid in a biological sample (e.g. an ovary-associated body fluid such as a urine sample). Such kits can be used to determine if a subject is suffering from or is at increased risk of developing ovarian cancer. For example, the kit can comprise a labeled compound or agent capable of detecting a marker protein or nucleic acid in a biological sample and means for determining the amount of the protein or

- 95 -

mRNA in the sample (*e.g.*, an antibody which binds the protein or a fragment thereof, or an oligonucleotide probe which binds to DNA or mRNA encoding the protein). Kits can also include instructions for interpreting the results obtained using the kit.

For antibody-based kits, the kit can comprise, for example: (1) a first
5 antibody (*e.g.*, attached to a solid support) which binds to a marker protein; and, optionally, (2) a second, different antibody which binds to either the protein or the first antibody and is conjugated to a detectable label.

For oligonucleotide-based kits, the kit can comprise, for example: (1) an
oligonucleotide, *e.g.*, a detectably labeled oligonucleotide, which hybridizes to a nucleic
10 acid sequence encoding a marker protein or (2) a pair of primers useful for amplifying a marker nucleic acid molecule. The kit can also comprise, *e.g.*, a buffering agent, a preservative, or a protein stabilizing agent. The kit can further comprise components necessary for detecting the detectable label (*e.g.*, an enzyme or a substrate). The kit can also contain a control sample or a series of control samples which can be assayed and
15 compared to the test sample. Each component of the kit can be enclosed within an individual container and all of the various containers can be within a single package, along with instructions for interpreting the results of the assays performed using the kit.

B. Pharmacogenomics

20 Agents or modulators which have a stimulatory or inhibitory effect on expression of a marker of the invention can be administered to individuals to treat (prophylactically or therapeutically) ovarian cancer in the patient. In conjunction with such treatment, the pharmacogenomics (*i.e.*, the study of the relationship between an individual's genotype and that individual's response to a foreign compound or drug) of
25 the individual may be considered. Differences in metabolism of therapeutics can lead to severe toxicity or therapeutic failure by altering the relation between dose and blood concentration of the pharmacologically active drug. Thus, the pharmacogenomics of the individual permits the selection of effective agents (*e.g.*, drugs) for prophylactic or therapeutic treatments based on a consideration of the individual's genotype. Such
30 pharmacogenomics can further be used to determine appropriate dosages and therapeutic regimens. Accordingly, the level of expression of a marker of the invention in an

- 96 -

individual can be determined to thereby select appropriate agent(s) for therapeutic or prophylactic treatment of the individual.

Pharmacogenomics deals with clinically significant variations in the response to drugs due to altered drug disposition and abnormal action in affected persons. See, e.g., Linder (1997) *Clin. Chem.* 43(2):254-266. In general, two types of pharmacogenetic conditions can be differentiated. Genetic conditions transmitted as a single factor altering the way drugs act on the body are referred to as "altered drug action." Genetic conditions transmitted as single factors altering the way the body acts on drugs are referred to as "altered drug metabolism". These pharmacogenetic conditions can occur either as rare defects or as polymorphisms. For example, glucose-6-phosphate dehydrogenase (G6PD) deficiency is a common inherited enzymopathy in which the main clinical complication is hemolysis after ingestion of oxidant drugs (antimalarials, sulfonamides, analgesics, nitrofurans) and consumption of fava beans.

As an illustrative embodiment, the activity of drug metabolizing enzymes is a major determinant of both the intensity and duration of drug action. The discovery of genetic polymorphisms of drug metabolizing enzymes (e.g., N-acetyltransferase 2 (NAT 2) and cytochrome P450 enzymes CYP2D6 and CYP2C19) has provided an explanation as to why some patients do not obtain the expected drug effects or show exaggerated drug response and serious toxicity after taking the standard and safe dose of a drug. These polymorphisms are expressed in two phenotypes in the population, the extensive metabolizer (EM) and poor metabolizer (PM). The prevalence of PM is different among different populations. For example, the gene coding for CYP2D6 is highly polymorphic and several mutations have been identified in PM, which all lead to the absence of functional CYP2D6. Poor metabolizers of CYP2D6 and CYP2C19 quite frequently experience exaggerated drug response and side effects when they receive standard doses. If a metabolite is the active therapeutic moiety, a PM will show no therapeutic response, as demonstrated for the analgesic effect of codeine mediated by its CYP2D6-formed metabolite morphine. The other extreme are the so called ultra-rapid metabolizers who do not respond to standard doses. Recently, the molecular basis of ultra-rapid metabolism has been identified to be due to CYP2D6 gene amplification.

Thus, the level of expression of a marker of the invention in an individual can be determined to thereby select appropriate agent(s) for therapeutic or prophylactic treatment of the individual. In addition, pharmacogenetic studies can be used to apply genotyping of polymorphic alleles encoding drug-metabolizing enzymes to the
5 identification of an individual's drug responsiveness phenotype. This knowledge, when applied to dosing or drug selection, can avoid adverse reactions or therapeutic failure and thus enhance therapeutic or prophylactic efficiency when treating a subject with a modulator of expression of a marker of the invention.

10 C. Monitoring Clinical Trials

Monitoring the influence of agents (*e.g.*, drug compounds) on the level of expression of a marker of the invention can be applied not only in basic drug screening, but also in clinical trials. For example, the effectiveness of an agent to affect marker expression can be monitored in clinical trials of subjects receiving treatment for ovarian
15 cancer. In a preferred embodiment, the present invention provides a method for monitoring the effectiveness of treatment of a subject with an agent (*e.g.*, an agonist, antagonist, peptidomimetic, protein, peptide, nucleic acid, small molecule, or other drug candidate) comprising the steps of (i) obtaining a pre-administration sample from a subject prior to administration of the agent; (ii) detecting the level of expression of one
20 or more selected markers of the invention in the pre-administration sample; (iii) obtaining one or more post-administration samples from the subject; (iv) detecting the level of expression of the marker(s) in the post-administration samples; (v) comparing the level of expression of the marker(s) in the pre-administration sample with the level of expression of the marker(s) in the post-administration sample or samples; and (vi)
25 altering the administration of the agent to the subject accordingly. For example, increased administration of the agent can be desirable to increase expression of the marker(s) to higher levels than detected, *i.e.*, to increase the effectiveness of the agent. Alternatively, decreased administration of the agent can be desirable to decrease expression of the marker(s) to lower levels than detected, *i.e.*, to decrease the
30 effectiveness of the agent.

D. Electronic Apparatus Readable Media and Arrays

Electronic apparatus readable media comprising a marker of the present invention is also provided. As used herein, "electronic apparatus readable media" refers to any suitable medium for storing, holding or containing data or information that can be
5 read and accessed directly by an electronic apparatus. Such media can include, but are not limited to: magnetic storage media, such as floppy discs, hard disc storage medium, and magnetic tape; optical storage media such as compact disc; electronic storage media such as RAM, ROM, EPROM, EEPROM and the like; general hard disks and hybrids of these categories such as magnetic/optical storage media. The medium is adapted or
10 configured for having recorded thereon a marker of the present invention.

As used herein, the term "electronic apparatus" is intended to include any suitable computing or processing apparatus or other device configured or adapted for storing data or information. Examples of electronic apparatus suitable for use with the present invention include stand-alone computing apparatus; networks, including a local
15 area network (LAN), a wide area network (WAN) Internet, Intranet, and Extranet; electronic appliances such as a personal digital assistants (PDAs), cellular phone, pager and the like; and local and distributed processing systems.

As used herein, "recorded" refers to a process for storing or encoding information on the electronic apparatus readable medium. Those skilled in the art can
20 readily adopt any of the presently known methods for recording information on known media to generate manufactures comprising the markers of the present invention.

A variety of software programs and formats can be used to store the marker information of the present invention on the electronic apparatus readable medium. For example, the marker nucleic acid sequence can be represented in a word
25 processing text file, formatted in commercially-available software such as WordPerfect and MicroSoft Word, or represented in the form of an ASCII file, stored in a database application, such as DB2, Sybase, Oracle, or the like, as well as in other forms. Any number of data processor structuring formats (*e.g.*, text file or database) may be employed in order to obtain or create a medium having recorded thereon the the markers
30 of the present invention.

- 99 -

By providing the markers of the invention in readable form, one can routinely access the marker sequence information for a variety of purposes. For example, one skilled in the art can use the nucleotide or amino acid sequences of the present invention in readable form to compare a target sequence or target structural
5 motif with the sequence information stored within the data storage means. Search means are used to identify fragments or regions of the sequences of the invention which match a particular target sequence or target motif.

The present invention therefore provides a medium for holding instructions for performing a method for determining whether a subject has ovarian
10 cancer or a pre-disposition to ovarian cancer, wherein the method comprises the steps of determining the presence or absence of a marker and based on the presence or absence of the marker, determining whether the subject has ovarian cancer or a pre-disposition to ovarian cancer and/or recommending a particular treatment for ovarian cancer or pre-ovarian cancer condition.

The present invention further provides in an electronic system and/or in a network, a method for determining whether a subject has ovarian cancer or a pre-disposition to ovarian cancer associated with a marker wherein the method comprises the steps of determining the presence or absence of the marker, and based on the presence or absence of the marker, determining whether the subject has ovarian cancer
20 or a pre-disposition to ovarian cancer, and/or recommending a particular treatment for the ovarian cancer or pre-ovarian cancer condition. The method may further comprise the step of receiving phenotypic information associated with the subject and/or acquiring from a network phenotypic information associated with the subject.

The present invention also provides in a network, a method for
25 determining whether a subject has ovarian cancer or a pre-disposition to ovarian cancer associated with a marker, said method comprising the steps of receiving information associated with the marker receiving phenotypic information associated with the subject, acquiring information from the network corresponding to the marker and/or ovarian cancer, and based on one or more of the phenotypic information, the marker, and the
30 acquired information, determining whether the subject has a ovarian cancer or a pre-disposition to ovarian cancer. The method may further comprise the step of

- 99 -

By providing the markers of the invention in readable form, one can routinely access the marker sequence information for a variety of purposes. For example, one skilled in the art can use the nucleotide or amino acid sequences of the present invention in readable form to compare a target sequence or target structural
5 motif with the sequence information stored within the data storage means. Search means are used to identify fragments or regions of the sequences of the invention which match a particular target sequence or target motif.

The present invention therefore provides a medium for holding instructions for performing a method for determining whether a subject has ovarian
10 cancer or a pre-disposition to ovarian cancer, wherein the method comprises the steps of determining the presence or absence of a marker and based on the presence or absence of the marker, determining whether the subject has ovarian cancer or a pre-disposition to ovarian cancer and/or recommending a particular treatment for ovarian cancer or pre-ovarian cancer condition.

The present invention further provides in an electronic system and/or in a network, a method for determining whether a subject has ovarian cancer or a pre-disposition to ovarian cancer associated with a marker wherein the method comprises the steps of determining the presence or absence of the marker, and based on the presence or absence of the marker, determining whether the subject has ovarian cancer
20 or a pre-disposition to ovarian cancer, and/or recommending a particular treatment for the ovarian cancer or pre-ovarian cancer condition. The method may further comprise the step of receiving phenotypic information associated with the subject and/or acquiring from a network phenotypic information associated with the subject.

The present invention also provides in a network, a method for
25 determining whether a subject has ovarian cancer or a pre-disposition to ovarian cancer associated with a marker, said method comprising the steps of receiving information associated with the marker receiving phenotypic information associated with the subject, acquiring information from the network corresponding to the marker and/or ovarian cancer, and based on one or more of the phenotypic information, the marker, and the
30 acquired information, determining whether the subject has a ovarian cancer or a pre-disposition to ovarian cancer. The method may further comprise the step of

- 100 -

recommending a particular treatment for the ovarian cancer or pre-ovarian cancer condition.

The present invention also provides a business method for determining whether a subject has ovarian cancer or a pre-disposition to ovarian cancer, said method comprising the steps of receiving information associated with the marker, receiving phenotypic information associated with the subject, acquiring information from the network corresponding to the marker and/or ovarian cancer, and based on one or more of the phenotypic information, the marker, and the acquired information, determining whether the subject has ovarian cancer or a pre-disposition to ovarian cancer. The method may further comprise the step of recommending a particular treatment for the ovarian cancer or pre-ovarian cancer condition.

The invention also includes an array comprising a marker of the present invention. The array can be used to assay expression of one or more genes in the array. In one embodiment, the array can be used to assay gene expression in a tissue to ascertain tissue specificity of genes in the array. In this manner, up to about 7600 genes can be simultaneously assayed for expression. This allows a profile to be developed showing a battery of genes specifically expressed in one or more tissues.

In addition to such qualitative determination, the invention allows the quantitation of gene expression. Thus, not only tissue specificity, but also the level of expression of a battery of genes in the tissue is ascertainable. Thus, genes can be grouped on the basis of their tissue expression *per se* and level of expression in that tissue. This is useful, for example, in ascertaining the relationship of gene expression between or among tissues. Thus, one tissue can be perturbed and the effect on gene expression in a second tissue can be determined. In this context, the effect of one cell type on another cell type in response to a biological stimulus can be determined. Such a determination is useful, for example, to know the effect of cell-cell interaction at the level of gene expression. If an agent is administered therapeutically to treat one cell type but has an undesirable effect on another cell type, the invention provides an assay to determine the molecular basis of the undesirable effect and thus provides the opportunity to co-administer a counteracting agent or otherwise treat the undesired effect. Similarly, even within a single cell type, undesirable biological effects can be

determined at the molecular level. Thus, the effects of an agent on expression of other than the target gene can be ascertained and counteracted.

In another embodiment, the array can be used to monitor the time course of expression of one or more genes in the array. This can occur in various biological contexts, as disclosed herein, for example development of ovarian cancer, progression of ovarian cancer, and processes, such a cellular transformation associated with ovarian cancer.

The array is also useful for ascertaining the effect of the expression of a gene on the expression of other genes in the same cell or in different cells. This provides, for example, for a selection of alternate molecular targets for therapeutic intervention if the ultimate or downstream target cannot be regulated.

The array is also useful for ascertaining differential expression patterns of one or more genes in normal and abnormal cells. This provides a battery of genes that could serve as a molecular target for diagnosis or therapeutic intervention.

15

E. Surrogate Markers

The markers of the invention may serve as surrogate markers for one or more disorders or disease states or for conditions leading up to disease states, and in particular, ovarian cancer. As used herein, a "surrogate marker" is an objective biochemical marker which correlates with the absence or presence of a disease or disorder, or with the progression of a disease or disorder (e.g., with the presence or absence of a tumor). The presence or quantity of such markers is independent of the disease. Therefore, these markers may serve to indicate whether a particular course of treatment is effective in lessening a disease state or disorder. Surrogate markers are of particular use when the presence or extent of a disease state or disorder is difficult to assess through standard methodologies (e.g., early stage tumors), or when an assessment of disease progression is desired before a potentially dangerous clinical endpoint is reached (e.g., an assessment of cardiovascular disease may be made using cholesterol levels as a surrogate marker, and an analysis of HIV infection may be made using HIV RNA levels as a surrogate marker, well in advance of the undesirable clinical outcomes of myocardial infarction or fully-developed AIDS). Examples of the use of surrogate

30

markers in the art include: Koomen *et al.* (2000) *J. Mass. Spectrom.* 35: 258-264; and James (1994) *AIDS Treatment News Archive* 209.

The markers of the invention are also useful as pharmacodynamic markers. As used herein, a "pharmacodynamic marker" is an objective biochemical marker which correlates specifically with drug effects. The presence or quantity of a pharmacodynamic marker is not related to the disease state or disorder for which the drug is being administered; therefore, the presence or quantity of the marker is indicative of the presence or activity of the drug in a subject. For example, a pharmacodynamic marker may be indicative of the concentration of the drug in a biological tissue, in that the marker is either expressed or transcribed or not expressed or transcribed in that tissue in relationship to the level of the drug. In this fashion, the distribution or uptake of the drug may be monitored by the pharmacodynamic marker. Similarly, the presence or quantity of the pharmacodynamic marker may be related to the presence or quantity of the metabolic product of a drug, such that the presence or quantity of the marker is indicative of the relative breakdown rate of the drug *in vivo*. Pharmacodynamic markers are of particular use in increasing the sensitivity of detection of drug effects, particularly when the drug is administered in low doses. Since even a small amount of a drug may be sufficient to activate multiple rounds of marker transcription or expression, the amplified marker may be in a quantity which is more readily detectable than the drug itself. Also, the marker may be more easily detected due to the nature of the marker itself; for example, using the methods described herein, antibodies may be employed in an immune-based detection system for a protein marker, or marker-specific radiolabeled probes may be used to detect a mRNA marker. Furthermore, the use of a pharmacodynamic marker may offer mechanism-based prediction of risk due to drug treatment beyond the range of possible direct observations. Examples of the use of pharmacodynamic markers in the art include: Matsuda *et al.* US 6,033,862; Hattis *et al.* (1991) *Env. Health Perspect.* 90: 229-238; Schentag (1999) *Am. J. Health-Syst. Pharm.* 56 Suppl. 3: S21-S24; and Nicolau (1999) *Am. J. Health-Syst. Pharm.* 56 Suppl. 3: S16-S20.

VI. Experimental Protocol for all OV markers and M352 - M360

A. Identification of markers

The markers of the present invention were identified by transcriptional
5 profiling using mRNA from 9 normal ovarian epithelia, 11 stage I/II ovarian cancer
tumors and 25 stage III/IV tumors. Clones having expression at least two-fold higher in
ovarian tumors as compared to their expression in non-ovarian tumor tissues in at least 4
tumor samples were selected to have their protein-encoding transcript sequences
determined.

10

B. Identification of Markers and Assembly of Their Sequences

Clones which displayed an increase in expression in ovarian tumor
samples over the corresponding average expression of non-tumor samples were used for
further study. Briefly, BLAST analysis, against both public and proprietary sequence
15 databases, of EST sequences known to be associated with each clone was performed,
either directly or in the context of automatically, high-stringency assembled contiguous
sequences. An identification of protein sequence corresponding to the clone was
accomplished by obtaining one of the following:

a) a direct match between the protein sequence and at least one EST
20 sequence in one of its 6 possible translations;

b) a direct match between the nucleotide sequence for the mRNA
corresponding to the protein sequence and at least one EST sequence;

c) a match between the protein sequence and a contiguous assembly
(contig) of the EST sequences with other available EST sequences in the databases in
25 one of its 6 possible translations; or

d) a match between the nucleotide sequence for the mRNA
corresponding to the protein sequence and a contiguous assembly of the EST sequences
with other available EST sequences in the databases in one of its 6 possible translations.

C. Identification of Markers Having Newly-Identified Nucleotide and Amino Acid Sequences.

The markers of Table 2 include newly-identified amino acid sequences.

- 5 These sequences were found to be novel based on one of the following criteria:
- a) the protein sequence found within available public databases was incomplete or erroneous, leading to the construction of an additional completed/corrected protein sequence that is not found as such in the public domain;
 - b) based on nucleotide evidence, variants of the protein sequence were
10 additionally constructed that are not found as such in the public domain; or
 - c) the contig for the EST sequences did not match any known protein, so that a novel protein sequence was derived from an open reading frame of the contig.

- 15 VII. Experimental Protocol for M68, M103, M138, M185, M312, M327-M328, M400, M430-M480, M559, M571-M573, M575-M576, M578-M583, M585-594, and M604-M617

A. Identification of Markers and Assembly of Their Sequences

- 20 The markers of the present invention were identified by transcription profiling using mRNA from 67 ovarian tumors of various histotypes and stage and 96 non-ovarian tumor tissues including normal ovarian epithelium, benign conditions, other normal tissues, and other abnormal tissues. Clones having expression at least three-fold higher in at least 10% of ovarian tumors, as compared to their expression in non-ovarian
25 tumor tissue, were designated as ovarian cancer specific markers. These cDNA clones were selected to have their protein-encoding transcript sequences determined. Briefly, BLAST analysis, against both public and proprietary sequence databases, of EST sequences known to be associated with each clone was performed, either directly or in the context of automatically, high-stringency assembled contiguous sequences. An
30 identification of protein sequence corresponding to the clone was accomplished by obtaining one of the following:
- a) a direct match between the protein sequence and at least one EST sequence in one of its 6 possible translations;

- b) a direct match between the nucleotide sequence for the mRNA corresponding to the protein sequence and at least one EST sequence;
- c) a match between the protein sequence and a contiguous assembly (contig) of the EST sequences with other available EST sequences in the databases in one of its 6 possible translations; or
- d) a match between the nucleotide sequence for the mRNA corresponding to the protein sequence and a contiguous assembly of the EST sequences with other available EST sequences in the databases in one of its 6 possible translations.

10 B. Identification of Markers Having Newly-Identified Amino Acid Sequences.

The markers of Table 2 include newly-identified amino acid sequences. These sequences were found to be novel based on one of the following criteria:

- a) the protein sequence found within available public databases was incomplete or erroneous, leading to the construction of an additional completed/corrected protein sequence that is not found as such in the public domain;
- b) based on nucleotide evidence, variants of the protein sequence were additionally constructed that are not found as such in the public domain; or
- c) the contig for the EST sequences did not match any known protein, so that a novel protein sequence was derived from an open reading frame of the contig.

VIII. Gene Expression Analysis

Total RNA from normal human tissue was obtained from commercial sources. The integrity of the RNA was verified by agarose gel electrophoresis and ethidium bromide staining. Cell lines were purchased from ATCC and grown under the conditions recommended by ATCC. Total RNA from a number of various cell lines was prepared using commercial kits (Qiagen). First strand cDNA was prepared using oligo-dT primer and standard conditions. Each RNA preparation was treated with DNase I (Ambion) at 37°C for 1 hour.

30 Novel gene expression was measured by TaqMan[®] quantitative PCR (Perkin Elmer Applied Biosystems) in cDNA prepared from the following normal human tissues: heart, kidney, skeletal muscle, pancreas, skin, dorsal root ganglion,

- 106 -

breast, ovary, prostate, salivary glands, lung, colon, liver and lymph node. Figure 1 graphically represents the results of the TaqMan® expression study. The columns labelled A to V depict the expression level observed for OV88 in the following tissues:

- Column A: Heart, normal tissue
- 5 Column B: Heart, CHF tissue
- Column C: Kidney, normal tissue
- Column D: Skeletal muscle, normal tissue
- Column E: Pancreas, normal tissue
- Column F: Skin, normal tissue
- 10 Column G: Dorsal root, normal tissue
- Column H: Breast, normal tissue
- Column I: Breast, tumor tissue
- Column J: Ovary, normal tissue
- Column K: Ovary, tumor tissue
- 15 Column L: Prostate, normal tissue
- Column M: Prostate, tumor tissue
- Column N: Salivary glands, normal tissue
- Column O: Lung, normal tissue
- Column P: Lung, tumor tissue
- 20 Column Q: Lung, COPD tissue
- Column R: Colon, IBD tissue
- Column S: Liver, normal tissue
- Column T: Liver fibrosis
- Column U: Lymph node, normal tissue
- 25 Column V: Positive control

IX. Summary of the Data Provided in the Tables

- Tables 1-3 list the markers of the present invention. In the Tables the markers are identified with a name ("Marker"), the name the gene is commonly known by, if applicable ("Gene Name"), the Sequence Listing identifier of the cDNA sequence of a nucleotide transcript encoded by or corresponding to the marker ("SEQ ID NO (nts)"), the Sequence Listing identifier of the amino acid sequence of a protein encoded

- 107 -

by the nucleotide transcript ("SEQ ID NO (AAs)"), and the location of the protein coding sequence within the cDNA sequence ("CDS").


Table 1 lists all of the markers of the invention, which are over-expressed in ovarian cancer cells compared to normal (*i.e.*, non-cancerous) ovarian cells and
5 comprises markers listed in Tables 2 and 3. Table 2 lists newly-identified nucleotide and amino acid sequences useful as ovarian cancer markers. Table 3 lists newly-identified nucleotide sequences useful as ovarian cancer markers.

Other Embodiments

10 Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims:

What is claimed:

1. A method of assessing whether a patient is afflicted with ovarian cancer, the method comprising comparing:
 - 5 a) the level of expression of a marker in a patient sample, wherein the marker is selected from Table 1, and
 - b) the normal level of expression of the marker in a control non-ovarian cancer sample,
- 10 wherein a significant increase in the level of expression of the marker in the patient sample and the normal level is an indication that the patient is afflicted with ovarian cancer.

Figure 1

SEQUENCE LISTING

<110> Millennium Pharmaceuticals, Inc. et al.

<120> Nucleic Acid Molecules and Proteins For The Identification,
Assessment, Prevention, and Therapy of Ovarian Cancer

<130> MRI-030FC

<150> 60/276,025

<151> 2001-03-14

<150> 60/325,149

<151> 2001-09-26

<150> 60/276,026

<151> 2001-03-14

<150> 60/324,967

<151> 2001/09/26

<150> 60/311,732

<151> 2001-08-10

<150> 60/325,102

<151> 2001-09-26

<150> 60/323,580

<151> 2001-09-19

<160> 363

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 4643

<212> DNA

<213> Homo sapiens

<400> 1

```

cctactctat tcagatatcc tccagattcc taaagattag agatcatttc tcattctoct 60
aggagtactc acttcaggaa gcaaccagat aaaagagagg tgcaacggaa gccagaacat 120
tctcctggga aattcaacct gtttcgcagt ttctcgagga atcagcattc agtcaatccg 180
ggccggggagc agtcactctgt ggtgaggctg attggtctgg caggaaacagc gccggggcgt 240
gggctgagca cagcgcttcg ctctctttgc cacaggaaagc ctgagctcat togagtagcg 300
gctcttccaa gctcaaaaga gcaagggccg ctgttcgttt cctttaggtc ttccactaa 360
agtgcgagta tcttcttcca agatttcaag tcttgggtgc cgttccaagg agcgcgaggt 420
cgggatggat cttgaagggg accgcaatgg aggagcaaaag aagaagaact tttttaact 480
gaacaataaa agtgaaaaag ataagaagga aaagaaccca actgtcagtg tattttcaat 540
gtttcgctat tcaaattggc ttgacaagtt gtatatggtg gtgggaactt tggctgccat 600
catccatggg gctggacttc ctctcatgat gctggtgttt ggagaaatga cagatatctt 660
tgcaaatgca ggaattttag aagatctgat gtcaaacatc actaatagaa gtgatatcaa 720
tgatacaggg ttcttcatga atctggagga agacatgacc aggtatgcct attattacag 780
tggaattggt gctggggtgc tggttgctgc ttacattcag gtttcatttt ggtgcctggc 840
agctggaaga caaatcacaa aaattagaaa acagtttttt catgctataa tgcacagga 900
gataggctgg ttgatgtgc acgatgttg ggagcttaac acccgactta catgatgtg 960
ctccaagatt aatgaaggaa ttggtgacaa aattggaatg ttctttcagt caatggcaac 1020

```


atttttcaact	gggttttatag	taggattttac	acgtggttgg	aagctaacc	ttgtgatttt	1080
ggccatcagt	ccgtgtcttg	gactgtcagc	tgctgtctgg	gcaaagatac	tatctcoatt	1140
tactgatataa	gaactcttag	cgtatgcata	agctggagca	gtagctgaag	aggtctctggc	1200
agcaattaga	cgtgtgattg	catttggagg	acaaaagaaa	gaacttgaaa	ggtacaaacaa	1260
aaatttagaa	gaagctcaaaa	gaattgggat	aaagaaagct	attacagcca	atattttctat	1320
aggtgtcgt	ttctcgtga	ttctatgcate	ttatgctctg	gccctctggg	atgggacac	1380
cttggctctc	tcagggggaat	attctatltgg	acaagtaotc	actgtatttt	ctgtatttaatt	1440
ttggggctttt	agtgttggac	aggaatctcc	aagcattgaa	gcatttgcac	gtccaaagag	1500
agcagcttat	gaaattttcca	agataattga	taataagcca	agtatgtgca	gctattcgaa	1560
gagtgggcac	naaacagata	atattaaggg	aaatttggaa	ttcagaatgt	ttcaactcag	1620
ttaccatct	cgaaaaagag	ttaagatctt	gaagggctcg	aacctgaagg	tcgacaggtg	1680
gcagacgggt	gccctgggtt	gaacacagtg	ctgtggggaag	agcacaacag	tcacagtgat	1740
gcagaggtct	tatgacccca	cagaggggat	ggctcagtggt	gatggacagg	atattagcat	1800
catataatga	aggtttctac	gggaatacat	tggtgtgtgtg	agtcagggaac	ctgtatttgtt	1860
tgccaccaog	atagctgaaa	acattcgcta	tgccogtgaa	aatgtcacca	tggtatgagat	1920
tgagaaagct	gtcaagggaag	ccaatgccta	tgacclttatc	atgaaactgc	ctcataaatt	1980
tgacacccgt	gtgtggagaga	gagggggccca	gttgagtggg	gggcagaagc	agaggtatgc	2040
catctgcagt	gccctgggtc	gcaaccccca	gatccctcgt	ctggatgagg	ccacgtcagc	2100
cttggacaca	gaaagcgaag	cagtggttca	ggtygctctg	gataaggcca	gaaaggtctg	2160
gaocacatt	gtgatgtctc	atcgtttgtg	tcacagtctg	taagctgagc	tcactgtcgt	2220
ttctgatgat	ggagtcaatt	tgagagaagg	naatcatgat	gaactcatga	aagagaaagg	2280
cttactactc	aaactgttca	caatgcagac	agcaggaat	gaagttgaat	tagaanaatgc	2340
agctgatgaa	tcocaaaagt	aaattgatgc	cttgaaaagt	tcctcaaatg	atccaagatc	2400
cagtcataata	agaaaagatg	caactcgtag	gagtgctcgt	ggatcacacag	ccaagagacag	2460
aaagcttagt	accaaaagag	ctctggatga	aagtataact	ccagtttctc	tttgaggagat	2520
tatgaagcta	aatttaactg	aatggccctt	ttttgttttt	gtgttatttt	gtgccattat	2580
aaatggaggg	ctgocaaacc	catttgcatt	aatattttca	aagattatag	gggttttttc	2640
aagattgat	gatcctgaaa	caaaaagca	gaatagttaac	ttgttttcaac	tattgtttct	2700
agoccttgg	attattttct	ttattacatt	tttctctcag	ggtttcacat	ttggcaaaagc	2760
ttggagagat	ctacacaaag	ggctccgata	catgttttct	cgatccatgc	tcagacagga	2820
ttgtagtgtg	ttttgatgacc	ctaaaaaac	cactggagca	ttgaclacca	tgctgtccaa	2880
tgatgtcgt	caagttaaag	gggtctatag	ttccagcctt	gctgtaat	ccagaaat	2940
agcaaatct	tgacagagaa	taattatata	cttcatctat	ggttggcaac	taacactgtt	3000
actcttagca	attgtaccac	tcattgcaat	agcaggaggt	gttgaaatga	aaatgttgtc	3060
tggaacaaga	ctgaagatga	agaaagaact	agaaggtgct	gggaagatcg	ctactgaagc	3120
aataagaaaac	ttccagaccg	ttgtttcttt	gactcaggag	cagaagtttg	aacatatgta	3180
tgctcagagt	ttgcaggtag	catacagaaa	ctctttgagg	aaagcacaca	ttcttggaat	3240
tacattttcc	ttcacccagg	caatgatgta	tttttctact	gctgtgattg	tcocgttttg	3300
agocctactg	gtggcacata	aaactcatgag	ctttgagtag	gttctctiag	tattttcagc	3360
tggtgtcttt	gggtgccatg	ccgtggggca	agtcaagtca	tttgctcctg	actatgcca	3420
agocaaaaat	tcagcagccc	acatcatcat	gatcattgaa	aaaacccctt	tgattgacag	3480
ctacagcag	gaagggccta	tgccgaacac	attggaagga	aatgtcacat	ttgggtgaag	3540
tgatttcaac	tatcccacc	gaocggacat	ccacgtgctt	cagggaactga	gocctggagg	3600
gaagaaggcc	cagacagctg	ctctgggtgg	cagcagctgt	ttggggaaga	gcacagtggt	3660
ccagctcctg	gagcgggtct	agcacccctt	ggcagggaaa	gtgctgcttg	atggcaagaa	3720
aataaagcga	ctgaattgtc	agtggctccg	agcacacctg	ggcatcgtgt	ccagaggacc	3780
catctctgtt	gaotcagca	ttgtcagaaa	cattgctcat	ggagacaaca	gccgggtggt	3840
gtcacaggaa	gagattgtga	gggcagcaaa	ggagcccaac	atacatgctc	tcacatgagtc	3900
actgctaat	aaatatagca	ctaaagttag	agacaaagga	actcagctct	ctgggtggcca	3960
gaacacacgc	attgtccatg	ctcgtgccct	ttgtagacag	otcatattt	tgcttttggga	4020
tgaagccacg	tcagctcttg	atacagaag	tgaaaaggtt	gtccaagaag	ccctggacaa	4080
agocagagaa	ggcgcacct	gcaattgtgat	tgctccacgc	ctgtccacca	tcacagaatgc	4140
agacttaata	gtgggttttc	agaattggcag	agtcnaagag	catggcagca	atcacagctg	4200
gctggcagac	aaaggcatct	atttttcaat	ggctcagctc	caggctggaa	caaaagccca	4260
gtgaactctg	actgtatgag	atgtttaata	ctttttaata	ttgttttaga	tatgacattt	4320
atccaaagtt	aaaagcaaac	acttacagaa	ttatgaagag	gtactctgtt	aaactttctc	4380
cagtcnaagtt	cagagtcttc	agagacttgc	taattaaagg	aaacagagtga	gagacatcat	4440
caagtggaga	gaaatcatag	tttaaacctgc	attataaatt	ttataacaga	attaaagtag	4500

attttaaaag ataaaaatgtg taatttttgtt tatattttcc catttggact gtaactgact 4560
 gccttgctaa aagattatag aagtagcaaa aagtattgaa atgtttgcat aaagtgtcta 4620
 taataaaact aaactttcat gtg 4643

<210> 2

<211> 1279

<212> PRT

<213> Homo sapiens

<400> 2

Met	Asp	Leu	Glu	Gly	Asp	Arg	Asn	Gly	Gly	Ala	Lys	Lys	Lys	Asn	Phe
1				5					10					15	
Phe	Lys	Leu	Asn	Asn	Lys	Ser	Glu	Lys	Asp	Lys	Lys	Glu	Lys	Lys	Pro
			20					25					30		
Thr	Val	Ser	Val	Phe	Ser	Met	Phe	Arg	Tyr	Ser	Asn	Trp		Leu	Asp
			35					40				45			
Leu	Tyr	Met	Val	Val	Gly	Thr	Leu	Ala	Ala	Ile	Ile	His	Gly	Ala	Gly
			50					55				60			
Leu	Pro	Leu	Met	Met	Leu	Val	Phe	Gly	Glu	Met	Thr	Asp	Ile	Phe	Ala
			65					70				75			80
Asn	Ala	Gly	Asn	Leu	Glu	Asp	Leu	Met	Ser	Asn	Ile	Thr	Asn	Arg	Ser
			85					90					95		
Asp	Ile	Asn	Asp	Thr	Gly	Phe	Phe	Met	Asn	Leu	Glu	Glu	Asp	Met	Thr
			100					105					110		
Arg	Tyr	Ala	Tyr	Tyr	Tyr	Ser	Gly	Ile	Gly	Ala	Gly	Val	Leu	Val	Ala
			115					120				125			
Ala	Tyr	Ile	Gln	Val	Ser	Phe	Trp	Cys	Leu	Ala	Ala	Gly	Arg	Gln	Ile
			130					135				140			
His	Lys	Ile	Arg	Lys	Gln	Phe	Phe	His	Ala	Ile	Met	Arg	Gln	Glu	Ile
			145					150			155			160	
Gly	Trp	Phe	Asp	Val	His	Asp	Val	Gly	Glu	Leu	Asn	Thr	Arg	Leu	Thr
			165					170					175		
Asp	Asp	Val	Ser	Lys	Ile	Asn	Glu	Gly	Ile	Gly	Asp	Lys	Ile	Gly	Met
			180					185					190		
Phe	Phe	Gln	Ser	Met	Ala	Thr	Phe	Phe	Thr	Gly	Phe	Ile	Val	Gly	Phe
			195					200				205			
Thr	Arg	Gly	Trp	Lys	Leu	Thr	Leu	Val	Ile	Leu	Ala	Ile	Ser	Pro	Val
			210					215				220			
Leu	Gly	Leu	Ser	Ala	Ala	Val	Trp	Ala	Lys	Ile	Leu	Ser	Ser	Phe	Thr
			225					230				235			240
Asp	Lys	Glu	Leu	Leu	Ala	Tyr	Ala	Lys	Ala	Gly	Ala	Val	Ala	Glu	Glu
			245					250					255		
Val	Leu	Ala	Ala	Ile	Arg	Thr	Val	Ile	Ala	Phe	Gly	Gly	Gln	Lys	Lys
			260					265					270		
Glu	Leu	Glu	Arg	Tyr	Asn	Lys	Asn	Leu	Glu	Glu	Ala	Lys	Arg	Ile	Gly
			275					280				285			
Ile	Lys	Lys	Ala	Ile	Thr	Ala	Asn	Ile	Ser	Ile	Gly	Ala	Ala	Phe	Leu
			290					295				300			
Leu	Ile	Tyr	Ala	Ser	Tyr	Ala	Leu	Ala	Phe	Trp	Tyr	Gly	Thr	Thr	Leu
			305					310				315			320
Val	Leu	Ser	Gly	Glu	Tyr	Ser	Ile	Gly	Gln	Val	Leu	Thr	Val	Phe	Ser
			325					330					335		
Val	Leu	Ile	Gly	Ala	Phe	Ser	Val	Gly	Gln	Ala	Ser	Pro	Ser	Ile	Glu
			340					345				350			
Ala	Phe	Ala	Asn	Ala	Arg	Gly	Ala	Ala	Tyr	Glu	Ile	Phe	Lys	Ile	Ile
			355					360				365			
Asp	Asn	Lys	Pro	Ser	Ile	Asp	Ser	Tyr	Ser	Lys	Ser	Gly	His	Lys	Pro
			370					375				380			

Asp Asn Ile Lys Gly Asn Leu Glu Phe Arg Asn Val His Phe Ser Tyr
 385 390 395 400
 Pro Ser Arg Lys Glu Val Lys Ile Leu Lys Gly Leu Asn Leu Lys Val
 405 410 415
 Gln Ser Gly Gln Thr Val Ala Leu Val Gly Asn Ser Gly Cys Gly Lys
 420 425 430
 Ser Thr Thr Val Gln Leu Met Gln Arg Leu Tyr Asp Pro Thr Glu Gly
 435 440 445
 Met Val Ser Val Asp Gly Gln Asp Ile Arg Thr Ile Asn Val Arg Phe
 450 455 460
 Leu Arg Glu Ile Ile Gly Val Val Ser Gln Glu Pro Val Leu Phe Ala
 465 470 475 480
 Thr Thr Ile Ala Glu Asn Ile Arg Tyr Gly Arg Glu Asn Val Thr Met
 485 490 495
 Asp Glu Ile Glu Lys Ala Val Lys Glu Ala Asn Ala Tyr Asp Phe Ile
 500 505 510
 Met Lys Leu Pro His Lys Phe Asp Thr Leu Val Gly Glu Arg Gly Ala
 515 520 525
 Gln Leu Ser Gly Gly Gln Lys Gln Arg Ile Ala Ile Ala Arg Ala Leu
 530 535 540
 Val Arg Asn Pro Lys Ile Leu Leu Leu Asp Glu Ala Thr Ser Ala Leu
 545 550 555 560
 Asp Thr Glu Ser Glu Ala Val Val Gln Val Ala Leu Asp Lys Ala Arg
 565 570 575
 Lys Gly Arg Thr Thr Ile Val Ile Ala His Arg Leu Ser Thr Val Arg
 580 585 590
 Asn Ala Asp Val Ile Ala Gly Phe Asp Asp Gly Val Ile Val Glu Lys
 595 600 605
 Gly Asn His Asp Glu Leu Met Lys Glu Lys Gly Ile Tyr Phe Lys Leu
 610 615 620
 Val Thr Met Gln Thr Ala Gly Asn Glu Val Glu Leu Glu Asn Ala Ala
 625 630 635 640
 Asp Glu Ser Lys Ser Glu Ile Asp Ala Leu Glu Met Ser Ser Asn Asp
 645 650 655
 Ser Arg Ser Ser Leu Ile Arg Lys Arg Ser Thr Arg Arg Ser Val Arg
 660 665 670
 Gly Ser Gln Ala Gln Asp Arg Lys Leu Ser Thr Lys Glu Ala Leu Asp
 675 680 685
 Glu Ser Ile Pro Pro Val Ser Phe Trp Arg Ile Met Lys Leu Asn Leu
 690 695 700
 Thr Glu Trp Pro Tyr Phe Val Val Gly Val Phe Cys Ala Ile Ile Asn
 705 710 715 720
 Gly Gly Leu Gln Pro Ala Phe Ala Ile Ile Phe Ser Lys Ile Ile Gly
 725 730 735
 Val Phe Thr Arg Ile Asp Asp Pro Glu Thr Lys Arg Gln Asn Ser Asn
 740 745 750
 Leu Phe Ser Leu Leu Phe Leu Ala Leu Gly Ile Ile Ser Phe Ile Thr
 755 760 765
 Phe Phe Leu Gln Gly Phe Thr Phe Gly Lys Ala Gly Glu Ile Leu Thr
 770 775 780
 Lys Arg Leu Arg Tyr Met Val Phe Arg Ser Met Leu Arg Gln Asp Val
 785 790 795 800
 Ser Trp Phe Asp Asp Pro Lys Asn Thr Thr Gly Ala Leu Thr Thr Arg
 805 810 815
 Leu Ala Asn Asp Ala Ala Gln Val Lys Gly Ala Ile Gly Ser Arg Leu
 820 825 830
 Ala Val Ile Thr Gln Asn Ile Ala Asn Leu Gly Thr Gly Ile Ile Ile
 835 840 845

Ser Phe Ile Tyr Gly Trp Gln Leu Thr Leu Leu Leu Ala Ile Val
 850 855 860
 Pro Ile Ile Ala Ile Ala Gly Val Val Glu Met Lys Met Leu Ser Gly
 865 870 875 880
 Gln Ala Leu Lys Asp Lys Lys Glu Leu Glu Gly Ala Gly Lys Ile Ala
 885 890 895
 Thr Glu Ala Ile Glu Asn Phe Arg Thr Val Val Ser Leu Thr Gln Glu
 900 905 910
 Gln Lys Phe Glu His Met Tyr Ala Gln Ser Leu Gln Val Pro Tyr Arg
 915 920 925
 Asn Ser Leu Arg Lys Ala His Ile Phe Gly Ile Thr Phe Ser Phe Thr
 930 935 940
 Gln Ala Met Met Tyr Phe Ser Tyr Ala Gly Cys Phe Arg Phe Gly Ala
 945 950 955 960
 Tyr Leu Val Ala His Lys Leu Met Ser Phe Glu Asp Val Leu Leu Val
 965 970 975
 Phe Ser Ala Val Val Phe Gly Ala Met Ala Val Gly Gln Val Ser Ser
 980 985 990
 Phe Ala Pro Asp Tyr Ala Lys Ala Lys Ile Ser Ala Ala His Ile Ile
 995 1000 1005
 Met Ile Ile Glu Lys Thr Pro Leu Ile Asp Ser Tyr Ser Thr Glu Gly
 1010 1015 1020
 Leu Met Pro Asn Thr Leu Glu Gly Asn Val Thr Phe Gly Glu Val Val
 1025 1030 1035 1040
 Phe Asn Tyr Pro Thr Arg Pro Asp Ile Pro Val Leu Gln Gly Leu Ser
 1045 1050 1055
 Leu Glu Val Lys Lys Gly Gln Thr Leu Ala Leu Val Gly Ser Ser Gly
 1060 1065 1070
 Cys Gly Lys Ser Thr Val Val Gln Leu Leu Glu Arg Phe Tyr Asp Pro
 1075 1080 1085
 Leu Ala Gly Lys Val Leu Leu Asp Gly Lys Glu Ile Lys Arg Leu Asn
 1090 1095 1100
 Val Gln Trp Leu Arg Ala His Leu Gly Ile Val Ser Gln Glu Pro Ile
 1105 1110 1115 1120
 Leu Phe Asp Cys Ser Ile Ala Glu Asn Ile Ala Tyr Gly Asp Asn Ser
 1125 1130 1135
 Arg Val Val Ser Gln Glu Glu Ile Val Arg Ala Ala Lys Glu Ala Asn
 1140 1145 1150
 Ile His Ala Phe Ile Glu Ser Leu Pro Asn Lys Tyr Ser Thr Lys Val
 1155 1160 1165
 Gly Asp Lys Gly Thr Gln Leu Ser Gly Gly Gln Lys Gln Arg Ile Ala
 1170 1175 1180
 Ile Ala Arg Ala Leu Val Arg Gln Pro His Ile Leu Leu Leu Asp Glu
 1185 1190 1195 1200
 Ala Thr Ser Ala Leu Asp Thr Glu Ser Glu Lys Val Val Gln Glu Ala
 1205 1210 1215
 Leu Asp Lys Ala Arg Glu Gly Arg Thr Cys Ile Val Ile Ala His Arg
 1220 1225 1230
 Leu Ser Thr Ile Gln Asn Ala Asp Leu Ile Val Val Phe Gln Asn Gly
 1235 1240 1245
 Arg Val Lys Glu His Gly Thr His Gln Gln Leu Leu Ala Gln Lys Gly
 1250 1255 1260
 Ile Tyr Phe Ser Met Val Ser Val Gln Ala Gly Thr Lys Arg Gln
 1265 1270 1275

<210> 3

<211> 3859

<212> DNA

<213> Homo sapiens

<400> 3

```

aatctatcag ggaacggcgg tggccgggtc ggcggtgttc gtgcgctctg gccgctcagg 60
ccgtgcggct ggggtgagcg acgcgaggcg gcgagcgggc aagcgtgttt cttagctcgtg 120
cgctcgggct tccggagcct tggcggcagc taggggagga tggcggagtc ttccggataag 180
ctctatcagag tcgagatcagc caaagagcgg cgcgctctct gcaagaaaag tcgagcagagc 240
atccccagag actcgcgcog gatggccatc atgtgtgcatt gccccatggt tcatgagaaa 300
gtccccacct ggtaccacct ctctcgcttc tggaaagttg gccactccat ccggcacccct 360
gacgttgagg tggatgggtt ctctgagctt cggtgggatg accagcagaa agtcaagaag 420
acagcggaag ctggaggagt gacaggcaaa ggccaagatg gaatttgtag caaggcagag 480
aagactctgg gtgactttgc agcagagtat gccaaagtcca acagaagtac gtgccaaggg 540
tgtatggaga agatagaaaa gggccagggt cgctgtcca agaagatgtg ggaccggcag 600
aagccacagc taggcatgat tgaccgcgtg taccatccag gctgctttgt caagaacagg 660
gaggagctgg gtttcggccc cgagtacagt gcgagtcagc tcaagggttt cagcctcctt 720
gctcacagag ataaagaagc cctgaagaag cagctcccag gagtcaagag tgaaggaaa 780
agaaaaaggc atgagtgaga tggagtggat gaagtggcga agaagaaatc taaaaagaa 840
aaagacaagg atagtaagct tgaaaaagcc ctaaaaggctc agaacgacct gatctggaac 900
atcaagagcg agctaaagaa agtgtgttca actaalgacc tgaaggagct actcatcttc 960
aacaagagct aagtcgcttc tggggagtgc gcgatcttgg accgagttagc tgatggcatg 1020
gtgttcgggt cctctcttcc ctgcgaggaa tgctcglttc caagagcgat 1080
gcctattact gcactgggga cgtcaactgc tggaccaagt gtatggtcac gacacagaca 1140
cccaacggga aggagtgggt aacccccaa gaaattccgag aaatctctta cctcaagaaa 1200
tgaagatta aaaaagcaga ccgtatatto cccccagaaa ccagcgcttc cgtggcgccc 1260
acgcctccgc cctccacagc ctcgctcctc gctgctgtga ctctctctgt ttcaagcagt 1320
aagccttatt ccaacatgaa gatcctgact ctcggaagc tgtccggaaa caaggatgaa 1380
gtgaaggcca tgaatgagaa actcgggggg aagttgacgg ggacggccaa caagccttc 1440
ctgtgcatac gccccaaaaa ggaggtggaa aagatgaata agaagtga ggaagttaa 1500
gaagcaaaa tccgagttgt gtctgaggac ttctcccgag acgtctccgc cctccacca 1560
agccttcagg agttgttctt agcgcacatc ttgtcccctt ggggggcaga ggtgaaggag 1620
gagcctgttg aagttgtggc cccaagaggg aagtcagggg ctgcgcttcc caaaaaaagc 1680
aagggccagg tcaaggagga aggtatcaac aaatctgaaa agagaatgaa attraactctt 1740
aaaggaggag cagctgtgga tctgtattct ggactggaa actctgcgca tgtcctggag 1800
aaagttggga aggtcttcag tgccaccctt ggctgtgtg acatcgttaa aggaaccaac 1860
tctactaca agctgcagct tctgagggac gacaaggaaa acaggtattg gatattcagg 1920
tctggggccc gtgtgggtac ggtgatcggt tggaaacaa acgaacaaac tggaaacagt ccgctccaa 1980
gaggatgcca ttgagcagtt catgaaatta tatgaagaaa aaaccgggaa cgttggcac 2040
tccaaaaatt tcaacgaagta tcccaaaaag tttaaccccc tggagattga ctatggcag 2100
gatgaagagg cagtgaaгаа gctcaacgta aatcctggca ccaagtccaa gctccccaa 2160
ccagttcagg acctcatcaa gatgatctt gatgtgaaa gtatgaagaa agocattggt 2220
gagtatgaga tcgacctca gaagatgcc ttggggagaa tgagcaaaag cgagatccag 2280
gcgcataact ccatctcagc tgaggtccag caggcggtgt ctccaggcag cagcgactct 2340
cagatctcgg atctctcaaa tgcgttttac acctgatcc gtccagactt tgggatgaag 2400
aagcctccgc tctgaacaa tgcaagcagt gtgcaggcca aggtggaatt gcttgaaca 2460
ctgtgggaca tcgaggtggc ctacagttcg ctacagggga ggtctgatg tagcagaag 2520
gcgcctcatg atgtcaacta tgaaagctc aaaaactgaca ttaagttgtg tgacagagat 2580
ctgaagaag ccgagatcat caggaaatga gtaagaaca ctcatgcaac cacacaagc 2640
cggtatgact tgaagtcac cgtatcttt aagatagagc gtgaaggcga atgcacaggt 2700
tacaagccct taaagcagct tcaaaccca agattgctg ggcacgggtc caggaccacc 2760
aactttgctg ggtactgtc ccagggtctt cggatagccc cgcctgaagc gccctgaca 2820
ggctacatgt ttggtaaagg gatctatttc gctgacatgg totccaagag tgccaactac 2880
taccatagct ctcaaggaga cccaataggc ttatcctgt ttggagaagt tgccttga 2940
aacatgtatg aactgaagca gccttcacat atcagcaggt taaccaagct caagcacagt 3000
ctcaagggt ttggcaaaac taccctgat ccttcagctc acattagctc ggaattgta 3060
gacgttctcc ttgggacggg gatttcatct acacotctct actataaac 3120
gagtaacatt tctatgat tgcacagga atctgtgaa actctgtaa actgaaattc 3180
aattttaaga cctcctgtg gtaattggga gaggtagcc agtcaacccc ggtgctgtg 3240

```

```

gtatgaattc acccggaagcg cttctgcacc aactcacctg gccgctaagt tgctgatggg 3300
tagtacctgt actaaaccac ctcagaaagg attttacaga aacgtgttaa aggtttttctc 3360
taactttctca agtcocctgt tttgtgtgtg gtctgtgggg aggggtgtgt ttgggggtgtg 3420
ttttgttttt tcttgccagg tagataaaac tgacatagag aaaaggctgg agagagattc 3480
tgttgcatag actagtcccta tggaaaaaac caaagcttgc ttagaalgtc tgccttactg 3540
gtttccccag ggaaggaaaa atacacttcc accctttttt ctaagtgttc gtctttagtt 3600
ttgatttttg aaagatgtta agcattttatt tttagttaaa ataaaaacta atttcatact 3660
atttagattt tcttttttat ctgtcaccta ttgtcccttt tttagttttt ttgttttgcc 3720
tctgtgtgtg aggggtgtgg gaagaccaa ggaaggaaac ctaacaattt ctcatactta 3780
gaacaaaaaa gagcttttct tctccaggaa tactgaacat gggagctctt gaaatatgta 3840
gtattaaaag ttgcattttg                                     3859

```

<210> 4

<211> 1014

<212> PRT

<213> Homo sapiens

<400> 4

```

Met Ala Glu Ser Ser Asp Lys Leu Tyr Arg Val Glu Tyr Ala Lys Ser
1 5 10 15
Gly Arg Ala Ser Cys Lys Lys Cys Ser Glu Ser Ile Pro Lys Asp Ser
20 25 30
Leu Arg Met Ala Ile Met Val Gln Ser Pro Met Phe Asp Gly Lys Val
35 40 45
Pro His Trp Tyr His Phe Ser Cys Phe Trp Lys Val Gly His Ser Ile
50 55 60
Arg His Pro Asp Val Glu Val Asp Gly Phe Ser Glu Leu Arg Trp Asp
65 70 75 80
Asp Gln Gln Lys Val Lys Lys Thr Ala Glu Ala Gly Gly Val Thr Gly
85 90 95
Lys Gly Gln Asp Gly Ile Gly Ser Lys Ala Glu Lys Thr Leu Gly Asp
100 105 110
Phe Ala Ala Glu Tyr Ala Lys Ser Asn Arg Ser Thr Cys Lys Gly Cys
115 120 125
Met Glu Lys Ile Glu Lys Gly Gln Val Arg Leu Ser Lys Lys Met Val
130 135 140
Asp Pro Glu Lys Pro Gln Leu Gly Met Ile Asp Arg Trp Tyr His Pro
145 150 155 160
Gly Cys Phe Val Lys Asn Arg Glu Glu Leu Gly Phe Arg Pro Glu Tyr
165 170 175
Ser Ala Ser Gln Leu Lys Gly Phe Ser Leu Leu Ala Thr Glu Asp Lys
180 185 190
Glu Ala Leu Lys Lys Gln Leu Pro Gly Val Lys Ser Glu Gly Lys Arg
195 200 205
Lys Gly Asp Glu Val Asp Gly Val Asp Glu Val Ala Lys Lys Lys Ser
210 215 220
Lys Lys Glu Lys Asp Lys Asp Ser Lys Leu Glu Lys Ala Leu Lys Ala
225 230 235 240
Gln Asn Asp Leu Ile Trp Asn Ile Lys Asp Glu Leu Lys Lys Val Cys
245 250 255
Ser Thr Asn Asp Leu Lys Glu Leu Leu Ile Phe Asn Lys Gln Gln Val
260 265 270
Pro Ser Gly Glu Ser Ala Ile Leu Asp Arg Val Ala Asp Gly Met Val
275 280 285
Phe Gly Ala Leu Leu Pro Cys Glu Glu Cys Ser Gly Gln Leu Val Phe
290 295 300
Lys Ser Asp Ala Tyr Tyr Cys Thr Gly Asp Val Thr Ala Trp Thr Lys
305 310 315 320

```

Cys Met Val Lys Thr Gln Thr Pro Asn Arg Lys Glu Trp Val Thr Pro
 325 330
 Lys Glu Phe Arg Glu Ile Ser Tyr Leu Lys Lys Leu Lys Val Lys Lys
 340 345 350
 Gln Asp Arg Ile Phe Pro Pro Glu Thr Ser Ala Ser Val Ala Ala Thr
 355 360 365
 Pro Pro Pro Ser Thr Ala Ser Ala Pro Ala Ala Val Asn Ser Ser Ala
 370 375 380
 Ser Ala Asp Lys Pro Leu Ser Asn Met Lys Ile Leu Thr Leu Gly Lys
 385 390 395 400
 Leu Ser Arg Asn Lys Asp Glu Val Lys Ala Met Ile Glu Lys Leu Gly
 405 410 415
 Gly Lys Leu Thr Gly Thr Ala Asn Lys Ala Ser Leu Cys Ile Ser Thr
 420 425 430
 Lys Lys Glu Val Glu Lys Met Asn Lys Lys Met Glu Glu Val Lys Glu
 435 440 445
 Ala Asn Ile Arg Val Val Ser Glu Asp Phe Leu Gln Asp Val Ser Ala
 450 455 460
 Ser Thr Lys Ser Leu Gln Glu Leu Phe Leu Ala His Ile Leu Ser Pro
 465 470 475 480
 Trp Gly Ala Glu Val Lys Ala Glu Pro Val Glu Val Val Ala Pro Arg
 485 490 495
 Gly Lys Ser Gly Ala Ala Leu Ser Lys Lys Ser Lys Gly Gln Val Lys
 500 505 510
 Glu Glu Gly Ile Asn Lys Ser Glu Lys Arg Met Lys Leu Thr Leu Lys
 515 520 525
 Gly Gly Ala Ala Val Asp Pro Asp Ser Gly Leu Glu His Ser Ala His
 530 535 540
 Val Leu Glu Lys Gly Gly Lys Val Phe Ser Ala Thr Leu Gly Leu Val
 545 550 555 560
 Asp Ile Val Lys Gly Thr Asn Ser Tyr Tyr Lys Leu Gln Leu Leu Glu
 565 570 575
 Asp Asp Lys Glu Asn Arg Tyr Trp Ile Phe Arg Ser Trp Gly Arg Val
 580 585 590 595
 Gly Thr Val Ile Gly Ser Asn Lys Leu Glu Gln Met Pro Ser Lys Glu
 595 600 605
 Asp Ala Ile Glu Gln Phe Met Lys Leu Tyr Glu Glu Lys Thr Gly Asn
 610 615 620
 Ala Trp His Ser Lys Asn Phe Thr Lys Tyr Pro Lys Lys Phe Tyr Pro
 625 630 635 640
 Leu Glu Ile Asp Tyr Gly Gln Asp Glu Glu Ala Val Lys Lys Leu Thr
 645 650 655
 Val Asn Pro Gly Thr Lys Ser Lys Leu Pro Lys Pro Val Gln Asp Leu
 660 665 670
 Ile Lys Met Ile Phe Asp Val Glu Ser Met Lys Lys Ala Met Val Glu
 675 680 685
 Tyr Glu Ile Asp Leu Gln Lys Met Pro Leu Gly Lys Leu Ser Lys Arg
 690 695 700
 Gln Ile Gln Ala Ala Tyr Ser Ile Leu Ser Glu Val Gln Gln Ala Val
 705 710 715 720
 Ser Gln Gly Ser Ser Asp Ser Gln Ile Leu Asp Leu Ser Asn Arg Phe
 725 730 735
 Tyr Thr Leu Ile Pro His Asp Phe Gly Met Lys Lys Pro Pro Leu Leu
 740 745 750
 Asn Asn Ala Asp Ser Val Gln Ala Lys Val Glu Met Leu Asp Asn Leu
 755 760 765
 Leu Asp Ile Glu Val Ala Tyr Ser Leu Leu Arg Gly Ser Asp Asp
 770 775 780

Ser Ser Lys Asp Pro Ile Asp Val Asn Tyr Glu Lys Leu Lys Thr Asp
 785 790 800
 Ile Lys Val Val Asp Arg Asp Ser Glu Glu Ala Glu Ile Ile Arg Lys
 805 810 815
 Tyr Val Lys Asn Thr His Ala Thr Thr His Ser Ala Tyr Asp Leu Glu
 820 825 830
 Val Ile Asp Ile Phe Lys Ile Glu Arg Glu Gly Glu Cys Gln Arg Tyr
 835 840 845
 Lys Pro Phe Lys Gln Leu His Asn Arg Arg Leu Leu Trp His Gly Ser
 850 855 860
 Arg Thr Thr Asn Phe Ala Gly Ile Leu Ser Gln Gly Leu Arg Ile Ala
 865 870 875 880
 Pro Pro Glu Ala Pro Val Thr Gly Tyr Met Phe Gly Lys Gly Ile Tyr
 885 890 895
 Phe Ala Asp Met Val Ser Lys Ser Ala Asn Tyr Tyr His Thr Ser Gln
 900 905 910
 Gly Asp Pro Ile Gly Leu Ile Leu Leu Gly Glu Val Ala Leu Gly Asn
 915 920 925
 Met Tyr Glu Leu Lys His Ala Ser His Ile Ser Arg Leu Pro Lys Gly
 930 935 940
 Lys His Ser Val Lys Gly Leu Gly Lys Thr Thr Pro Asp Pro Ser Ala
 945 950 955 960
 Asn Ile Ser Leu Asp Gly Val Asp Val Pro Leu Gly Thr Gly Ile Ser
 965 970 975
 Ser Gly Val Ile Asp Thr Ser Leu Leu Tyr Asn Glu Tyr Ile Val Tyr
 980 985 990
 Asp Ile Ala Gln Val Asn Leu Lys Tyr Leu Leu Lys Leu Lys Phe Asn
 995 1000 1005
 Phe Lys Thr Ser Leu Trp
 1010

<210> 5
 <211> 1465
 <212> DNA
 <213> Homo sapiens

<400> 5
 ggacagaggg ggacgctctc agctctctggc gcacggccca gggttatcttg tagcatagca 60
 acttcggatt tcactctacc cggagagttt cccgcttggt tgaacacatt ggccctcagga 120
 agcttccttc aaaaatgtcta ctgttcacga aatcctgtgc aagctcagct tggagggtga 180
 tcactctaca ccccacagtg catatgggtc tgctaaagcc tatactaaat ttgatgtcta 240
 cggggatgct ttgaacattg aaacagccat caagaccaaa ggtgtggatg aggtcaccat 300
 tgtcaacatt ttgaccaacc gcagcaatgc acagagacag gatattgctt tgcctacca 360
 gagaaggacc aaaaaggaaac ttgcatcagc actgaagtca gccttatctg gccacctgga 420
 gacgttgatt ttgggcoctat tgaagacacc tgctcagtat gacgctctct agctaaaagc 480
 ttccatgaag gggtcgggaa ccgacgagga ctctctcatt gagatcatct gctcagaac 540
 caaccaggag ctgcaggaaa ttaacagagt ctacaaggaa atgtacaaga ctgatctgga 600
 gaaggacatt atttcggaca catctggtag ctccgcaag ctgatgtgtg cccctggcaaa 660
 gggtagaaga cgagagtagt gctctgtcat tgattatgaa ctgattgacc aagatgctcg 720
 ggatctctat gaogctggag tgaagaggaa aggaactgat gttcccaagt ggatcagcat 780
 catgcagcag cggagcgtgc cccacctcca gaaaglatat gataggtaca agagttaacag 840
 cccatagac atgttggaaa gcacacagaa agaggttaaa ggagacctgg aaaaatgcttt 900
 cctgaacctg gttcagtgca ttcagaaaca gccctgtat tttgctgac ggctgtatga 960
 ctccatgaag ggcaaggggga cgcgagataa ggtcctgac agaactatgg tctcccgacg 1020
 tgaagtggac atgttgaaaa ttaggctctga attcaagaga aagcacggca agtccctgta 1080
 ctattatctc cagcaagaca ctaaggcgca ctaccagaaa ccgctgctgt acctgtgttg 1140
 tggagatgac tgaagccoga caggcctgta gcgtccagaa atggtgctca ccatgctctc 1200


```

agctaacagc tctagaaaaac cagcttgcca ataacagtc cctgtggccat cctgtgagg 1260
gtgaagttag cattaccccc aacotcattt tagttgccta agcattgcct ggccttctcg 1320
tctagtctct cctgtgaagcc aaagaatatga acattccaag gatttgaagc tgaagtctat 1380
gatgtgaaac actttgcctc ctgtgtactg tgtcataaac agatgaataa actgaatttg 1440
tactttaaaa aaaaaaaaaa aaaaaa                                     1465

```

```

<210> 6
<211> 339
<212> PRT
<213> Homo sapiens

```

```

<400> 6
Met Ser Thr Val His Glu Ile Leu Cys Lys Leu Ser Leu Glu Gly Asp
1 5 10 15
His Ser Thr Pro Ser Ala Tyr Gly Ser Val Lys Ala Tyr Thr Asn
20 25 30
Phe Asp Ala Glu Arg Asp Ala Leu Asn Ile Glu Thr Ala Ile Lys Thr
35 40 45
Lys Gly Val Asp Glu Val Thr Ile Val Asn Ile Leu Thr Asn Arg Ser
50 55 60
Asn Ala Gln Arg Gln Asp Ile Ala Phe Ala Tyr Gln Arg Arg Thr Lys
65 70 75 80
Lys Glu Leu Ala Ser Ala Leu Lys Ser Ala Leu Ser Gly His Leu Glu
85 90 95
Thr Val Ile Leu Gly Leu Leu Lys Thr Pro Ala Gln Tyr Asp Ala Ser
100 105 110
Glu Leu Lys Ala Ser Met Lys Gly Leu Gly Thr Asp Glu Asp Ser Leu
115 120 125
Ile Glu Ile Ile Cys Ser Arg Thr Asn Gln Glu Leu Gln Glu Ile Asn
130 135 140
Arg Val Tyr Lys Glu Met Tyr Lys Thr Asp Leu Glu Lys Asp Ile Ile
145 150 155 160
Ser Asp Thr Ser Gly Asp Phe Arg Lys Leu Met Val Ala Leu Ala Lys
165 170 175
Gly Arg Arg Ala Glu Asp Gly Ser Val Ile Asp Tyr Glu Leu Ile Asp
180 185 190
Gln Asp Ala Arg Asp Leu Tyr Asp Ala Gly Val Lys Arg Lys Gly Thr
195 200 205
Asp Val Pro Lys Trp Ile Ser Ile Met Thr Glu Arg Ser Val Pro His
210 215 220
Leu Gln Lys Val Phe Asp Arg Tyr Lys Ser Tyr Ser Pro Tyr Asp Met
225 230 235 240
Leu Glu Ser Ile Arg Lys Glu Val Lys Gly Asp Leu Glu Asn Ala Phe
245 250 255
Leu Asn Leu Val Gln Cys Ile Gln Asn Lys Pro Leu Tyr Phe Ala Asp
260 265 270
Arg Leu Tyr Asp Ser Met Lys Gly Lys Gly Thr Arg Asp Lys Val Leu
275 280 285
Ile Arg Ile Met Val Ser Arg Ser Glu Val Asp Met Leu Lys Ile Arg
290 295 300
Ser Glu Phe Lys Arg Lys Tyr Gly Lys Ser Leu Tyr Tyr Ile Gln
305 310 315 320
Gln Asp Thr Lys Gly Asp Tyr Gln Lys Ala Leu Leu Tyr Leu Cys Gly
325 330 335
Gly Asp Asp

```

11

<210> 7
 <211> 1362
 <212> DNA
 <213> Homo sapiens

<400> 7
 catttgggga cgtctctcagc tctcggcgca cggcccagct tctctcaaaa tgtctactgt 60
 tcacgaaatc ctgtgcaagc tcagcttgga ggggtgatcac tctacacccc caagtgcata 120
 tgggtctgtc aaagcctata ctaactttga tgcctgagcg gatgctttga acattgaaac 180
 agccatcaag accaaaggtg tggatgaggt caccattgtc aacattttga ccaaccgcag 240
 caatgcacag agacaggata ttgccttcgc ctaccagaga aggacaaaa aggaacttgc 300
 atcagcactg aagtcagcct tatctggcca cctggagacg gtgattttgg gcctattgaa 360
 gacacctgct cagtatgacg ctctcgagct aaaagcttcc atgaaaggcg tgggaaccga 420
 cgaggactct ctcaattgaga tcatctgctc cagaaccaac caggagctgc aggaatttaa 480
 cagagtctac aaggaaatgt acaagactga tctggagaag gacattattt cggacacatc 540
 tggtagcttc cgcaagctga tgggtgccct ggcaagggtt agaagagcag aggaatggctc 600
 tglcatgat tatgaaactga ttgaccaaga tgcctgggat ctctatgacg ctggagtgaa 660
 gaggaaggga actgatgttc ccaagtggat cagcatcatg accgagcgga gcgtgcccca 720
 cctccagaaa gtatttgata ggtacaagag ttacagccct tatgacatgt tggaaagcat 780
 caggaagag gttaaaggag acctggaaaa tgccttctcg aacctggctc agtgcatcca 840
 gaacaagccc ctgtattttg ctgatcggtc gtatgactcc atgaagggca aggggacgcg 900
 agataaggct ctgatcagaa tcatggtctc ccgcagtga gtggacatgt tgaaaattag 960
 tctctgaattc aagagaaggt acggcaagtc cctgtactat tataccagc aagacactaa 1020
 gggcgactac cagaagaagc tgctgtacct gtgtggtgga gatgactgaa gcccgacagc 1080
 gctgagcgt ccagaaatgg tgctcaccat gcttcacagt aacaggtcta gaaaaccagc 1140
 ttgcgaaatg cagctcccggt ggccatccct gtgagggtga cgttagcatt acccccacac 1200
 tcaatttagt tgccctaagca ttgctgtgcc ttctgtctta gtctctctcg taagccaaag 1260
 aaatgaacat tccaaggagt tgggaagtga gtctatgat tgaaacactt tgcctcctgt 1320
 gtactgtgtc ataaacagat gaataaactg aatttgtact tt 1362

<210> 8
 <211> 339
 <212> PRT
 <213> Homo sapiens

<400> 8
 Met Ser Thr Val His Glu Ile Leu Cys Lys Leu Ser Leu Glu Gly Asp
 1 5 10 15
 His Ser Thr Pro Pro Ser Ala Tyr Gly Ser Val Lys Ala Tyr Thr Asn
 20 25 30
 Phe Asp Ala Glu Arg Asp Ala Leu Asn Ile Glu Thr Ala Ile Lys Thr
 35 40 45
 Lys Gly Val Asp Glu Val Thr Ile Val Asn Ile Leu Thr Asn Arg Ser
 50 55 60
 Asn Ala Gln Arg Gln Asp Ile Ala Phe Ala Tyr Gln Arg Arg Thr Lys
 65 70 75 80
 Lys Glu Leu Ala Ser Ala Leu Lys Ser Ala Leu Ser Gly His Leu Glu
 85 90 95
 Thr Val Ile Leu Gly Leu Leu Lys Thr Pro Ala Gln Tyr Asp Ala Ser
 100 105 110
 Glu Leu Lys Ala Ser Met Lys Gly Leu Gly Thr Asp Glu Asp Ser Leu
 115 120 125
 Ile Glu Ile Ile Cys Ser Arg Thr Asn Gln Glu Leu Gln Glu Ile Asn
 130 135 140
 Arg Val Tyr Lys Glu Met Tyr Lys Thr Asp Leu Glu Lys Asp Ile Ile
 145 150 155 160
 Ser Asp Thr Ser Gly Asp Phe Arg Lys Leu Met Val Ala Leu Ala Lys
 165 170 175

Gly Arg Arg Ala Glu Asp Gly Ser Val Ile Asp Tyr Glu Leu Ile Asp
 180 185 190
 Gln Asp Ala Arg Asp Leu Tyr Asp Ala Gly Val Lys Arg Lys Gly Thr
 195 200 205
 Asp Val Pro Lys Trp Ile Ser Ile Met Thr Glu Arg Ser Val Pro His
 210 215 220
 Leu Gln Lys Val Phe Asp Arg Tyr Lys Ser Tyr Ser Pro Tyr Asp Met
 225 230 235 240
 Leu Glu Ser Ile Arg Lys Glu Val Lys Gly Asp Leu Glu Asn Ala Phe
 245 250 255
 Leu Asn Leu Val Gln Cys Ile Gln Asn Lys Pro Leu Tyr Phe Ala Asp
 260 265 270
 Arg Leu Tyr Asp Ser Met Lys Gly Lys Gly Thr Arg Asp Lys Val Leu
 275 280 285
 Ile Arg Ile Met Val Ser Arg Ser Glu Val Asp Met Leu Lys Ile Arg
 290 295 300
 Ser Glu Phe Lys Arg Lys Tyr Gly Lys Ser Leu Tyr Tyr Tyr Ile Gln
 305 310 315 320
 Gln Asp Thr Lys Gly Asp Tyr Gln Lys Ala Leu Leu Tyr Leu Cys Gly
 325 330 335
 Gly Asp Asp

<210> 9

<211> 1982

<212> DNA

<213> Homo sapiens

<400> 9

gcagaggagg agcgcacgcc ggcctcgaag aactttctgt tgggtggctg aaactctgac 60
 ttgaactaga gtcattggcga tggcaaccaa aggaggtagt gtcaaaagctg cttcaggatt 120
 caatgccatg gaagatgccc agaccctgag gaaggccatg aaagggctcg gccccgatga 180
 agacgccatt attagcgtcc ttgcctaccg caacaccccg cagcgccagg agatcaggac 240
 agcctacaag agcaaccatcg gcagggaact gatagacgac ctgaagtcag aactgagtg 300
 caacttcgag caggtgattg tggggatgat gacgcocag gtgctgtatg acgtgcaaga 360
 gctgcgaagg gccatgaagg gagcgcgcac tgatgagggc tgcccaattg agatcctggc 420
 ctccoggagcc cctgaggaga tccggcgcat aagccaaacc taccagcagc aaatgtggac 480
 gagccttgaa gatgacattc gctctgacac atcgttcatg ttccagcgag tgctgggtgc 540
 tctgtcagct ggtgggaggg atgaaggaaa ttatctggac gatgctctcg tgagacagga 600
 tgccagggac ctgtatgagg ctggagagaa gaaatggggg acagatgagg tgaattctc 660
 aactgttctc tgttcccggg accgaaatca cctgttgcat gtgtttgatg aatacaaaaag 720
 gatatacagc aaggatattg aacagagtat taaatctgaa acatctgtga gctttgaaga 780
 tgctctgctg gctatagtaa agtgcatgag gaacaaatct gcataatttg ctgaaaagct 840
 ctataaatcg atgaagggtc tgggcaccca tgatacaccc ctcatcagag tgatggtttc 900
 tcgagcagaa attgacatgt tggatatccg ggacacctc aagagactct atggaaagtc 960
 tctgtactcg ttoatcaagg gtgacacatc tggagactac aggaaagtac tggttgttct 1020
 ctgtggagga gatgattaaa ataaaaatcc cagaaggaca gaggagttct caacactttg 1080
 aattttttta acttcaattt tctacactgc tattatcatt atctcagaat gcttatcttc 1140
 aattaaaaag cctacaagctg cctcctagaa tatagactgt ctgtattatt attcaactat 1200
 aattagtcac tatgatgctt taaagctgta cttgcatttc aaagcttata agatataaat 1260
 ggagatttta aagtataaat aaatatgtat tccatgtttt taaaagatta cttctcactt 1320
 tggattttcc agacatgtga tatattaaat tattccattt ttctttttta gtgaaaattt 1380
 ttttaaatgg aagactgttc taaaatcaact tttttcccta atccaatttt tagagtggct 1440
 agtattttct tcatittgaa ttgtaaagca ccggtcagta gaagtcgcca tctcagttttc 1500
 tatatttcat agtccaaagcc ttgaaaagcat ctacaaatct ctttttttag gttttgtcca 1560
 tagcatcagt tgatccttca taagtttttc atgggagact tcttccatca catcttatgt 1620
 tgaatcaact tctgttagtc aaagtatacc aaacccaatt tatctgaact aaattctaaa 1680

```

gtatgggttat acaaaccata tacatctggt taccaaacat aaatgctgaa cattccatat 1740
tattatagtt aatgtcttaa tccagcttgc aagtgaatgg aaaaaaaat aagcttcaaa 1800
ctaggtattc tgggaatgat gtaatgctct gaatttagta tgatataaag aaaacttttt 1860
tgtgtcaaaa atacttttta aaatcaattt tgttgattgt agtaattctt atttgcactg 1920
tgccittcaa ctccagaaac attctaagat gtacttggat ttaattaaaa agttcacttt 1980
gt

```

```

<210> 10
<211> 321
<212> PRT
<213> Homo sapiens

```

```

<400> 10
Met Ala Met Ala Thr Lys Gly Gly Thr Val Lys Ala Ala Ser Gly Phe
1 5 10 15
Asn Ala Met Glu Asp Ala Gln Thr Leu Arg Lys Ala Met Lys Gly Leu
20 25 30
Gly Thr Asp Glu Asp Ala Ile Ile Ser Val Leu Ala Tyr Arg Asn Thr
35 40 45
Ala Gln Arg Gln Glu Ile Arg Thr Ala Tyr Lys Ser Thr Ile Gly Arg
50 55 60
Asp Leu Ile Asp Asp Leu Lys Ser Glu Leu Ser Gly Asn Phe Glu Gln
65 70 75 80
Val Ile Val Gly Met Met Thr Pro Thr Val Leu Tyr Asp Val Gln Glu
85 90 95
Leu Arg Arg Ala Met Lys Gly Ala Gly Thr Asp Glu Gly Cys Leu Ile
100 105 110
Glu Ile Leu Ala Ser Arg Thr Pro Glu Glu Ile Arg Arg Ile Ser Gln
115 120 125
Thr Tyr Gln Gln Gln Tyr Gly Arg Ser Leu Glu Asp Asp Ile Arg Ser
130 135 140
Asp Thr Ser Phe Met Phe Gln Arg Val Leu Val Ser Leu Ser Ala Gly
145 150 155 160
Gly Arg Asp Glu Gly Asn Tyr Leu Asp Asp Ala Leu Val Arg Gln Asp
165 170 175
Ala Gln Asp Leu Tyr Glu Ala Gly Glu Lys Lys Trp Gly Thr Asp Glu
180 185 190
Val Lys Phe Leu Thr Val Leu Cys Ser Arg Asn Arg Asn His Leu Leu
195 200 205
His Val Phe Asp Glu Tyr Lys Arg Ile Ser Gln Lys Asp Ile Glu Gln
210 215 220
Ser Ile Lys Ser Glu Thr Ser Gly Ser Phe Glu Asp Ala Leu Leu Ala
225 230 235 240
Ile Val Lys Cys Met Arg Asn Lys Ser Ala Tyr Phe Ala Glu Lys Leu
245 250 255
Tyr Lys Ser Met Lys Gly Leu Gly Thr Asp Asn Thr Leu Ile Arg
260 265 270
Val Met Val Ser Arg Ala Glu Ile Asp Met Leu Asp Ile Arg Ala His
275 280 285
Phe Lys Arg Leu Tyr Gly Lys Ser Leu Tyr Ser Phe Ile Lys Gly Asp
290 295 300
Thr Ser Gly Asp Tyr Arg Lys Val Leu Leu Val Leu Cys Gly Gly Asp
305 310 315 320
Asp

```

```

<210> 11

```

14

<211> 1316
 <212> DNA
 <213> Homo sapiens

<400> 11
 agctagacgc cccgaggctg gactgaagcg ccgggacoga gcccgtctc ccaggagatc 60
 cggggcgcac ggcacgcgag agagcgcggg agccaacctg ggccatcat gcgcagggcc 120
 cgggacgctg ggccggtcta caccgcgcgc tgggtcacgt ggccgggacg ggccggcggc 180
 tgccccgcgc ggggggcggg ggtcgcccg gggttgcgct ggacgacoga gacggcgggg 240
 ccgcgagcgg cctggagcct cccaacccgc gccgcgctgg cctcgagcgg taggagccgc 300
 cccctgcccc cccgcgcggg cccgcgcgcc gccgcgcgcg cccctatata gcgcgcccca 360
 cgagggcgcg cgcgcaggcg ccagcctcgg agtggggcgg ggacagtgg ccgcgccccg 420
 cagccaggcc cccgcgcgcg ccgcatccac ctctccgcc gcccgcgacc caacggcggc 480
 cccccgcggg cagctgcgcg cgggccccgc cggccaccat gaagaaggag gtgtgctccg 540
 tggccttcct caaggccgtg ttgcgaaggt tcttggccac cctcatcttc gtcttctttg 600
 gcctgggctc ggccctcaa tggccgtcgg cgtgctcac catcctgca atcgcgctgg 660
 cgtttgcctt ggccatagg acgctggccc aggcctctgg acccgtgagc ggccggccca 720
 tcaacccccc catcacctg gccctcttgg tgggcaacca gatctcgctg ctccgggctt 780
 tctcttaagt gggcggccag ctgtgtgggg ccattgcggg ggctggcacc ctctacggtg 840
 tggcacgcct caatgccggg ggcaatctgg ccgtcaacgc gctcaacaac aacacaaacg 900
 agggccaggc catgtgtgtg gactgattc tgaccttcca gctggcactc tgcactcttcg 960
 cctccactga cctccgcgcg accagcctg tgggctcccc agccctgtcc atggcctgt 1020
 ctgtcaacct ggccacactt gtccgaatct acttcaactg ctgctccatg aaccagcccc 1080
 gctctttttg cctcgcggtg gtcaatgaat ggttcagccc cgtcactcgg gtttcttggg 1140
 tagggcccat cgtggggcgg gtccctgctg ccactcctta ctctacactg cttctcccca 1200
 actcctcgag ccttgagtgag cgtgtggcca tcatcaagg cactgatgag cctgcaggag 1260
 actgggagga gcagcgggaa gagcgggaaga agaccatgga gctgaccacc cgtctga 1316

<210> 12
 <211> 265
 <212> PRT
 <213> Homo sapiens

<400> 12
 Met Lys Lys Glu Val Cys Ser Val Ala Phe Leu Lys Ala Val Phe Ala
 1 5 10 15
 Glu Phe Leu Ala Thr Leu Ile Phe Val Phe Phe Gly Leu Gly Ser Ala
 20 25 30
 Leu Lys Trp Pro Ser Ala Leu Pro Thr Ile Leu Gln Ile Ala Leu Ala
 35 40 45
 Phe Gly Leu Ala Ile Gly Thr Leu Ala Gln Ala Leu Gly Pro Val Ser
 50 55 60
 Gly Gly His Ile Asn Pro Ala Ile Thr Leu Ala Leu Leu Val Gly Asn
 65 70 75 80
 Gln Ile Ser Leu Leu Arg Ala Phe Phe Tyr Val Ala Ala Gln Leu Val
 85 90 95
 Gly Ala Ile Ala Gly Ala Gly Ile Leu Tyr Gly Val Ala Pro Leu Asn
 100 105 110
 Ala Arg Gly Asn Leu Ala Val Asn Ala Leu Asn Asn Asn Thr Thr Gln
 115 120 125
 Gly Gln Ala Met Val Val Glu Leu Ile Leu Thr Phe Gln Leu Ala Leu
 130 135 140
 Cys Ile Phe Ala Ser Thr Asp Ser Arg Arg Thr Ser Pro Val Gly Ser
 145 150 155 160
 Pro Ala Leu Ser Ile Gly Leu Ser Val Thr Leu Gly His Leu Val Gly
 165 170 175
 Ile Tyr Phe Thr Gly Cys Ser Met Asn Pro Ala Arg Ser Phe Gly Pro
 180 185 190

15

Ala	Val	Val	Met	Asn	Arg	Phe	Ser	Pro	Ala	His	Trp	Val	Phe	Trp	Val
	195						200					205			
Gly	Pro	Ile	Val	Gly	Ala	Val	Leu	Ala	Ala	Ile	Leu	Tyr	Phe	Tyr	Leu
	210					215					220				
Leu	Phe	Pro	Asn	Ser	Leu	Ser	Leu	Ser	Glu	Arg	Val	Ala	Ile	Ile	Lys
	225				230				235						240
Gly	Thr	Tyr	Glu	Pro	Asp	Glu	Asp	Trp	Glu	Glu	Gln	Arg	Glu	Glu	Arg
			245					250						255	
Lys	Lys	Thr	Met	Glu	Leu	Thr	Thr	Arg							
			260					265							

<210> 13
 <211> 1653
 <212> DNA
 <213> Homo sapiens

<400> 13
 acgtccgggg agggggccagg tgagcggcag acccggcag caggtggggg ccggcggggt 60
 cgtgtggcag agctgcagag agacaaggcg gggcggtgt cgtgtgtggg tgcagtggag 120
 aagaggccct cgtgtgtgccc catggctggc caggatcctg cgtgtgacac gactcaccgc 180
 ttctacgacg tggccagaca tggcattctg caggtggcag gggatgaccg ctttgggaaga 240
 cgtgttgtca cgttcagctg ctgcgggatg ccgccctccc acgagctgga ccaccagcgg 300
 ctgctggagt atttgaagta cacactggac caatacgttg agaacgatta taccatcgtc 360
 tatttccaac acgggctgaa cagccgggaa aagccttccc tgggctgggt ccagagcgca 420
 tacaaggagt tcgataggaa agacggggat ctcaactatgt ggcccaggct ggtctcgaaac 480
 tccaagctca agcgatcctc ccactcagc ctcccaagt actgggatta caggtacaag 540
 aagaacttga aggcctctta cgtggtgcac cccaccagct tcactcaaggt cctgtggaaac 600
 atcttgaagc cctcatcag tcacaagttt gggaagaaag tcactctattt caactacctg 660
 agttagctcc acgaacaact taaatacagc cagctgtgtca tccctcccgca agttttgcgg 720
 tacgatgaga agctccagag cctgcaacgag ggccgggaagc cgctctctac caagacacca 780
 ccgcgcgggc ccccgctgccc cacacagcag tttggcgtca gtctgcaata cctcaaaagac 840
 aaaaatcaag gcgaactcat ccccccgtgt ctgagggttca cagtgaagta cctgagagag 900
 aaaggcctgc gcaccgaggg cctgttccgg agatccgccca gcgtgcagac cgtccgcgag 960
 atccagaggg tctacaacca agggaaagccc gtgaactttg acgactacgg ggacattcac 1020
 atccctcgcc tgatcctgaa gaccttctcg cgagagctgc ccagccctcg tctgaccttc 1080
 caggcctacg agcagattct cgggatcacc tgtgtggaga gcagcctcg tgtcaactggc 1140
 tggcccgaga tcttaccgag cctcccagag cacaactacg tctgtctcgg ctacctcatg 1200
 gctcttctgc atgcggtgtc ccgggagagc acttccaaca aaatgaacag ctctaacctg 1260
 gctgtgtgtc tcgggctgaa tttgatctgg ccattcccag ggtgtctctc cctgagtgcc 1320
 ctgtgtcccc tgaacatggt cactgaactg ctgatcgagt acatgaaaa gatcttcagc 1380
 accccggagg caactgggga gcacggcctg gccactagg aacaggggag cagggcagcc 1440
 cctttgcagg aggtgtgccc acggacacaa gccacgggcc taccacaagc taccctacct 1500
 ccgagctccc tgatggcagc cagaagacgt ctctagtgtt ggaacaactc tgbatgttct 1560
 gagctacctc ccacacctgt ctgtgcactt gtatgttttg taaacttggc atctgtaaaa 1620
 ataaccagcc attagatgaa ttcagaacct tct 1653

<210> 14
 <211> 464
 <212> PRT
 <213> Homo sapiens

<400> 14
 Met Ala Gly Gln Asp Pro Ala Leu Ser Thr Ser His Pro Phe Tyr Asp
 1 5 10 15
 Val Ala Arg His Gly Ile Leu Gln Val Ala Gly Asp Asp Arg Phe Gly
 20 25 30

<212> DNA

<213> Homo sapiens

<400> 15

```

atgaggactc  tccgcaggtt  gaagttcatg  agttcgccca  gcctcagtag  cctggggcaag  60
agagagccgg  ccgcgcgcgc  ggaagcagcg  ggcacgcagc  agcgcccgcc  ctgcccacac  120
acagacctga  acagatccca  caacgggggt  atgcgcgtct  tccagcgcaa  ggggctgccc  180
gaccagagag  tcttcagcct  caacgagggc  gtcgcggcag  tgttgaagac  agagctgggg  240
tcctctttca  cggtagctac  gcagaaccag  cgtctgacaa  aaggcatlgt  gatccttcgg  300
gacaaatgac  gcttctatga  gggacagaag  ctgctggact  cactggcaga  gacctgggac  360
gtgacgtggt  gtgacgtgct  gccatgctg  caggccatct  tctaccoggt  gcagggcaag  420
gagccatcgg  tgcgccagct  gccctgctg  cacttcogga  atgccatcac  cctcagtggt  480
aagctagagg  atgcgtggc  ccgggcccat  gccctgtgct  ccctgccat  cgtgcagatg  540
ctgctgtgtg  tgcagggggt  acatgagtc  agggcgctga  ctgaggacta  cctgcgctgt  600
gagacgtggt  tccagaaggt  ggtgtcgcca  tacctgggca  cctacggcct  cactccagc  660
gaggggcccc  tccccattc  ctgcatcctg  gagctgcaga  gagacaaggc  ggcggcggt  720
gctgtgtgtg  gtgcagtgag  gaagaggccc  tcgggtgtgc  ccattggctg  ccagatcct  780
gcgtcgagca  cgaagtaccc  gttctacgac  gtggccagac  atggcattct  gcaggtggca  840
gggatgacc  gctttggaag  acgtgtgtc  acgttcagct  gctgccggat  gccaccctcc  900
cacagctgtg  accacnagcg  cgtgctggag  tcaagaaga  acttgaagcg  cctctacgtg  960
gtgcacccca  ccagcttcat  caaggtcctg  tggaaactct  tgaagccct  catcagtcac  1020
aagtttggga  agaaagtcac  ctatttcaac  tactctgagt  agctccagca  acactttaa  1080
tacgacacgc  ttgtctatcc  tccgaagtt  ttgcggtacg  atgagaagct  ccagagcctg  1140
cacagagggc  ggacgcgcgc  tccacacaa  accacacgc  ccggccccc  cgtgcccca  1200
cagcagtttg  gcgtcagctc  gcaatacct  aaagacaaa  atcaaggcga  actcatcccc  1260
ctctgtctga  ggttcacagt  gacgtacct  agagagaaa  gctctgcgac  cgaggcgctg  1320
ttccggagat  ccgcccagct  gcagaccgt  cgcgagatcc  agaggctcta  caaccaaggg  1380
aagccgtgga  actttgacga  ctacggggac  attcacatc  ctgcctgtat  ctgaagacc  1440
ttcctgcgag  agctgcccc  gcccttctg  accttcagg  cctacgagca  gattctcgg  1500
atcacctgtg  ttgagagcag  cctgcgtgtc  actggctgcc  gccagatctt  accgagcctc  1560
ccagagcaca  actacgtcgt  cctcogctac  ctcatgggt  tcttgcatgc  ggtgtccgg  1620
gagagcatct  tcaacaaaat  gaacagctct  aacctggcct  gtgtcttcgg  gctgaatttg  1680
atctggccat  ccaggggggt  ctctccctg  agtgcccttg  tgccctgaa  catgttact  1740
gaactgtgta  tcgagtacta  tgaaaagatc  ttccagccac  ccgaggcacc  tggggagcac  1800
ggcctggcac  catgggaaca  ggggagcagg  gcagccctt  tgcaggagc  tgtgccaccg  1860
acacaagcca  cgggctcacc  caagcctacc  ctacctccga  gtccctgat  ggagccacga  1920
agacgtctct  agtgttgcca  acactctgta  tatttcgagc  tacctccacc  acctgtctgt  1980
gcacttgat  gttttgtaaa  ctggcatct  gtaaaaataa  ccagccatta  gatgaattca  2040
gaa  2043

```

<210> 16

<211> 643

<212> PRT

<213> Homo sapiens

<400> 16

```

Met Arg Thr  Leu Arg Arg  Leu Lys Phe  Met Ser Ser  Pro Ser Leu Ser
1           5           10          15
Asp Leu Gly  Lys Arg Glu  Pro Ala Ala  Ala Asp Glu Arg Gly Thr
20          25          30          35
Gln Gln Arg  Arg Ala Cys  Ala Asn Ala  Thr Trp Asn Ser Ile His Asn
35          40          45          50
Gly Val Ile  Ala Val Phe  Gln Arg Lys  Gly Leu Pro Asp Gln Glu Leu
50          55          60          65
Phe Ser Leu  Asn Glu Gly Val Arg  Gln Leu Leu  Lys Thr Glu Leu Gly
65          70          75          80
Ser Phe Phe  Thr Glu Tyr  Leu Gln Asn  Gln Leu Leu  Thr Lys Gly Met
85          90          95

```

Val Ile Leu Arg Asp Lys Ile Arg Phe Tyr Glu Gly Gln Lys Leu Leu
 100 105 110
 Asp Ser Leu Ala Glu Thr Trp Asp Phe Phe Phe Ser Asp Val Leu Pro
 115 120 125
 Met Leu Gln Ala Ile Phe Tyr Pro Val Gln Gly Lys Glu Pro Ser Val
 130 135 140
 Arg Gln Leu Ala Leu Leu His Phe Arg Asn Ala Ile Thr Leu Ser Val
 145 150 155 160
 Lys Leu Glu Asp Ala Leu Ala Arg Ala His Ala Arg Val Pro Pro Ala
 165 170 175
 Ile Val Gln Met Leu Leu Val Leu Gln Gly Val His Glu Ser Arg Gly
 180 185 190
 Val Thr Glu Asp Tyr Leu Arg Leu Glu Thr Leu Val Gln Lys Val Val
 195 200 205
 Ser Pro Tyr Leu Gly Thr Tyr Gly Leu His Ser Ser Glu Gly Pro Phe
 210 215 220
 Thr His Ser Cys Ile Leu Glu Leu Gln Arg Asp Lys Ala Ala Ala Ala
 225 230 235 240
 Ala Val Leu Gly Ala Val Arg Lys Arg Pro Ser Val Val Pro Met Ala
 245 250 255
 Gly Gln Asp Pro Ala Leu Ser Thr Ser His Pro Phe Tyr Asp Val Ala
 260 265 270
 Arg His Gly Ile Leu Gln Val Ala Gly Asp Asp Arg Phe Gly Arg Arg
 275 280 285
 Val Val Thr Phe Ser Cys Cys Arg Met Pro Pro Ser His Glu Leu Asp
 290 295 300
 His Gln Arg Leu Leu Glu Tyr Lys Lys Asn Leu Lys Ala Leu Tyr Val
 305 310 315 320
 Val His Pro Thr Ser Phe Ile Lys Val Leu Trp Asn Ile Leu Lys Pro
 325 330 335
 Leu Ile Ser His Lys Phe Gly Lys Lys Val Ile Tyr Phe Asn Tyr Leu
 340 345 350
 Ser Glu Leu His Glu His Leu Lys Tyr Asp Gln Leu Val Ile Pro Pro
 355 360 365
 Glu Val Leu Arg Tyr Asp Glu Lys Leu Gln Ser Leu His Glu Gly Arg
 370 375 380
 Thr Pro Pro Pro Thr Lys Thr Pro Pro Pro Arg Pro Pro Leu Pro Thr
 385 390 395 400
 Gln Gln Phe Gly Val Ser Leu Gln Tyr Leu Lys Asp Lys Asn Gln Gly
 405 410 415
 Glu Leu Ile Pro Pro Val Leu Arg Phe Thr Val Thr Tyr Leu Arg Glu
 420 425 430
 Lys Gly Leu Arg Thr Glu Gly Leu Phe Arg Arg Ser Ala Ser Val Gln
 435 440 445
 Thr Val Arg Glu Ile Gln Arg Leu Tyr Asn Gln Gly Lys Pro Val Asn
 450 455 460
 Phe Asp Asp Tyr Gly Asp Ile His Ile Pro Ala Val Ile Leu Lys Thr
 465 470 475 480
 Phe Leu Arg Glu Leu Pro Gln Pro Leu Leu Thr Phe Gln Ala Tyr Glu
 485 490 495
 Gln Ile Leu Gly Ile Thr Cys Val Glu Ser Ser Leu Arg Val Thr Gly
 500 505 510
 Cys Arg Gln Ile Leu Arg Ser Leu Pro Glu His Asn Tyr Val Val Leu
 515 520 525
 Arg Tyr Leu Met Gly Phe Leu His Ala Val Ser Arg Glu Ser Ile Phe
 530 535 540
 Asn Lys Met Asn Ser Ser Asn Leu Ala Cys Val Phe Gly Leu Asn Leu
 545 550 555 560

Ile	Trp	Pro	Ser	Gln	Gly	Val	Ser	Ser	Leu	Ser	Ala	Leu	Val	Pro	Leu
				565					570					575	
Asn	Met	Phe	Thr	Glu	Leu	Leu	Ile	Glu	Tyr	Tyr	Glu	Lys	Ile	Phe	Ser
			580					585					590		
Thr	Pro	Glu	Ala	Pro	Gly	Glu	His	Gly	Leu	Ala	Pro	Trp	Glu	Gln	Gly
		595					600					605			
Ser	Arg	Ala	Ala	Pro	Leu	Gln	Glu	Ala	Val	Pro	Arg	Thr	Gln	Ala	Thr
	610					615					620				
Gly	Leu	Thr	Lys	Pro	Thr	Leu	Pro	Pro	Ser	Pro	Leu	Met	Ala	Ala	Arg
	625				630					635					640
Arg	Arg	Leu													

<210> 17
 <211> 2274
 <212> DNA
 <213> Homo sapiens

<400> 17
 atgaggactc tccgcagggt gaagttcatg agttcgccca gcctcagtg cctgggcaag 60
 agagagcccg ccgcccgcgc ggacgagcgg ggcacgcagc agcgccggg ctgcgccaac 120
 gccacctcga acagcatcca caacgggggt atcgccgtct tccagcgcaa ggggctgccc 180
 gaccaggagc tcttcagcct caacgagggc gtccggcagc tgttgaagac agagctgggg 240
 tctctcttca cggagtaact cagaaaccaa ctgctgacaa aagcatgggt gatcctcgg 300
 gacaagattc gcttctatga gggacagaag ctgctggact cactggcaga gaactggggc 360
 ttctcttcca gtgagtgct gccaatgctg caggccatct tctaccoggt gcagggcaag 420
 gagccatcgg tgcgcagctt ggcctgctg caottccgga atgccatcac cctcagtggt 480
 aagctagagg atgcgtggc ccgggcccac gccctgtgct cccctgccat cgtgcagatg 540
 ctgctgtgtg tgcagggggt acatgagtc agggcggtga ctgaggacta cctgcgcctg 600
 gagacgtggg tccagaaggt ggtgtcgcca tacctgggca cctacggcct ccactccagc 660
 gaggggcccc tcacccattc ctgcatcctg gagctgcaga gagacaagg gcggcgggct 720
 gctgtgctgg gtgcagttag gaagaggccc tccgtggtgc ccatggctgg ccaggatcct 780
 cgctgagaga cagtcacccc gttctacgac gtggccagac atggcatctt gcaggtygca 840
 ggggatgacc gctttggaag acgtgttgct acgttcagct gctgccggat gccaccctcc 900
 cacgagctgg accaccagcg cctgctggag tatttgaagt acacactgga ccaataogtt 960
 gagaacgatt ataccatcgt ctatttccac tacgggctga acagccggaa caagccttcc 1020
 ctgggctggc tccagagcgc atacaaggag ttcatagga aagacgggga tctcactatg 1080
 tggccacagg tggctctgaa ctccaaagtc aagcgtacct ccacactcac cctcccaaat 1140
 tactgggatt acaggtacaa gaagaaactg aaggccctct acgtggtgca cccaccacag 1200
 ttcattcaag tctgttgaaa catcttgaag cccctcatca gtccacaagt tgggaagaaa 1260
 gtcatctatt ttaactacat gagttagctc acgaaacacc ttaaatacga tcaactggctc 1320
 atccctcccg aagtttttgc gtaacgttag aagctccaga gctgtcacga gggccggagc 1380
 ccgctcccca ccaagacacc accgcgcggc ccccgctgca cccacacaga gtttggcgct 1440
 agtctcaaat acctcaaaaga caaaaatcaa ggcgaactca tccccctgtg gctgaggttc 1500
 acagtgcagt acctgacaga gaaaggcctg cgcaccgagc gctgttccg gctgacogcc 1560
 agcgtgcaga ccgtccggga gatccagagg ctctacaacc aagggaagcc cgtgaacttt 1620
 gagactacg gggacattca catccctgcc gtgactccta agaccttctc gcgagatcgt 1680
 cccacgcggc ttctgacctt ccaggcctac gagcagattc tcgggatcac ctgtgtggag 1740
 agcagcctgc gtgtcactgg ctgcgcggag atcttaacga gcctccaga gcaacaactc 1800
 gtgctcctcc gctacotcat gggcttctgt catgcgggtg cccgggagag catcttcaac 1860
 aaaatgaaca gctctaacctt ggcctgtgtc ttccgggtga atttgtatgt gccatccagc 1920
 ggggtctcct cctgtactgc cctgtgtccc ctgaactcgt tcaactgaat gctgatcgag 1980
 tactatgaaa agatcttcag caccocggag gacctctggg agcacggcct ggacccatgg 2040
 gaacagggga gcaggtcgag cccctttgac caccggacac agccacgggc 2100
 ctccacaagc ctaccctacc tccagtcctc ctgactggcg ccagagagac tctctagtgt 2160
 tgcgaacaot ctgttatatt cagactacct cctgtgcact tgtatgtttt 2220
 gtaaacctgg catctgtaaa aataaccagc cattagatga attcagaac tttc 2274

<210> 18
 <211> 751
 <212> PRT
 <213> Homo sapiens

<220>
 <221> VARIANT
 <222> (1)...(751)
 <223> Xaa = Any Amino Acid

<400> 18
 Met Arg Thr Leu Arg Arg Leu Lys Phe Met Ser Ser Pro Ser Leu Ser
 1 5 10 15
 Asp Leu Gly Lys Arg Glu Pro Ala Ala Asp Glu Arg Gly Thr
 20 25 30
 Gln Gln Arg Arg Ala Cys Ala Asn Ala Thr Trp Asn Ser Ile His Asn
 35 40 45
 Gly Val Ile Ala Val Phe Gln Arg Lys Gly Leu Pro Asp Gln Glu Leu
 50 55 60
 Phe Ser Leu Asn Glu Gly Val Arg Gln Leu Leu Lys Thr Glu Leu Gly
 65 70 75 80
 Ser Phe Phe Thr Glu Tyr Leu Gln Asn Gln Leu Leu Thr Lys Gly Met
 85 90 95
 Val Ile Leu Arg Asp Lys Ile Arg Phe Tyr Glu Gly Gln Lys Leu Leu
 100 105 110
 Asp Ser Leu Ala Glu Thr Trp Asp Phe Phe Phe Ser Asp Val Leu Pro
 115 120 125
 Met Leu Gln Ala Ile Phe Tyr Pro Val Gln Gly Lys Glu Pro Ser Val
 130 135 140
 Arg Gln Leu Ala Leu Leu His Phe Arg Asn Ala Ile Thr Leu Ser Val
 145 150 155 160
 Lys Leu Glu Asp Ala Leu Ala Arg Ala His Ala Arg Val Pro Pro Ala
 165 170 175
 Ile Val Gln Met Leu Leu Val Leu Gln Gly Val His Glu Ser Arg Gly
 180 185 190
 Val Thr Glu Asp Tyr Leu Arg Leu Glu Thr Leu Val Gln Lys Val Val
 195 200 205
 Ser Pro Tyr Leu Gly Thr Tyr Gly Leu His Ser Ser Glu Gly Pro Phe
 210 215 220
 Thr His Ser Cys Ile Leu Glu Leu Gln Arg Asp Lys Ala Ala Ala Ala
 225 230 235 240
 Ala Val Leu Gly Ala Val Arg Lys Arg Pro Ser Val Val Pro Met Ala
 245 250 255
 Gly Gln Asp Pro Ala Leu Ser Thr Ser His Pro Phe Tyr Asp Val Ala
 260 265 270
 Arg His Gly Ile Leu Gln Val Ala Gly Asp Asp Arg Phe Gly Arg Arg
 275 280 285
 Val Val Thr Phe Ser Cys Cys Arg Met Pro Pro Ser His Glu Leu Asp
 290 295 300
 His Gln Arg Leu Leu Glu Tyr Leu Lys Tyr Thr Leu Asp Gln Tyr Val
 305 310 315 320
 Glu Asn Asp Tyr Thr Ile Val Tyr Phe His Tyr Gly Leu Asn Ser Arg
 325 330 335
 Asn Lys Pro Ser Leu Gly Trp Leu Gln Ser Ala Tyr Lys Glu Phe Asp
 340 345 350
 Arg Lys Asp Gly Asp Leu Thr Met Trp Pro Arg Leu Val Ser Asn Ser
 355 360 365

Lys Leu Lys Arg Ser Ser His Leu Ser Leu Pro Lys Tyr Trp Asp Tyr
 370 375 380
 Arg Tyr Lys Lys Asn Leu Lys Ala Leu Tyr Val Val His Pro Thr Ser
 385 390 395
 Phe Ile Lys Val Leu Trp Asn Ile Leu Lys Pro Leu Ile Ser His Lys
 405 410 415
 Phe Gly Lys Lys Val Ile Tyr Phe Asn Tyr Leu Ser Glu Leu His Glu
 420 425 430
 His Leu Lys Tyr Asp Gln Leu Val Ile Pro Pro Glu Val Leu Arg Tyr
 435 440 445
 Asp Glu Lys Leu Gln Ser Leu His Glu Gly Arg Thr Pro Pro Pro Thr
 450 455 460
 Lys Thr Pro Pro Pro Arg Pro Pro Leu Pro Thr Gln Gln Phe Gly Val
 465 470 475
 Ser Leu Gln Tyr Leu Lys Asp Lys Asn Gln Gly Glu Leu Ile Pro Pro
 485 490 495
 Val Leu Arg Phe Thr Val Thr Tyr Leu Arg Glu Lys Gly Leu Arg Thr
 500 505 510
 Glu Gly Leu Phe Arg Arg Ser Ala Ser Val Gln Thr Val Arg Glu Ile
 515 520 525
 Gln Arg Leu Tyr Asn Gln Gly Lys Pro Val Asn Phe Asp Asp Tyr Gly
 530 535 540
 Asp Ile His Ile Pro Ala Val Ile Leu Lys Thr Phe Leu Arg Glu Leu
 545 550 555
 Pro Gln Pro Leu Leu Thr Phe Gln Ala Tyr Glu Gln Ile Leu Gly Ile
 565 570 575
 Thr Cys Val Glu Ser Ser Leu Arg Val Thr Gly Cys Arg Gln Ile Leu
 580 585 590
 Arg Ser Leu Pro Glu His Asn Tyr Val Val Leu Arg Tyr Leu Met Gly
 595 600 605
 Phe Leu His Ala Val Ser Arg Glu Ser Ile Phe Asn Lys Met Asn Ser
 610 615 620
 Ser Asn Leu Ala Cys Val Phe Gly Leu Asn Leu Ile Trp Pro Ser Gln
 625 630 635
 Gly Val Ser Ser Leu Ser Ala Leu Val Pro Leu Asn Met Phe Thr Glu
 645 650 655
 Leu Leu Ile Glu Tyr Tyr Glu Lys Ile Phe Ser Thr Pro Glu Ala Pro
 660 665 670
 Gly Glu His Gly Leu Ala Pro Trp Glu Gln Gly Ser Arg Ala Ala Pro
 675 680 685
 Leu Gln Glu Ala Val Pro Arg Thr Gln Ala Thr Gly Leu Thr Lys Pro
 690 695 700
 Thr Leu Pro Pro Ser Pro Leu Met Ala Ala Arg Arg Arg Leu Xaa Cys
 705 710 715
 Cys Glu His Ser Val Tyr Phe Glu Leu Pro Pro Thr Pro Val Cys Ala
 725 730 735
 Leu Val Cys Phe Val Asn Leu Ala Ser Val Lys Ile Thr Ser His
 740 745 750

<210> 19

<211> 718

<212> PRT

<213> Homo sapiens

<400> 19

Met Arg Thr Leu Arg Arg Leu Lys Phe Met Ser Ser Pro Ser Leu Ser
 1 5 10 15

Asp Leu Gly Lys Arg Glu Pro Ala Ala Ala Asp Glu Arg Gly Thr
 20 25 30
 Gln Gln Arg Arg Ala Cys Ala Asn Ala Thr Trp Asn Ser Ile His Asn
 35 40 45
 Gly Val Ile Ala Val Phe Gln Arg Lys Gly Leu Pro Asp Gln Glu Leu
 50 55 60
 Phe Ser Leu Asn Glu Gly Val Arg Gln Leu Leu Lys Thr Glu Leu Gly
 65 70 75 80
 Ser Phe Phe Thr Glu Tyr Leu Gln Asn Gln Leu Leu Thr Lys Gly Met
 85 90 95
 Val Ile Leu Arg Asp Lys Ile Arg Phe Tyr Glu Gly Gln Lys Leu Leu
 100 105 110
 Asp Ser Leu Ala Glu Thr Trp Asp Phe Phe Ser Asp Val Leu Pro
 115 120 125
 Met Leu Gln Ala Ile Phe Tyr Pro Val Gln Gly Lys Glu Pro Ser Val
 130 135 140
 Arg Gln Leu Ala Leu Leu His Phe Arg Asn Ala Ile Thr Leu Ser Val
 145 150 155 160
 Lys Leu Glu Asp Ala Leu Ala Arg Ala His Ala Arg Val Pro Pro Ala
 165 170 175
 Ile Val Gln Met Leu Leu Val Leu Gln Gly Val His Glu Ser Arg Gly
 180 185 190
 Val Thr Glu Asp Tyr Leu Arg Leu Glu Thr Leu Val Gln Lys Val Val
 195 200 205
 Ser Pro Tyr Leu Gly Thr Tyr Gly Leu His Ser Ser Glu Gly Pro Phe
 210 215 220
 Thr His Ser Cys Ile Leu Glu Leu Gln Arg Asp Lys Ala Ala Ala Ala
 225 230 235 240
 Ala Val Leu Gly Ala Val Arg Lys Arg Pro Ser Val Val Pro Met Ala
 245 250 255
 Gly Gln Asp Pro Ala Leu Ser Thr Ser His Pro Phe Tyr Asp Val Ala
 260 265 270
 Arg His Gly Ile Leu Gln Val Ala Gly Asp Asp Arg Phe Gly Arg Arg
 275 280 285
 Val Val Thr Phe Ser Cys Cys Arg Met Pro Pro Ser His Glu Leu Asp
 290 295 300
 His Gln Arg Leu Leu Glu Tyr Leu Lys Tyr Thr Leu Asp Gln Tyr Val
 305 310 315 320
 Glu Asn Asp Tyr Thr Ile Val Tyr Phe His Tyr Gly Leu Asn Ser Arg
 325 330 335
 Asn Lys Pro Ser Leu Gly Trp Leu Gln Ser Ala Tyr Lys Glu Phe Asp
 340 345 350
 Arg Lys Asp Gly Asp Leu Thr Met Trp Pro Arg Leu Val Ser Asn Ser
 355 360 365
 Lys Leu Lys Arg Ser Ser His Leu Ser Leu Pro Lys Tyr Trp Asp Tyr
 370 375 380
 Arg Tyr Lys Lys Asn Leu Lys Ala Leu Tyr Val Val His Pro Thr Ser
 385 390 395 400
 Phe Ile Lys Val Leu Trp Asn Ile Leu Lys Pro Leu Ile Ser His Lys
 405 410 415
 Phe Gly Lys Lys Val Ile Tyr Phe Asn Tyr Leu Ser Glu Leu His Glu
 420 425 430
 His Leu Lys Tyr Asp Gln Leu Val Ile Pro Pro Glu Val Leu Arg Tyr
 435 440 445
 Asp Glu Lys Leu Gln Ser Leu His Glu Gly Arg Thr Pro Pro Thr
 450 455 460
 Lys Thr Pro Pro Pro Arg Pro Pro Leu Pro Thr Gln Gln Phe Gly Val
 465 470 475 480

Ser Leu Gln Tyr Leu Lys Asp Lys Asn Gln Gly Glu Leu Ile Pro Pro
 485 490
 Val Leu Arg Phe Thr Val Thr Tyr Leu Arg Glu Lys Gly Leu Arg Thr
 500 505 510
 Glu Gly Leu Phe Arg Arg Ser Ala Ser Val Gln Thr Val Arg Glu Ile
 515 520 525
 Gln Arg Leu Tyr Asn Gln Gly Lys Pro Val Asn Phe Asp Asp Tyr Gly
 530 535 540
 Asp Ile His Ile Pro Ala Val Ile Leu Lys Thr Phe Leu Arg Glu Leu
 545 550 555 560
 Pro Gln Pro Leu Leu Thr Phe Gln Ala Tyr Glu Gln Ile Leu Gly Ile
 565 570 575
 Thr Cys Val Glu Ser Ser Leu Arg Val Thr Thr Gly Cys Arg Gln Ile Leu
 580 585 590
 Arg Ser Leu Pro Glu His Asn Tyr Val Val Leu Arg Tyr Leu Met Gly
 595 600 605
 Phe Leu His Ala Val Ser Arg Glu Ser Ile Phe Asn Lys Met Asn Ser
 610 615 620
 Ser Asn Leu Ala Cys Val Phe Gly Leu Asn Leu Ile Trp Pro Ser Gln
 625 630 635 640
 Gly Val Ser Ser Leu Ser Ala Leu Val Pro Leu Asn Met Phe Thr Glu
 645 650 655
 Leu Leu Ile Glu Tyr Tyr Glu Lys Ile Phe Ser Thr Pro Glu Ala Pro
 660 665 670
 Gly Glu His Gly Leu Ala Pro Trp Glu Gln Gly Ser Arg Ala Ala Pro
 675 680 685
 Leu Gln Glu Ala Val Pro Arg Thr Gln Ala Thr Gly Leu Thr Lys Pro
 690 695 700
 Thr Leu Pro Pro Ser Pro Leu Met Ala Ala Arg Arg Arg Leu
 705 710 715

<210> 20

<211> 1431

<212> DNA

<213> Homo sapiens

<400> 20

acgtcgcggg aggggcccagg tgagcggcag acccggcagc caggtggggg ccggcgggggt 60
 ccgtggccag agctgcagag agacaaggcg gccggcgctg ctgtgtctgg tgcagtgaag 120
 aagaggccct cgggtggtgcc catggctggc caggatccct cgtcagcac gagtccaccg 180
 ttctacgacg tggccagaca tggcattctg caggtggcag gggatgaccg ctttggaaag 240
 cgtgtgtgca cgttcagctg ctgcgggatg ccacctccc acgagctgga ccaccagcgg 300
 ctgctggaca ggtacaagaa gaacttgaag gccctctacg tggcgacccc caccagcttc 360
 atcaaggctc tgtggaacat cttgaagccc ctcatcagtc acaagtttgg gaagaaagtc 420
 atctatttca actactgag tgagctccac gaacacctta aatcagacca gctggtcatc 480
 cctcccgag ttttgcggta cgtagagaag ctccagagcc tgcacgaggg ccggacgccc 540
 cctcccacca agacaccacc gccgcggccc ccgctgccca cacagcagtt tggcgtcagt 600
 ctgcaatacc tcaagacaaa aaatcaaggc gaactcatcc cccctgtgct gaggttcaca 660
 gtgacgtacc tgagagagaa aggcctgcgc accgagggcc ttctccggag atccgccagc 720
 gtgcagacog tccgcgagat ccagaggctc tacaaccaag ggaagccctg gaactttgac 780
 gactacgggg acattcacat ccttgccgtg atcctgaaga ccttccctgc agagctgccg 840
 cagccgcttc ttacccttcca ggccacagag cagattctcg ggaacacctg tgtggagagc 900
 agcctgcgtg tcaactggctg ccgccagatc ttacggagcc tccagagaca caactacgtc 960
 gtctcccgct acctcatggg ctctcctgat cgggtgtccc gggagagcat cttcaacaaa 1020
 atgaacagct ctaacctggc ctgtgtcttc gggctgaatt tgactctgcc atcccagggg 1080
 gtctctcccg ttgctgcctt tgtgcctcgt aacatgttca ctgaactgct gatcgagtac 1140
 tatgaaaaga tcttcagcac ccggaggaca cctggggagc acggcctggc accatgggaa 1200

```

caggggagca gggcagcccc ttgtcaggag gctgtgccac ggacacaagc caccggcctc 1260
accaagccta cctacctcc gagtcccctg atggcagcca gaagacgtct ctagtgttgc 1320
gaacactctg tatatttcga gctacctccc acacctgtct gtgcacttgt atgttttgta 1380
aacttggcat ctgtaaaaat aaccagccat tagatgaatt cagaaccttc t      1431

```

<210> 21

<211> 390

<212> FRT

<213> Homo sapiens

<400> 21

```

Met Ala Gly Gln Asp Pro Ala Leu Ser Thr Ser His Pro Phe Tyr Asp
 1          5          10          15
Val Ala Arg His Gly Ile Leu Gln Val Ala Gly Asp Asp Arg Phe Gly
 20          25          30
Arg Arg Val Val Thr Phe Ser Cys Cys Arg Met Pro Pro Ser His Glu
 35          40          45
Leu Asp His Gln Arg Leu Leu Asp Arg Tyr Lys Lys Asn Leu Lys Ala
 50          55          60
Leu Tyr Val Val His Pro Thr Ser Phe Ile Lys Val Leu Trp Asn Ile
 65          70          75
Leu Lys Pro Leu Ile Ser His Lys Phe Gly Lys Lys Val Ile Tyr Phe
 85          90          95
Asn Tyr Leu Ser Glu Leu His Glu His Leu Lys Tyr Asp Gln Leu Val
100          105          110
Ile Pro Pro Glu Val Leu Arg Tyr Asp Glu Lys Leu Gln Ser Leu His
115          120          125
Glu Gly Arg Thr Pro Pro Pro Thr Lys Thr Pro Pro Arg Pro Pro
130          135          140
Leu Pro Thr Gln Gln Phe Gly Val Ser Leu Gln Tyr Leu Lys Asp Lys
145          150          155
Asn Gln Gly Glu Leu Ile Pro Pro Val Leu Arg Phe Thr Val Thr Tyr
165          170          175
Leu Arg Glu Lys Gly Leu Arg Thr Glu Gly Leu Phe Arg Arg Ser Ala
180          185          190
Ser Val Gln Thr Val Arg Glu Ile Gln Arg Leu Tyr Asn Gln Gly Lys
195          200          205
Pro Val Asn Phe Asp Asp Tyr Gly Asp Ile His Ile Pro Ala Val Ile
210          215          220
Leu Lys Thr Phe Leu Arg Glu Leu Pro Gln Pro Leu Leu Thr Phe Gln
225          230          235
Ala Tyr Glu Gln Ile Leu Gly Ile Thr Cys Val Glu Ser Ser Leu Arg
245          250          255
Val Thr Gly Cys Arg Gln Ile Leu Arg Ser Leu Pro Glu His Asn Tyr
260          265          270
Val Val Leu Arg Tyr Leu Met Gly Phe Leu His Ala Val Ser Arg Glu
275          280          285
Ser Ile Phe Asn Lys Met Asn Ser Ser Asn Leu Ala Cys Val Phe Gly
290          295          300
Leu Asn Leu Ile Trp Pro Ser Gln Gly Val Ser Ser Leu Ser Ala Leu
305          310          315
Val Pro Leu Asn Met Phe Thr Glu Leu Leu Ile Glu Tyr Tyr Glu Lys
325          330          335
Ile Phe Ser Thr Pro Glu Ala Pro Gly Glu His Gly Leu Ala Pro Trp
340          345          350
Glu Gln Gly Ser Arg Ala Ala Pro Leu Gln Glu Ala Val Pro Arg Thr
355          360          365
Gln Ala Thr Gly Leu Thr Lys Pro Thr Leu Pro Pro Ser Pro Leu Met

```

370
Ala Ala Arg Arg Arg Leu
385 390

375

380

<210> 22
<211> 2019
<212> DNA
<213> Homo sapiens

<400> 22
atgaggactc tccgcagggt gaagtctatg agttcgccca gcctcagtg cctggggcaag 60
agagagccgg ccgcccggcg ggaacgagogg ggaacgagcg agcgccgggg ctcgcgcaac 120
gccacctgga acagcatcca caacgggggtg atcgccgtct tccagcgcaa ggggctgccc 180
gaccaggagc tcttcagcct caacgagggc gtccggcagc tggtgaagac agagctgggg 240
tcctttctca cggagtacct gcagaacca ggcacgacaa aaggcatggt gatcctctgg 300
gacaagattc gcttctatga gggacagaag ctgctggact cactggcaga gacctgggac 360
ttctttctca gtgacgtgct gcccatgctg caggccatct tctaccgggt gcagggcgaag 420
gagccatcgg tgcgcacgct ggcctgctg cacttcogga atgccatcac cctcagtggt 480
aagctagagg atgcgctggc ccgggcccct gcccgtgtgc cccctgccat cgtgcagatg 540
ctgctggtgc tgcagggggt acatgagtc agggcgctga ctgaggacta cctggcgcctg 600
gagacgctgg tccagaaggt ggtgtgcgca tacctgggca cctacggcct ccactccagc 660
gagggggccc tcacccattc ctgcatcctg gagctgcaga gagacaaggc ggcggcggtg 720
gtgtggtgtg gtgcagtgag gaagaggccc togggtgtgc ccatggctgg ccaggatcct 780
cgctcgagca cgaatcaccc gttctacgac gtggccagac atggcattct gcaggtggca 840
ggggatgaac gctttggaag acgtgtgtgc acgttcagct gctgcgggat gccaccctcc 900
cacgagctgg accaccagcg gctgctggag tatltgaagt acacactgga ccaatcagct 960
gagaacgatt ataccatgct ctatltccac tacgggtgta acagccggaa caagccttcc 1020
ctgggctggc tccagagcgc atacaaggag ttcgatagga aagacgggga tctcactatg 1080
tgggccaggc tgggtctgaa ctccaagctc aagcatctct cccacctcag cctccccaag 1140
tactgggatt acaggtacaa gaagaacttg aaggccctct acgtgtgca ccccaccagc 1200
ttcatcaagg tctctgtgaa catcttgaag cccctcatca gtccacaagt tgggaagaaa 1260
gtcatctatt tcaactacct gactgagctc cacgaacacc ttaatacga ccagctggtc 1320
atccctcccg aagttttgag gtacgatgag aagctccaga gctgcacga gggccggacg 1380
ccgctcccca ccaagacacc acgcgcggcg ccccgctgca ccacacagca gtttggcgtc 1440
actctgcaat acctcaaga caaaaatcaa ggccaactca tccccctgtg gctgaggttc 1500
acagtgacgt acctgagaga gaaaggccct ccaagacaca actacgtcgt cctccgctac 1560
ctcatgggct tctgtcatgc ggtgtccogg gagagcatct tcaacaaaaa gaacagctct 1620
aacctggcct gtgtctctgg gctgaatttg atctggccat cccaggggtg ctctccctg 1680
agtgccttg tggccctgaa catgttccat gaactgtgta togagtacta tgaaaagatc 1740
ttcagacacc cggaggcacc tggggagcac ggctggcac catgggaa ca ggggagcagg 1800
gcagcccctt tgcaggaggg ttgtccacgg acacaagcga cgggctcac aggcctacc 1860
ctaacctcga gtccctgat ggcagccaga agacgtctct agtgttgca acactctgta 1920
tatttcgagc taacctccac acctgtctgt gacttgtat gttttgtaaa cttggcatct 1980
gtaaaaataa ccagccatta gatgaattca gaacctct 2019

<210> 23
<211> 633
<212> PRT
<213> Homo sapiens

<400> 23
Met Arg Thr Leu Arg Arg Leu Lys Phe Met Ser Ser Pro Ser Leu Ser
1 5 10 15
Asp Leu Gly Lys Arg Glu Pro Ala Ala Ala Asp Glu Arg Gly Thr
20 25 30
Gln Gln Arg Arg Ala Cys Ala Asn Ala Thr Trp Asn Ser Ile His Asn
35 40 45

Gly Val Ile Ala Val Phe Gln Arg Lys Gly Leu Pro Asp Gln Glu Leu
 50 55 60
 Phe Ser Leu Asn Glu Gly Val Arg Gln Leu Leu Lys Thr Glu Leu Gly
 65 70 75 80
 Ser Phe Phe Thr Glu Tyr Leu Gln Asn Gln Leu Leu Thr Lys Gly Met
 85 90 95
 Val Ile Leu Arg Asp Lys Ile Arg Phe Tyr Glu Gly Gln Lys Leu Leu
 100 105 110
 Asp Ser Leu Ala Glu Thr Trp Asp Phe Phe Phe Ser Asp Val Leu Pro
 115 120 125
 Met Leu Gln Ala Ile Phe Tyr Pro Val Gln Gly Lys Glu Pro Ser Val
 130 135 140
 Arg Gln Leu Ala Leu Leu His Phe Arg Asn Ala Ile Thr Leu Ser Val
 145 150 155 160
 Lys Leu Glu Asp Ala Leu Ala Arg Ala His Ala Arg Val Pro Pro Ala
 165 170 175
 Ile Val Gln Met Leu Leu Val Leu Gln Gly Val His Glu Ser Arg Gly
 180 185 190
 Val Thr Glu Asp Tyr Leu Arg Leu Glu Thr Leu Val Gln Lys Val Val
 195 200 205
 Ser Pro Tyr Leu Gly Thr Tyr Gly Leu His Ser Ser Glu Gly Pro Phe
 210 215 220
 Thr His Ser Cys Ile Leu Glu Leu Gln Arg Asp Lys Ala Ala Ala Ala
 225 230 235 240
 Ala Val Leu Gly Ala Val Arg Lys Arg Pro Ser Val Val Pro Met Ala
 245 250 255
 Gly Gln Asp Pro Ala Leu Ser Thr Ser His Pro Phe Tyr Asp Val Ala
 260 265 270
 Arg His Gly Ile Leu Gln Val Ala Gly Asp Asp Arg Phe Gly Arg Arg
 275 280 285
 Val Val Thr Phe Ser Cys Cys Arg Met Pro Pro Ser His Glu Leu Asp
 290 295 300
 His Gln Arg Leu Leu Glu Tyr Leu Lys Tyr Thr Leu Asp Gln Tyr Val
 305 310 315 320
 Glu Asn Asp Tyr Thr Ile Val Tyr Phe His Tyr Gly Leu Asn Ser Arg
 325 330 335
 Asn Lys Pro Ser Leu Gly Trp Leu Gln Ser Ala Tyr Lys Glu Phe Asp
 340 345 350
 Arg Lys Asp Gly Asp Leu Thr Met Trp Pro Arg Leu Val Ser Asn Ser
 355 360 365
 Lys Leu Lys Arg Ser Ser His Leu Ser Leu Pro Lys Tyr Trp Asp Tyr
 370 375 380
 Arg Tyr Lys Lys Asn Leu Lys Ala Leu Tyr Val Val His Pro Thr Ser
 385 390 395 400
 Phe Ile Lys Val Leu Trp Asn Ile Leu Lys Pro Leu Ile Ser His Lys
 405 410 415
 Phe Gly Lys Lys Val Ile Tyr Phe Asn Tyr Leu Ser Glu Leu His Glu
 420 425 430
 His Leu Lys Tyr Asp Gln Leu Val Ile Pro Pro Glu Val Leu Arg Tyr
 435 440 445
 Asp Glu Lys Leu Gln Ser Leu His Glu Gly Arg Thr Pro Pro Pro Thr
 450 455 460
 Lys Thr Pro Pro Pro Arg Pro Pro Leu Pro Thr Gln Gln Phe Gly Val
 465 470 475 480
 Ser Leu Gln Tyr Leu Lys Asp Lys Asn Gln Gly Glu Leu Ile Pro Pro
 485 490 495
 Val Leu Arg Phe Thr Val Thr Tyr Leu Arg Glu Lys Gly Leu Pro Glu
 500 505 510

27

His Asn Tyr Val Val Leu Arg Tyr Leu Met Gly Phe Leu His Ala Val
 515 520 525
 Ser Arg Glu Ser Ile Phe Asn Lys Met Asn Ser Ser Asn Leu Ala Cys
 530 535 540
 Val Phe Gly Leu Asn Leu Ile Trp Pro Ser Gln Gly Val Ser Ser Leu
 545 550 555 560
 Ser Ala Leu Val Pro Leu Asn Met Phe Thr Glu Leu Leu Ile Glu Tyr
 565 570 575
 Tyr Glu Lys Ile Phe Ser Thr Pro Glu Ala Pro Gly Glu His Gly Leu
 580 585 590
 Ala Pro Trp Glu Gln Gly Ser Arg Ala Ala Pro Leu Gln Glu Ala Val
 595 600 605
 Pro Arg Thr Gln Ala Thr Gly Leu Thr Lys Pro Thr Leu Pro Pro Ser
 610 615 620
 Pro Leu Met Ala Ala Arg Arg Arg Leu
 625 630

<210> 24

<211> 1398

<212> DNA

<213> Homo sapiens

<400> 24

acgtccgggg agggggccagg tgagcggcag acccggcag caggtggggg ccggcggggg 60
 cegtggccag agctgcagag agacaaggcg gcggcgccgtg ctgtgctggg tgcagtggg 120
 aagaggccct cgggtgtgcc catggctggc caggatcctg cgtgagcac gagtcaaccc 180
 ttctacgaag tggccagaca tggcattctg caggtggcag gggatgacgc ctttgaaga 240
 cgtgttgta cgttcagctg ctgcgggatg ccaccctccc acgagctgga ccaccagcgg 300
 ctgctggagt atttgaagta cacactggac caataccttg agaacgatta taccatcgtc 360
 tatttcgact acgggctgaa cagccgggaa aagccttccc tgggctggct ccagagcgca 420
 tacaaggagt tgcgataggaa agacggggat ctcaactatg ggcccaggct ggtctcgaa 480
 tccaagctca agcgatcctc ccacctcagc ctcccaagt actgggatta caggtacaag 540
 aagaacttga aggcctccta cgtgtgcac ccaccagct tcatcaaggt cctgtggaac 600
 atcttgaaag cctcatcag tcacaagttt gggaagaaag tcatctattt caactaactg 660
 agtgagctcc acgaacacct taatacagac cagctgtgta tccctcccgag agttttcgg 720
 tacgatgaga agctccagag cctgcacgag ggccggacgc cgctccac caagacacca 780
 ccgcgcggcg ccccgctgcc cacacagcag tttggcgtca gtctgcaata cctcaaaagac 840
 aaaaaatcaa gcgaactcat ccccccgtg ctgaggttca cagtgaagta cctgagagag 900
 aaaggcctcc cagagcacaa ctacgtgtgc ctccgtacc tcatgggctt cctgcatgag 960
 tgtgtccggg agagactctt caacaaaatg aacagctcta acctggcctg tgtcttcggg 1020
 ctgaatttga tctggccatc ccagggggtc tcttccctga gtgccttgg gccctgaac 1080
 atgttcactg aactgctgat cgagtactat gaaaagatct tcagcaccgc ggaggcaact 1140
 ggggagcagc gctggcaccc atgggaacag gggagcaggg cagcccttt gcaggaggtt 1200
 gtgccacgga cacaagccac gggcctcacc aagcctaccc tacctcagat tccctgatg 1260
 cagccagaaa gactgtctta gtgttgogaa cactctgat atttgcagc aotccocaca 1320
 cctgtctgtg caattgtatg ttttgaatac ttggcatctg taaaaataac cagccattag 1380
 atgaattcag aaccttct 1398

<210> 25

<211> 379

<212> PRT

<213> Homo sapiens

<400> 25

Met Ala Gly Gln Asp Pro Ala Leu Ser Thr Ser His Pro Phe Tyr Asp
 1 5 10 15
 Val Ala Arg His Gly Ile Leu Gln Val Ala Gly Asp Asp Arg Phe Gly


```

gacaagattc gcttctatga gggacagaag ctgctggact cactggcaga gacctgggac 360
ttcttcttca gtgacgtgct gccatgctg caggccatct tctaccgggt gcaggggcaag 420
gagccatcgg tgcgccagct ggcctgctg cacttccgga atgcatcac cctcagtggt 480
aagctagagg atcgctggc ccgggoccat gccctgtgc cccctgccat ctgcagatg 540
ctgctgggtg tgcagggggt acatgagtc aggggcgtga ctgaggacta cctgcgcctg 600
gagacgctgg tcacaagagt ggtgtgcga tacttgggca cctacggcct caactccagc 660
gaggggcccct tcaccattc ctgcatcct gagctgcaga gagacaaggc gcggcgagggt 720
cgtgtgctgg gtgcagtgag gaagaggccc tcggtggtgc coactggctgg ccaggatcct 780
gcgtgagca cgagtcaccc gttctacgac gtggccagac atggcattct gcaggtggca 840
gggatgacc gctttggaag acgtgttgtc acgttcagct gctgccgat gccaccctcc 900
cacgagctgg accaccagcg cgtctggag tactaagaaga acttgaaggc cctctacgtg 960
gtgcacccca ccagcttcat caaggtcctg tgaacatct tgaagccct catcagtcac 1020
aagtttggga aaaaagtcat ctatttcaac tacttgagtg agotccacga acactctaaa 1080
tacgaccagg tggatcatcc tccgaagt ttgggtacg atgagaagct ccagagacctg 1140
cacagggggc ggacgcccgc tcccaacca agaccaccgc ccggcgcccc gctgccaca 1200
cagcagtttg gctgcagct gcaatacctc aaagacaaaa atcaaggoga actcatcccc 1260
cctgtgctga ggttcacagt gacgtacctg agagagaaa cctccagag cacaactacg 1320
tcgtcctccg ctacctcatg ggtctctctc atcggtgtgc ccgggagagc atctcaaca 1380
aaatgaacag ctctaacctg ccctgtgtct tcgggctgaa ttgatctgg ccacccagg 1440
gggtctcttc cctgagtgcc ctgtgcccc tgaacatgt cactgaactg ctgactcgag 1500
actatgaaaa gatcttcagc accccggagg cactggggg gcacggcctg gcaccatggg 1560
aacagggggc caggcgagcc cctttgcagg aggctgtgcc acggacacaa gccacggggc 1620
tcaccaagcc tacctacct ccgagtcctc tgatggcagc cagaagacgt cctcagtggt 1680
gcgaacactc tgtatatctc gagctacctc ccacacctgt ctgtgcact gtatgtttt 1740
taaacttggc atctgtaaaa ataaccagcc attagatgaa ttcagaa 1787

```

<210> 27

<211> 461

<212> PRT

<213> Homo sapiens

<400> 27

```

Met Arg Thr Leu Arg Arg Leu Lys Phe Met Ser Ser Pro Ser Leu Ser
1 5 10 15
Asp Leu Gly Lys Arg Glu Pro Ala Ala Ala Asp Glu Arg Gly Thr
20 25 30
Gln Gln Arg Arg Ala Cys Ala Asn Ala Thr Trp Asn Ser Ile His Asn
35 40 45
Gly Val Ile Ala Val Phe Gln Arg Lys Gly Leu Pro Asp Gln Glu Leu
50 55 60
Phe Ser Leu Asn Glu Gly Val Arg Gln Leu Leu Lys Thr Glu Leu Gly
65 70 75 80
Ser Phe Phe Thr Glu Tyr Leu Gln Asn Gln Leu Leu Thr Lys Gly Met
85 90 95
Val Ile Leu Arg Asp Lys Ile Arg Phe Tyr Glu Gly Gln Lys Leu Leu
100 105 110
Asp Ser Leu Ala Glu Thr Trp Asp Phe Phe Phe Ser Asp Val Leu Pro
115 120 125
Met Leu Gln Ala Ile Phe Tyr Pro Val Gln Gly Lys Glu Pro Ser Val
130 135 140
Arg Gln Leu Ala Leu Leu His Phe Arg Asn Ala Ile Thr Leu Ser Val
145 150 155 160
Lys Leu Glu Asp Ala Leu Ala Arg Ala His Ala Arg Val Pro Pro Ala
165 170 175
Ile Val Gln Met Leu Leu Val Leu Gln Gly Val His Glu Ser Arg Gly
180 185 190
Val Thr Glu Asp Tyr Leu Arg Leu Glu Thr Leu Val Gln Lys Val Val
195 200 205

```

30

Ser Pro Tyr Leu Gly Thr Tyr Gly Leu His Ser Ser Glu Gly Pro Phe
 210 215 220
 Thr His Ser Cys Ile Leu Glu Leu Gln Arg Asp Lys Ala Ala Ala Ala
 225 230 240
 Ala Val Leu Gly Ala Val Arg Lys Arg Pro Ser Val Val Pro Met Ala
 245 250 255
 Gly Gln Asp Pro Ala Leu Ser Thr Ser His Pro Phe Tyr Asp Val Ala
 260 265 270
 Arg His Gly Ile Leu Gln Val Ala Gly Asp Asp Arg Phe Gly Arg Arg
 275 280 285
 Val Val Thr Phe Ser Cys Cys Arg Met Pro Pro Ser His Glu Leu Asp
 290 295 300
 His Gln Arg Leu Leu Glu Tyr Lys Lys Asn Leu Lys Ala Leu Tyr Val
 305 310 315 320
 Val His Pro Thr Ser Phe Ile Lys Val Leu Trp Asn Ile Leu Lys Pro
 325 330 335
 Leu Ile Ser His Lys Phe Gly Lys Lys Val Ile Tyr Phe Asn Tyr Leu
 340 345 350
 Ser Glu Leu His Glu His Leu Lys Tyr Asp Gln Leu Val Ile Pro Pro
 355 360 365
 Glu Val Leu Arg Tyr Asp Glu Lys Leu Gln Ser Leu His Glu Gly Arg
 370 375 380
 Thr Pro Pro Pro Thr Lys Thr Pro Pro Pro Arg Pro Pro Leu Pro Thr
 385 390 395 400
 Gln Gln Phe Gly Val Ser Leu Gln Tyr Leu Lys Asp Lys Asn Gln Gly
 405 410 415
 Glu Leu Ile Pro Pro Val Leu Arg Phe Thr Val Thr Tyr Leu Arg Glu
 420 425 430
 Lys Ala Ser Gln Ser Thr Thr Thr Ser Ser Ser Ala Thr Ser Trp Ala
 435 440 445
 Ser Cys Met Arg Cys Pro Gly Arg Ala Ser Ser Thr Lys
 450 455 460

<210> 28
 <211> 1176
 <212> DNA
 <213> Homo sapiens

<400> 28
 acgtccgggg agggggccagg tgagcggcag acccggcacg caggtggggg ccggcggggg 60
 ccgtggccag agctgcagag agacaaggcg gggcgcggtg ctgtgctggg tgcagtgagg 120
 aagaggccct ccgtgggtgcc catggctggc caggatcctg cgctgagcac gagtccacccg 180
 ttctacgacg tggccagaca tggcattctg caggtggcag gggatgaccg ctttgggaaga 240
 cgtgttgtca cgttcagctg ctgcoggatg ccaccctccc acgagctgga ccaccagcgg 300
 ctgtcggaac ggtacaagaa gaacttgaag gccctctacg tggtgacccc caccagcttc 360
 atcaaggctc tgtggaacat cttgaagccc ctcatcagtc acaagtttgg gaagaagctc 420
 atctatttca actacctgag tgagctccac gaacacctta aatacgacca gctggtcatc 480
 cctccogaag ttttgcggtg cgtagagaag ctccagagcc tgcacgaggg ccggacgccg 540
 cctccacaca agacacaccc gccgcggccc ccgctgccca cacagcagtt tggcgtcagt 600
 ctgcaatacc tcaaaagacaa aaatcaaaggc gaactcatcc cccctgtgct gagggtcaca 660
 gtgacgtacc tgagagagaa aggcctccca gaggcacaact acgtcgctcc ccgctacctc 720
 atgggctctc tgcattcggt gtccoggggag agcatcttca acaaaatgaa cagctctaac 780
 ctggcctgtg tcttcgggct gaatttgatc tggcatccc aggggggtctc ctccctgagt 840
 gcccttgtgc cctgaaacat gtctcaactgaa ctgctgatcg agtactatga aaagatcttc 900
 gcccccggcg aggcacctgg ggagcaccgg ctggcaccat ggggaacagg gaggcaggga 960
 gcccttttgc aggaggtgtg gccacggaca caagccacgg gcttcaccaa gctaccctta 1020
 cctcogagtc ccttgatggc agccagaaga cgtctctagt gttgcgaaca ctctgtatat 1080

ttcgagctac ctccacacacc tgtctgtgca cttgtatgtt ttgtaaactt ggcattctgta 1140
 aaataacca gccattagat gaattcagaa ccttct 1176

<210> 29
 <211> 305
 <212> PRT
 <213> Homo sapiens

<400> 29
 Met Ala Gly Gln Asp Pro Ala Leu Ser Thr Ser His Pro Phe Tyr Asp
 1 5 10 15
 Val Ala Arg His Gly Ile Leu Gln Val Ala Gly Asp Asp Arg Phe Gly
 20 25 30
 Arg Arg Val Val Thr Phe Ser Cys Cys Arg Met Pro Ser His Glu
 35 40 45
 Leu Asp His Gln Arg Leu Leu Asp Arg Tyr Lys Lys Asn Leu Lys Ala
 50 55 60
 Leu Tyr Val Val His Pro Thr Ser Phe Ile Lys Val Leu Trp Asn Ile
 65 70 75 80
 Leu Lys Pro Leu Ile Ser His Lys Phe Gly Lys Lys Val Ile Tyr Phe
 85 90 95
 Asn Tyr Leu Ser Glu Leu His Glu His Leu Lys Tyr Asp Gln Leu Val
 100 105 110
 Ile Pro Pro Glu Val Leu Arg Tyr Asp Glu Lys Leu Gln Ser Leu His
 115 120 125
 Glu Gly Arg Thr Pro Pro Pro Thr Lys Thr Pro Pro Arg Pro Pro
 130 135 140
 Leu Pro Thr Gln Gln Phe Gly Val Ser Leu Gln Tyr Leu Lys Asp Lys
 145 150 155 160
 Asn Gln Gly Glu Leu Ile Pro Pro Val Leu Arg Phe Thr Val Thr Tyr
 165 170 175
 Leu Arg Glu Lys Gly Leu Pro Glu His Asn Tyr Val Val Leu Arg Tyr
 180 185 190
 Leu Met Gly Phe Leu His Ala Val Ser Arg Glu Ser Ile Phe Asn Lys
 195 200 205
 Met Asn Ser Ser Asn Leu Ala Cys Val Phe Gly Leu Asn Leu Ile Trp
 210 215 220
 Pro Ser Gln Gly Val Ser Ser Leu Ser Ala Leu Val Pro Leu Asn Met
 225 230 235 240
 Phe Thr Glu Leu Leu Ile Glu Tyr Tyr Glu Lys Ile Phe Ser Thr Pro
 245 250 255
 Glu Ala Pro Gly Glu His Gly Leu Ala Pro Trp Glu Gln Gly Ser Arg
 260 265 270
 Ala Ala Pro Leu Gln Glu Ala Val Pro Arg Thr Gln Ala Thr Gly Leu
 275 280 285
 Thr Lys Pro Thr Leu Pro Pro Ser Pro Leu Met Ala Ala Arg Arg Arg
 290 295 300
 Leu
 305

<210> 30
 <211> 3257
 <212> DNA
 <213> Homo sapiens

<400> 30
 atttccttct ccctttccccc gccagcttgc catccatctc ccccccacccg taacccctc 60

ctgcctccat	ccaccggggc	tatggccgca	gaagaggtat	tgcagacggt	ggaccattat	120
aagactgaga	tagagaggct	aaccaaggag	ctcacggaga	ccaccacaga	gaagatccag	180
gctgcocgag	acgggctggt	ggtgctggag	gagaagctga	ccctcaaaa	gcagtatgat	240
gaactcgagg	ctgagtacga	cagcctcaaa	caggagctgg	agcagctcaa	agggccattt	300
gggacgtcc	tctccatcca	cgggaaggtt	gctgaagatg	gagagactcg	ggaggaacag	360
cttctgcagg	atgcagcadc	gaaggaggct	tactatctgg	ggaagatctg	ggagatgcag	420
aacgagctga	aacagagccg	ggctgtggct	actaatgtac	aggcagaaaa	cgagaggctc	480
accgcagctg	tgcagagctc	gaaggagaac	aatgagatgg	tggagctaca	gagaatacgg	540
atgaaggatg	aaatccgaga	atataagttc	cgggaggcac	ggctccttca	ggactatact	600
gaattggaag	aagaaaaat	caatctgcag	aaactagtgt	ccacgttgaa	gcagaaaccg	660
gttgaatcac	aaggcttaaa	gcattgagatt	aagcgatttg	aggaggagac	ggtactgctg	720
aacagccagc	tggaaagtgc	catccgattg	aaagagattg	ctgagcacca	actggaagaa	780
gccctcgaga	ctttaaaaaa	tgaagagag	caaaagaaca	acctgcggaa	ggagctctcc	840
cagtatatca	gcctcaatga	taaccatata	agcatctcag	tagatggact	caaatctgac	900
gaggaatggga	gtgaaccaaa	caatgatgac	aaatgaacg	gtcatatcca	tgggctctct	960
gtgaaactga	gtggagacta	tccgactccc	acctaaagga	aaggagagtc	cttgaacctc	1020
gtctctgact	tattcagtga	gctgaacatt	tcagaaatac	agaagtggaa	gcagcagctt	1080
atgcaggtag	agcgggaaaa	ggccattctt	tggccaacc	tacaggagtc	acagacaacg	1140
ctggaacaca	ccaagggggc	actgacggag	cagcatgagc	gggtgacccg	gctccacagag	1200
cacgtcaagt	cagctgaggg	cctgcaaaag	caagaaggag	tcaaggctga	gtggaacggg	1260
gagaaggggc	gggactcagg	ggaggaggcc	catgatctatg	aggtggacat	caatgggtta	1320
gagatccttg	aattgcaaaa	caggggtggca	gtaactgagc	tgattgatct	gaagctgaa	1380
attaaggcct	taaaaggaaa	atataataaa	tctgtagaata	actacactga	tgagaaggcc	1440
aagtatgaga	gtataaatcca	gatgtatgat	gagcaggtga	caagccttga	gaagaccacc	1500
gacgaagagt	gtgagaagat	ggccacatgc	gagaaggagt	tgcataaagt	gaccagcata	1560
gccaaagaaa	atcacagctac	ccttaatacg	gccacggatg	agttagtgtac	attcagtgag	1620
gagttagctc	agctttacca	ccatgtgtgt	ctatgtataa	atgaaactcc	caacagggtc	1680
atgctgactt	actataggca	gagcagagtc	accgcagtg	gcagcctgaa	agggcccgat	1740
gatcccagag	gactttgttc	cccacgatta	gccaggcggt	gtgtgtctac	cccggtagaa	1800
acaaggacct	cctctgaacc	agttgcaaaa	gaaagcacag	agccccgcaa	agaaaccaagt	1860
ccaactaaga	cccaccaaat	ctctcctgtt	attactgccc	caccgtccac	tcagtlattg	1920
gatacaagtg	acatccgcaa	agagccaatg	aatatctaca	accttaatgc	cataatccgg	1980
gaccaaatca	gaaagctgtg	gaaagctgtg	gacggctcct	tgcactgtct	tctcaaaaga	2040
gcagcagctc	gggagctagc	ccccatgatt	gataaagaca	aggaagccct	aatggaagag	2100
atcctcaagc	taaagtcctc	gctgagcacc	aaacggggag	agatcgccac	attgaggggc	2160
gtgttgaaag	ccaacaagca	gacagctgag	gtggcgctag	ctaattccaa	gaacaatat	2220
gaaaatgaaa	aagcaatggg	gactgaaaac	atgaogaaag	ttagaaatga	actgaaggct	2280
ttgaaaagag	gtctgcacac	cttctcatcc	ctgagaaaca	tggttgcaac	agaagtgtgat	2340
gaatatgtca	cccagtttga	tgaatgcag	acagagttag	cagctgcaga	ggatgagaag	2400
aagactctga	acacttttgt	acgaatggct	atccagcaaa	aactcgccct	gcaccaagag	2460
ctggaggact	tagagtttga	ccatgagcag	tcccgcagca	gcaaaggcaa	acttggaaag	2520
agcaacatgc	cagcgcctaa	agtaagtggc	gaggcatacg	tcacgtgcag	ccccatagac	2580
acttacctcc	tgcatagtca	gggcccacag	accaccaaca	ttcggttcag	cagttggcact	2640
cagagggaaa	gacaattttc	accttccctt	tgatgacaga	gcgtccccag	gacttcaggg	2700
gcttctcacc	tacagaattt	atlaagagtt	ccccctgac	ccactccac	agaattcattt	2760
cttctgaagg	gcccctcttc	catgagttaa	ttcatcgaag	ggcaccggct	cagcaaggaa	2820
aaaagggttaa	cgtgtctccc	accagattgt	cagcagcctg	ctgcctccgt	accgccacag	2880
tgtctcaaac	tgcgcgggag	gcaagactgc	coactgcaca	gtctgtacac	agctctccct	2940
gaggagcagc	caacttccag	ctccagctgc	gcccctctcc	actgtctctc	caagctccct	3000
caccctcagt	cttcatctcc	tgtgtagcga	catctggggg	ggaggttttg	tgcgaacaca	3060
caggatactg	cccaagatcc	agcgggtgtt	ttcttctcgg	ttgttagatg	tacaatttga	3120
ttaatgtcca	tctgttttga	agacgagaaa	tgtgagaaga	acacagagaa	cagaccctga	3180
tgtgataaaa	cattttctgc	tttctctgag	tcacagataa	acttctgcga	tcgaattggc	3240
acagttcatt	taaaattt					3257

<210> 31

<211> 975

<212> PRT

<213> Homo sapiens

<400> 31

```

Met Ala Ala Glu Glu Val Leu Gln Thr Val Asp His Tyr Lys Thr Glu
 1      5      10      15
Ile Glu Arg Leu Thr Lys Glu Leu Thr Glu Thr Thr His Glu Lys Ile
 20      25      30
Gln Ala Ala Glu Tyr Gly Leu Val Val Leu Glu Glu Lys Leu Thr Leu
 35      40      45
Lys Gln Gln Tyr Asp Glu Leu Glu Ala Glu Tyr Asp Ser Leu Lys Gln
 50      55      60
Glu Leu Glu Gln Leu Lys Glu Ala Phe Gly Gln Ser Phe Ser Ile His
 65      70      75      80
Arg Lys Val Ala Glu Asp Gly Glu Thr Arg Glu Glu Thr Leu Leu Gln
 85      90      95
Glu Ser Ala Ser Lys Glu Ala Tyr Tyr Leu Gly Lys Ile Leu Glu Met
100      105      110
Gln Asn Glu Leu Lys Gln Ser Arg Ala Val Val Thr Asn Val Gln Ala
115      120      125
Glu Asn Glu Arg Leu Thr Ala Val Val Gln Asp Leu Lys Glu Asn Asn
130      135      140
Glu Met Val Glu Leu Gln Arg Ile Arg Met Lys Asp Glu Ile Arg Glu
145      150      155      160
Tyr Lys Phe Arg Glu Ala Arg Leu Leu Gln Asp Tyr Thr Glu Leu Glu
165      170      175
Glu Glu Asn Ile Thr Leu Gln Lys Leu Val Ser Thr Leu Lys Gln Asn
180      185      190
Gln Val Glu Tyr Glu Gly Leu Lys His Glu Ile Lys Arg Phe Glu Glu
195      200      205
Glu Thr Val Leu Leu Asn Ser Gln Leu Glu Asp Ala Ile Arg Leu Lys
210      215      220
Glu Ile Ala Glu His Gln Leu Glu Glu Ala Leu Glu Thr Leu Lys Asn
225      230      235      240
Glu Arg Glu Gln Lys Asn Asn Leu Arg Lys Glu Leu Ser Gln Tyr Ile
245      250      255
Ser Leu Asn Asp Asn His Ile Ser Ile Ser Val Asp Gly Leu Lys Phe
260      265      270
Ala Glu Asp Gly Ser Glu Pro Asn Asn Asp Asp Lys Met Asn Gly His
275      280      285
Ile His Gly Pro Leu Val Lys Leu Asn Gly Asp Tyr Arg Thr Pro Thr
290      295      300
Leu Arg Lys Gly Glu Ser Leu Asn Pro Val Ser Asp Leu Phe Ser Glu
305      310      315      320
Leu Asn Ile Ser Glu Ile Gln Lys Leu Lys Gln Gln Leu Met Gln Val
325      330      335
Glu Arg Glu Lys Ala Ile Leu Leu Ala Asn Leu Gln Glu Ser Gln Thr
340      345      350
Gln Leu Glu His Thr Lys Gly Ala Leu Thr Glu Gln His Glu Arg Val
355      360      365
His Arg Leu Thr Glu His Val Asn Ala Met Arg Gly Leu Gln Ser Ser
370      375      380
Lys Glu Leu Lys Ala Glu Leu Asp Gly Glu Lys Gly Arg Asp Ser Gly
385      390      395      400
Glu Glu Ala His Asp Tyr Glu Val Asp Ile Asn Gly Leu Glu Ile Leu
405      410      415
Glu Cys Lys Tyr Arg Val Ala Val Thr Glu Val Ile Asp Leu Lys Ala
420      425      430
Glu Ile Lys Ala Leu Lys Glu Lys Tyr Asn Lys Ser Val Glu Asn Tyr

```

		435							440						445							
Thr	Asp	Glu	Lys	Ala	Lys	Tyr	Glu	Ser	Lys	Ile	Gln	Met	Tyr	Asp	Glu							
	450					455					460											
Gln	Val	Thr	Ser	Leu	Gly	Lys	Thr	Thr	Lys	Glu	Ser	Gly	Glu	Lys	Met	480						
	465				470					475												
Ala	His	Met	Glu	Lys	Glu	Leu	Gln	Lys	Met	Thr	Ser	Ile	Ala	Asn	Glu	495						
				485					490													
Asn	His	Ser	Thr	Leu	Asn	Thr	Ala	Gln	Asp	Glu	Leu	Val	Thr	Phe	Ser							
			500					505					510									
Glu	Glu	Leu	Ala	Gln	Leu	Tyr	His	His	Val	Cys	Leu	Cys	Asn	Asn	Gln							
	515						520					525										
Thr	Pro	Asn	Arg	Val	Met	Leu	Asp	Tyr	Tyr	Arg	Gln	Ser	Arg	Val	Thr							
	530					535					540											
Arg	Ser	Gly	Ser	Leu	Lys	Gly	Pro	Asp	Asp	Pro	Arg	Gly	Leu	Leu	Ser							
	545				550				555						560							
Pro	Arg	Leu	Ala	Arg	Arg	Gly	Val	Ser	Ser	Pro	Val	Glu	Thr	Arg	Thr							
				565					570					575								
Ser	Ser	Glu	Pro	Val	Ala	Lys	Glu	Ser	Thr	Glu	Pro	Ser	Lys	Glu	Pro							
			580					585					590									
Ser	Pro	Thr	Lys	Thr	Pro	Thr	Ile	Ser	Pro	Val	Ile	Thr	Ala	Pro	Pro							
	595					600					605											
Ser	Ser	Pro	Val	Leu	Asp	Thr	Ser	Asp	Ile	Arg	Lys	Glu	Pro	Met	Asn							
	610					615					620											
Ile	Tyr	Asn	Leu	Asn	Ala	Ile	Ile	Arg	Asp	Gln	Ile	Lys	His	Leu	Gln							
	625				630				635					640								
Lys	Ala	Val	Asp	Arg	Ser	Leu	Gln	Leu	Ser	Arg	Gln	Arg	Ala	Ala								
				645					650					655								
Arg	Glu	Leu	Ala	Pro	Met	Ile	Asp	Lys	Asp	Lys	Glu	Ala	Leu	Met	Glu							
			660					665					670									
Glu	Ile	Leu	Lys	Leu	Lys	Ser	Leu	Leu	Ser	Thr	Lys	Arg	Glu	Gln	Ile							
	675						680					685										
Ala	Thr	Leu	Arg	Ala	Val	Leu	Lys	Ala	Asn	Lys	Gln	Thr	Ala	Glu	Val							
	690					695					700											
Ala	Leu	Ala	Asn	Leu	Lys	Asn	Lys	Tyr	Glu	Asn	Glu	Lys	Ala	Met	Val							
	705				710				715					720								
Thr	Glu	Thr	Met	Thr	Lys	Leu	Arg	Asn	Glu	Leu	Lys	Ala	Leu	Lys	Glu							
			725						730					735								
Asp	Thr	Ala	Thr	Phe	Ser	Ser	Leu	Arg	Thr	Met	Phe	Ala	Thr	Arg	Cys							
			740					745					750									
Asp	Glu	Tyr	Val	Thr	Gln	Leu	Asp	Glu	Met	Gln	Arg	Gln	Leu	Ala	Ala							
	755						760					765										
Ala	Glu	Asp	Glu</																			

900										905					910				
Glu	Lys	Arg	Leu	Thr	Val	Ala	Pro	Asp	Cys	Gln	Gln	Pro	Ala	Ala					
915										920					925				
Ser	Val	Pro	Pro	Gln	Cys	Ser	Gln	Leu	Ala	Gly	Arg	Gln	Asp	Cys	Pro				
930										935					940				
Thr	Val	Ser	Pro	Asp	Thr	Ala	Leu	Pro	Glu	Glu	Gln	Pro	His	Ser	Ser				
945										950					955				
Ser	Gln	Cys	Ala	Pro	Leu	His	Cys	Leu	Ser	Lys	Pro	Pro	His	Pro	Ser				
965										970					975				

```
<210> 32
<211> 2717
<212> DNA
<213> Homo sapiens
```

[illegible]

36

```

gottaatctt gtctctcttc totttctttat gatgttctact ttaaaaaacaa caaaaccctt 2520
gagctggact gttgagcagg cctgtctctc ctattaagta aaaataaata gtagtagtat 2580
gtttgtaagc tattctgaca gaaaagacaa aggtacttaa ttgtatgata gtgtttttat 2640
atggaagaat gtacagctta tggacaaatg tacacctttt tgttacttta ataaaaatgt 2700
agtaggataa aaaaaaa 2717

```

```

<210> 33
<211> 158
<212> PRT
<213> Homo sapiens

```

```

<400> 33
Met Ser His Gly Lys Gly Thr Asp Met Leu Pro Glu Ile Ala Ala Ala
1 5 10 15
Val Gly Phe Leu Ser Ser Leu Leu Arg Thr Arg Gly Cys Val Ser Glu
20 25 30
Gln Arg Leu Lys Val Phe Ser Gly Ala Leu Glu Ala Leu Thr Glu
35 40 45
His Tyr Lys His His Trp Phe Pro Glu Lys Pro Ser Lys Gly Ser Gly
50 55 60
Tyr Arg Cys Ile Arg Ile Asn His Lys Met Asp Pro Ile Ile Ser Arg
65 70 75 80
Val Ala Ser Gln Ile Gly Leu Ser Gln Pro Gln Leu His Gln Leu Leu
85 90 95
Pro Ser Glu Leu Thr Leu Trp Val Asp Pro Tyr Glu Val Ser Tyr Arg
100 105 110
Ile Gly Glu Asp Gly Ser Ile Cys Val Leu Tyr Glu Glu Ala Pro Leu
115 120 125
Ala Ala Ser Cys Gly Leu Leu Thr Cys Lys Asn Gln Val Leu Leu Gly
130 135 140
Arg Ser Ser Pro Ser Lys Asn Tyr Val Met Ala Val Ser Ser
145 150 155

```

```

<210> 34
<211> 5471
<212> DNA
<213> Homo sapiens

```

```

<400> 34
acagctgtgca acactcggca tcttttctgg aggcgccttc ttcagcagcc gcagatggca 60
tcgggtcgog ggcctggggc tcgcaattga ttctccccc tgcccacctc gactccacgg 120
acgcacotct cctctccctt cctcccttgc cgtctctggg tctgagccca gctcgcgacc 180
gcggggcaga gnatcagtcg cggcgccga ggctgagcag cagcgtcttc gctccctgac 240
gtggggagaa gcgcccaacc gggagagctg atcccggtct gcctccagcg ccccccacct 300
ttgcactcc aagccggggg ctcagagac ccgctcctcc aggcgcact atgtcggaac 360
cttgctccag cgaagaagaa tcggatgaga tcgtggagga ggagagcggc aaggaggtgc 420
tcgctcgggc ccgctcgggc gcgcgcctgt ctccagccg taccagcgag gctcggccg 480
gcagcgcggc gctggggggc ggcgcgcggc gcgcgggagc cggggtgggt gcagcggcg 540
ggggggcgag cggcgcgagg agcggcgggc gggcggggg gctgcaacc agcaccccg 600
ctggcgggcg ccggccctcc agccccagcc cgtcggtggg gaggagagag gagaaggaag 660
agttgagagc gctgcagaaa gaggagggag agaggaagaa gaggctgcag ctgtatgtgt 720
ctgtgatcgc ctgcctgccc taccctttta atgccaaaga gccacacgac agctcgcgc 780
ggcagcagaa gatcagcaaa cagcagctgc agacagtcaa ggaacgggtt caggctttcc 840
tcaatgggga aaccagatc atggctgagc aagccttcac gaaacgctgt cagagttact 900
atgagtggtt cctgaagagc gacggtgtgg ccgcgatggt tcagagtgga ggcgtgtccg 960
ccaacgactc ccgggaggtc ttcaagaagc acattgagaa gagagtgccg agcctgcctg 1020
agattgacgg cctcagcaag gagactgtgc tgagctcctg gatggccaaa ttgatgcc 1080

```

tctaccgttg	agaagaggac	cgcgggaagc	agcaggcccg	gatgacagcc	agcgcagcct	1140
ccgagctgat	tctgagcaag	gagcaactct	atgagatgtt	ccagaacatt	cttgggattc	1200
agaagcttga	acatcagctc	ctttacaatg	cctgccagct	ggacaatcca	gatggagcaag	1260
cagcccaagt	cagcagcagag	ctgtgatggc	gtctacaaat	ggcagaccac	atgagccaggc	1320
aacgcacaa	tcccaagtgt	gtatccaaag	aaatggaaaa	catgtacatt	gaggagctga	1380
agtcatctgt	caacctgtct	atggccaaact	tggagagcat	gccggtatcc	aaagccgggg	1440
agttcaagct	ccagaaaactc	aaacgcagcc	acaatgcttc	catcatcgac	atgggcgagg	1500
atagtgagaa	ccagctctccc	aagtcagatg	tcgtgtctgc	tttctcatgt	gaggttggtaa	1560
ttatggaa	ccaaagctctc	aaatcttttg	ctccaaatcg	catcgtatat	tgcaaatagg	1620
aggtggaagg	aggagagaaa	ctacagactg	atcaggccga	ggcttctaaa	ccaacctggg	1680
gcacccagg	tgactctccc	acaacccatg	cactgccagc	tgtgaaggtg	aagctgttca	1740
cagagagcac	aggcgtcctg	gcgttggagg	acaaggagct	tggggcggtt	attctccatc	1800
ccaccocgaa	cagccccaaa	cagtcagagt	ggcacaacat	gacagctccc	aaaaactggc	1860
ccgacaaaga	tctcaaaatc	aaacttgctg	tccgaatgga	taagcctcaa	aacatgaagc	1920
attctgggta	tttatgggcc	atcggtaaga	atgtctggaa	gagatgggaag	aaaagggttt	1980
ttgtatttgt	gcaggtcagt	cagtcacagt	ttgccatgtg	cagttatcgg	gagaagaaa	2040
cggagcctca	ggaactctca	caattggagt	gctacactgt	ggattacacc	gaccccagc	2100
caggttttga	gggtggccga	gccttcttca	atcgtgtcaa	ggagggagac	accgtgatct	2160
ttgccagtga	cgatgaacaa	gaccgcattc	tgtgggtcca	ggccatgtat	cggggccacg	2220
ggcagtcaca	caagctctcc	ccccgacccc	aagtcacgaa	actcaacggc	agagagtgaa	2280
atgtacctca	gctggatgcc	cttatctctc	aattttaccg	agatagagct	caaaaacatg	2340
gcattggatga	atttatctct	tccaaacctc	gtacatttga	ccacgcttcc	ctctttgaga	2400
tggtacaacg	ccctactttg	gatcacagac	ttaatgattc	ctattctctg	ctgggctggg	2460
tcagttctgg	ccaggtcttt	gtactagacg	agtaattgcg	ccgaaatgga	gtccggcggt	2520
gtcacccaga	ctctctgtac	ctcagagact	tgcttgaacg	ggcagaaaat	ggcgccatga	2580
tcgacccccc	ctctctcac	tacagctttg	ctctctgtgc	atcccatgtc	catggggaaca	2640
ggcctgatgg	aattggaact	gtgactgttg	aagaaaagga	acgttttgaa	gaaatcaaa	2700
agaggctccg	agttctgcta	gaaaatcaga	ttacacattt	taggtattgc	tttccattg	2760
gtcgacctga	aggtgctttg	aaagctactc	tctcactctt	ggaaaggttt	tttgtgaaag	2820
atatttgtac	cccagtgcca	caagaggagg	taaaaacagt	tatccgtaaa	tgtctggaac	2880
aggctgcgtt	agtcacactat	tctcggctct	cagagtatgc	caaaaatcgaa	gagaatccaa	2940
aggatgcaga	aaatgtaggc	cggtttaatca	ctcctgccaa	aaagcttgaa	gatcaaatac	3000
gtcttgctga	actagctatt	gaagttcttc	agcaaaaatga	ggagcaccac	ctagactggc	3060
atgttgtata	aggagaagcc	tttgcgtgtg	ggtcagattt	aatgggtggg	catgcgagaa	3120
cgttctctgc	actctttgca	gtagacatgg	atgcagcctt	agaggtgcac	cctccagaca	3180
catgggacag	tttttcaact	tttcagctgc	tgaatgattt	tctccgtact	gactataatt	3240
tgtgcgaatg	aaaattttcac	aaacacctgc	aagacotggt	tgccccactt	gttgttagat	3300
atgtggatct	gatggagtcc	tcaattgcac	aatccattca	caggggcttt	gagcgggagt	3360
catgggaacc	agtcgaaggt	ttaaccagta	acctacccaa	tgtgaacctc	cccaatgtga	3420
accttcccaa	agttccaaat	ctaccagtta	acalccctta	aggcatccca	caaatgtcca	3480
ctttttcggc	accgtctagg	atggctgcta	tatatgatgc	ggataatggg	tcagccacct	3540
cagaagcttg	gtttttgaaa	cttgacgccc	ttcagacctt	cattegggac	ctgacactgc	3600
ctgaagaaga	gttttggaa	caactggaac	aacggctgaa	gttgatggca	agtgacatga	3660
tcgaattgtg	tgttcaaaa	accaggatg	catttgaa	taagctgcac	aaaaccagtc	3720
catcaacaga	tttttcagtc	ccacagtcaa	tatgacacat	gtttaatggt	atgggttgat	3780
ccaaagctca	atcaacaaaa	ctttgcagca	tggaaatggg	ccaagagcat	caataccatt	3840
caaaaataga	cgaactcaatt	gaagaaactg	ttaaagaaat	gataaacact	ttggttgcac	3900
agttcgttca	tatcttggaa	ggagtgtcgt	caaaattatc	cagatatgac	gaaggagact	3960
tgttttcttc	ttttctgtca	tttaccgtga	agggcagctt	caaatatgtg	gatgtacctc	4020
aaacccggat	ggagcgtggc	gagccctacg	tgactttcgt	cgcgcatctc	caggatgtcc	4080
tgctgtgata	ggttcaatgag	gagatgtaca	tagaaaaggt	atttgatcaa	tggatcaaca	4140
gctccatgaa	cgtgatctgc	acctggtttg	cggacccgat	ggacttacag	cttcatattt	4200
atcagttgaa	aacactaa	aggggtgtaa	agaaaacctc	cagagatttc	cgatgtcaag	4260
gggtctctga	ctccacctta	aacagcaaga	cctatgaaac	gatccgggac	cgtctcactg	4320
tggaggaagc	cacagcatca	gtgagtgaag	gtggggcagc	gcagggcagc	agcatggaag	4380
acacgcatga	ggagagcaga	gaagacgatt	agaccatttg	gtcctagagt	ctgctgggac	4440
agagtctctg	aatcagtgca	tgctcctagt	ctgttagtia	aaccocatag	gaattttcgt	4500
tcaactacca	tgcccatgag	atgtttatca	atacaactgc	cattttagct	atgtgtgacc	4560

```

aagattagca aatgaccttc atatccactg atttccctgat gtccatgtct atattgtttac 4620
aagcaatatg gagcaccatt cttttaaatc tgttcatgga gaatacatag tctaacccact 4680
aggcgtgtcc ctgttatcag caaagatcaa tgatgcttca ttcattgtact atgtatgcat 4740
tggtgggtaaa tggatgtgag ggcaagtaca tcaagtacat tcactctgtt tcacgtatgt 4800
ggatgccagtg taattaaatg agtacgtaaa taaataatatt aaaacacata gatctgtcttt 4860
gtgttttttat tttttttttt tgaaaaacaa aaggccaagtc tccacaattt aacttttgat 4920
gctttctgtt cccctaaaaac caaaaaatga accccttgtg tcgttgttaa cccatccctt 4980
catttactaca tataattagc caaaaaaaa aggatgggcta cataccaatg gattgattct 5040
cttaattgcc accgcaaggg ggcgatccta tcatgactta acatcaagcg cgcaggttcaa 5100
aactactgtc tctgtgtcaa gttttctcct cttaaatgtt attttgtctt tacgtctcaa 5160
ctgtgtatgt aaaaaaaacg aatatttaaa ttacaacctt agactaaaaa tgtgtttata 5220
ataagatgtg gatatttctc tcatagattt gtaaccataa tttaaattat ttgtttccac 5280
acgtgtttttt atactgttca tgtacattgc attttgatct gtaactgcac aacctggggg 5340
tttctctcag agctatttct ttccatgtaa agtagtggtt ccatcttgct ttgtccttat 5400
ataaagccta cagttatgga agtgtggaaa actgtggctt ctcaataaat attcagatgt 5460
cctaagaata t 5471

```

```

<210> 35
<211> 1390
<212> PRT
<213> Homo sapiens

```

```

<400> 35
Pro Gly Glu Lys Arg Pro Pro Gly Arg Ala Asp Pro Arg Leu Pro Pro
1 5 10 15
Ala Pro Pro Thr Phe Cys Thr Pro Ser Arg Gly Leu Gln Arg Pro Arg
20 25 30
Ser Pro Gly Ala Thr Met Leu Asp Pro Ser Ser Ser Glu Glu Ser
35 40 45
Asp Glu Ile Val Glu Glu Glu Ser Gly Lys Glu Val Leu Gly Ser Ala
50 55 60
Pro Ser Gly Ala Arg Leu Ser Pro Ser Arg Thr Ser Glu Gly Ser Ala
65 70 75 80
Gly Ser Ala Gly Leu Gly Gly Gly Ala Gly Ala Gly Ala Gly Val
85 90 95
Gly Ala Gly Gly Gly Gly Gly Ser Gly Ala Ser Ser Gly Gly Ala
100 105 110
Gly Gly Leu Gln Pro Ser Ser Arg Ala Gly Gly Gly Arg Pro Ser Ser
115 120 125
Pro Ser Pro Ser Val Val Ser Glu Lys Glu Lys Glu Glu Leu Glu Arg
130 135 140
Leu Gln Lys Glu Glu Glu Glu Arg Lys Lys Arg Leu Gln Leu Tyr Val
145 150 155 160
Phe Val Met Arg Cys Ile Ala Tyr Pro Phe Asn Ala Lys Gln Pro Thr
165 170 175
Asp Met Ala Arg Arg Gln Gln Lys Ile Ser Lys Gln Gln Leu Gln Thr
180 185 190
Val Lys Asp Arg Phe Gln Ala Phe Leu Asn Gly Glu Thr Gln Ile Met
195 200 205
Ala Asp Glu Ala Phe Met Asn Ala Val Gln Ser Tyr Tyr Glu Val Phe
210 215 220
Leu Lys Ser Asp Arg Val Ala Arg Met Val Gln Ser Gly Gly Cys Ser
225 230 235 240
Ala Asn Asp Ser Arg Glu Val Phe Lys Lys His Ile Glu Lys Arg Val
245 250 255
Arg Ser Leu Pro Glu Ile Asp Gly Leu Ser Lys Glu Thr Val Leu Ser
260 265 270
Ser Trp Met Ala Lys Phe Asp Ala Ile Tyr Arg Gly Glu Glu Asp Pro

```

275	280	285
Arg Lys Gln Gln Ala Arg Met Thr Ala Ser Ala Ala Ser Glu Leu Ile		
290	295	300
Leu Ser Lys Glu Gln Leu Tyr Glu Met Phe Gln Asn Ile Leu Gly Ile		
305	310	315
Lys Lys Phe Glu His Gln Leu Leu Tyr Asn Ala Cys Gln Leu Asp Asn		
	320	330
Pro Asp Glu Gln Ala Ala Gln Ile Arg Arg Glu Leu Asp Gly Arg Leu		
	340	350
Gln Met Ala Asp Gln Ile Ala Arg Glu Arg Lys Phe Pro Lys Phe Val		
	355	365
Ser Lys Glu Met Glu Asn Met Tyr Ile Glu Glu Leu Lys Ser Ser Val		
	370	380
Asn Leu Leu Met Ala Asn Leu Glu Ser Met Pro Val Ser Lys Gly Gly		
385	390	395
Glu Phe Lys Leu Gln Lys Leu Lys Arg Ser His Asn Ala Ser Ile Ile		
	405	415
Asp Met Gly Glu Glu Ser Glu Asn Gln Leu Ser Lys Ser Asp Val Val		
	420	430
Leu Ser Phe Ser Leu Glu Val Val Ile Met Glu Val Gln Gly Leu Lys		
	435	445
Ser Leu Ala Pro Asn Arg Ile Val Tyr Cys Thr Met Glu Val Glu Gly		
	450	460
Gly Glu Lys Leu Gln Thr Asp Gln Ala Glu Ala Ser Lys Pro Thr Trp		
465	470	475
Gly Thr Gln Gly Asp Phe Ser Thr Thr His Ala Leu Pro Ala Val Lys		
	485	490
Val Lys Leu Phe Thr Glu Ser Thr Gly Val Leu Ala Leu Glu Asp Lys		
	500	505
Glu Leu Gly Arg Val Ile Leu His Pro Thr Pro Asn Ser Pro Lys Gln		
	515	520
Ser Glu Trp His Lys Met Thr Val Ser Lys Asn Cys Pro Asp Gln Asp		
	530	535
Leu Lys Ile Lys Leu Ala Val Arg Met Asp Lys Pro Gln Asn Met Lys		
545	550	555
His Ser Gly Tyr Leu Trp Ala Ile Gly Lys Asn Val Trp Lys Arg Trp		
	565	570
Lys Lys Arg Phe Phe Val Leu Val Gln Val Ser Gln Tyr Thr Phe Ala		
	580	585
Met Cys Ser Tyr Arg Glu Lys Lys Ala Glu Pro Gln Glu Leu Leu Gln		
	595	600
Leu Asp Gly Tyr Thr Val Asp Tyr Thr Asp Pro Gln Pro Gly Leu Glu		
	610	615
Gly Gly Arg Ala Phe Phe Asn Ala Val Lys Glu Gly Asp Thr Val Ile		
625	630	635
Phe Ala Ser Asp Asp Glu Gln Asp Arg Ile Leu Trp Val Gln Ala Met		
	645	650
Tyr Arg Ala Thr Gly Gln Ser His Lys Pro Val Pro Pro Thr Gln Val		
	660	665
Gln Lys Leu Asn Ala Lys Gly Gly Asn Val Pro Gln Leu Asp Ala Pro		
	675	680
Ile Ser Gln Phe Tyr Ala Asp Arg Ala Gln Lys His Gly Met Asp Glu		
	690	695
Phe Ile Ser Ser Asn Pro Cys Asn Phe Asp His Ala Ser Leu Phe Glu		
705	710	715
Met Val Gln Arg Leu Thr Leu Asp His Arg Leu Asn Asp Ser Tyr Ser		
	725	730
Cys Leu Gly Trp Phe Ser Pro Gly Gln Val Phe Val Leu Asp Glu Tyr		
	735	

740					745					750					
Cys	Ala	Arg	Asn	Gly	Val	Arg	Gly	Cys	His	Arg	His	Leu	Cys	Tyr	Leu
		755					760					765			
Arg	Asp	Leu	Leu	Glu	Arg	Ala	Glu	Asn	Gly	Ala	Met	Ile	Asp	Pro	Thr
		770					775					780			
Leu	Leu	His	Tyr	Ser	Phe	Ala	Phe	Cys	Ala	Ser	His	Val	His	Gly	Asn
		785					790					795			800
Arg	Pro	Asp	Gly	Ile	Gly	Thr	Val	Thr	Val	Glu	Glu	Lys	Glu	Arg	Phe
				805					810					815	
Glu	Glu	Ile	Lys	Glu	Arg	Leu	Arg	Val	Leu	Leu	Glu	Asn	Gln	Ile	Thr
			820					825					830		
His	Phe	Arg	Tyr	Cys	Phe	Pro	Phe	Gly	Arg	Pro	Glu	Gly	Ala	Leu	Lys
			835					840					845		
Ala	Thr	Leu	Ser	Leu	Leu	Glu	Arg	Val	Leu	Met	Lys	Asp	Ile	Val	Thr
			850				855					860			
Pro	Val	Pro	Gln	Glu	Glu	Val	Lys	Thr	Val	Ile	Arg	Lys	Cys	Leu	Glu
				870						875				880	
Gln	Ala	Ala	Leu	Val	Asn	Tyr	Ser	Arg	Leu	Ser	Glu	Tyr	Ala	Lys	Ile
				885						890				895	
Glu	Glu	Asn	Gln	Lys	Asp	Ala	Glu	Asn	Val	Gly	Arg	Leu	Ile	Thr	Pro
			900					905					910		
Ala	Lys	Lys	Leu	Glu	Asp	Thr	Ile	Arg	Leu	Ala	Glu	Leu	Val	Ile	Glu
			915					920					925		
Val	Leu	Gln	Gln	Asn	Glu	Glu	His	His	Ala	Glu	Pro	His	Val	Asp	Lys
			930				935					940			
Gly	Glu	Ala	Phe	Ala	Trp	Ser	Ser	Asp	Leu	Met	Val	Glu	His	Ala	Glu
			945			950				955				960	
Thr	Phe	Leu	Ser	Leu	Phe	Ala	Val	Asp	Met	Asp	Ala	Ala	Leu	Glu	Val
				965					970					975	
Gln	Pro	Pro	Asp	Thr	Trp	Asp	Ser	Phe	Pro	Leu	Phe	Gln	Leu	Leu	Asn
			980						985				990		
Asp	Phe	Leu	Arg	Thr	Asp	Tyr	Asn	Leu	Cys	Asn	Gly	Lys	Phe	His	Lys
			995					1000					1005		
His	Leu	Gln	Asp	Leu	Phe	Ala	Pro	Leu	Val	Val	Arg	Tyr	Val	Asp	Leu
			1010				1015					1020			
Met	Glu	Ser	Ser	Ile	Ala	Gln	Ser	Ile	His	Arg	Gly	Phe	Glu	Arg	Glu
			1025			1030				1035				1040	
Ser	Trp	Glu	Pro	Val	Lys	Ser	Leu	Thr	Ser	Asn	Leu	Pro	Asn	Val	Asn
				1045						1050				1055	
Leu	Pro	Asn	Val	Asn	Leu	Pro	Lys	Val	Pro	Asn	Leu	Pro	Val	Asn	Ile
				1060				1065					1070		
Pro	Leu	Gly	Ile	Pro	Gln	Met	Pro	Thr	Phe	Ser	Ala	Pro	Ser	Trp	Met
			1075				1080						1085		
Ala	Ala	Ile	Tyr	Asp	Ala	Asp	Asn	Gly	Ser	Gly	Thr	Ser	Glu	Asp	Leu
			1090												

	1205		1210		1215
Leu Leu Val Ala Lys Phe Val Thr Ile Leu Glu Gly Val Leu Ala Lys					
	1220		1225		1230
Leu Ser Arg Tyr Asp Glu Gly Thr Leu Phe Ser Ser Phe Leu Ser Phe					
	1235		1240		1245
Thr Val Lys Ala Ala Ser Lys Tyr Val Asp Val Pro Lys Pro Gly Met					
	1250		1255		1260
Asp Val Ala Asp Ala Tyr Val Thr Phe Val Arg His Ser Gln Asp Val					
	1265		1270		1275
Leu Arg Asp Lys Val Asn Glu Glu Met Tyr Ile Glu Arg Leu Phe Asp					
	1285		1290		1295
Gln Trp Tyr Asn Ser Ser Met Asn Val Ile Cys Thr Trp Leu Thr Asp					
	1300		1305		1310
Arg Met Asp Leu Gln Leu His Ile Tyr Gln Leu Lys Thr Leu Ile Arg					
	1315		1320		1325
Val Val Lys Lys Thr Tyr Arg Asp Phe Arg Leu Gln Gly Val Leu Asp					
	1330		1335		1340
Ser Thr Leu Asn Ser Lys Thr Tyr Glu Thr Ile Arg Asn Arg Leu Thr					
	1345		1350		1355
Val Glu Glu Ala Thr Ala Ser Val Ser Glu Gly Gly Leu Gln Gly					
	1365		1370		1375
Ile Ser Met Lys Asp Ser Asp Glu Glu Asp Glu Glu Asp Asp					
	1380		1385		1390

<210> 36

<211> 4828

<212> DNA

<213> Homo sapiens

<400> 36

agtggcgtcg	gaactgcaaa	gcacctgtga	gottgcggaa	gtcagttcag	actccagccc	60
gctccagccc	ggcccgaacc	gaccgcaacc	ggcgctgcc	ctcgctcggc	gtcccggccc	120
agccatgggc	ccttgaggcc	gcagcctctc	ggcgctgctg	ctgctgctgc	aggtctctctc	180
ttggctctgc	caggagccgg	agccctgcc	ccctggcttt	gacgcggaga	gctacacgtt	240
cacggtgccc	cgcgccacc	tggagagagg	cgcgctcgtg	ggcagagtga	attttgaaga	300
ttgcacgggt	cgacaaagga	cagcctattt	ttccctcgac	acccgattca	aagtggccac	360
agatgggttg	attacagtca	aaaggcctct	acggtttcat	aaccacaga	tccatttctt	420
ggctctacgc	tgggactcca	ctcacagaaa	gtttccacc	aaagtcaagg	tgaatacagt	480
ggggcaccac	caccgcccc	cgccccatca	ggcctccgtt	tctggaatcc	aagcagaatt	540
gctcacattt	cccaactcct	ctcctggcct	cagaagacag	aagagagact	gggttatcc	600
tcctatcagc	tgccccgaaa	atgaaaaagg	cccatcttct	aaaaacctgg	ttcatgatca	660
atccacaaca	gacaaaagag	gcaaggtttt	ctacagcctc	actggccaag	gagctgacac	720
accocctggt	ggtgtcttta	ttattgaaag	agaaacagga	tggctgaagg	tgacagagcc	780
cttggtataga	gaacgcattg	ccacatacac	tctctctctc	cacgctgtgt	catccaaacg	840
gaatgcagtt	gaggatccaa	tggagatttt	gatcacggta	accgatcaga	atgacacaaa	900
gcccgaaatc	accagggagg	tctttaaggg	gtctgtcatg	gaaggtgtct	ttccaggaac	960
ctctgtgatg	gaggtcacag	ccacagacgc	ggacgatgat	gtgaacacct	acaatggccc	1020
catcgcttac	accatctcca	gccaaagatcc	tgagctccct	gacaaaataa	tgctccacct	1080
taacagggaac	acaggagtca	tcagttgtgt	caccactggg	ctggaccgag	agagtctccc	1140
tacgtatacc	ctggtgtgtc	aaagtcgtga	ccttcaaggt	gaggggttaa	gcacaacagc	1200
aaacagctgt	atcacagtca	ctgacaccaa	cgataactct	ccgatctcca	atcccaccac	1260
gtacacgggt	caggtgcctg	agaacgaggc	tacaagtctc	atcacacac	tgaagtgcac	1320
tgatgctgat	gcccccaata	ccccagcgtg	ggaggtcgtg	tacacacat	tgaatgatga	1380
tgggtgacaa	ttgtgtcgtc	ccacaatacc	agtgaaacac	gatggcattt	tgaaaacagc	1440
aaagggcctg	gattttgagg	ccaagcagca	gtacattcta	cacgtagacg	tgacgaatgt	1500
ggtacacttt	gaggtctctc	tcaccaacct	cacagccacc	gtcaccgtgt	atgtcgtgga	1560
tgtgaatgaa	gcccccatct	ttgtgcctcc	tgaaaagaga	gtggaagtgt	ccgaggacct	1620

tggcggtgggc	caggaaatca	catcctacac	tgcccaggag	ccagacacat	ttatggaaca	1680
gaaaataaca	tatcgagattt	ggagagacac	tgcccaactgg	ctggagatta	atccggacac	1740
tggtgccaatt	tcocactcggg	ctgagctgga	caggggaggat	tttgagcacg	tgaagaacag	1800
cacgtacaca	gcocataatca	tagctacaga	caatggttct	ccagttgcta	ctggacacagg	1860
gacactctctg	ctgatccogt	ctgatgtgaa	tgacaacgcc	cccatcaccg	aacctcgaa	1920
tatatctctc	tgtgagagga	atccaaagcc	tcaggtccta	aacatcatgt	atcgacacct	1980
tcctcccaat	acatctccct	tcacagcaga	actaacacac	ggggcgagtg	ccaactggac	2040
cattcagtag	aagcaccaca	cccaagaatc	tatcattttg	aagccaaaga	tggtccttaga	2100
gggggtgac	tacaaataac	atctcaagct	catggataac	acataataag	accaagttagc	2160
caccttagag	gtacogctgt	gtgactgtga	aggggccgcc	ggcgctctga	ggaaggcaca	2220
gcctgtcgaa	gcaggattgc	aaattcctgc	catctcgggg	attcttggag	gaattcttgc	2280
tttgtaatt	ctgatctctg	tgctcttgct	gtttcttcgg	aggagagcgg	tggtcaaaaga	2340
gccttctactg	ccocaggag	atgacacccg	ggacaaagtt	tattactatg	atgaagaagg	2400
aggcgagaa	gaggccaagg	actttgaact	gagccagctg	cacaggggcc	tgagcgctcg	2460
gcctgaagt	actgtaacg	acgttgccac	aacctcatg	agtgcccccc	ggtatcttcc	2520
ccgcctgcc	aatcccgatg	aaattggaaa	ttttattgat	gaaaatctga	aagcggtctga	2580
tactgacccc	acagccccc	cttatgattc	tctgctctgt	tttgactatg	aagggaagcgg	2640
ttccgaagct	gcgatcttga	gctccctgaa	ctcctcagag	tcagacaaag	accaggacta	2700
tgactacttg	aacgaatggg	gcaatcgctt	caagaagctg	gctgacatgt	acggaggcgg	2760
cgaggccagac	taggggactc	gagagaggcg	ggccccaagc	caattgtctg	ggaattcgag	2820
aaatcacggt	gctggtgggt	tttcagctcc	cttcctctga	gatgagtttc	tggggaaaaa	2880
aaagagactg	gtttagtgat	cagtttagtat	agctttatag	ctctccactc	ttatagctct	2940
ataaagtgtt	tgtagaaaaa	gtttogaact	atttcttaaa	gctttttttt	ttttcccatc	3000
actctttaca	tggtgtggat	gtccaaaaga	taccaaaatt	ttatattcag	agaagaagag	3060
ctttagcatc	agaaggttica	cccagcacct	tgacagattt	cttaaggaaat	tttgtctcac	3120
ttttaaaaag	aagggtgaaa	gtcagctact	tgttttctgt	gtttttctgt	ataataattt	3180
ttaaaaaaa	tttgtgtgct	tctgctcaat	actacactgg	tggtgcccct	tgccctttttt	3240
ttttttttta	agacagggtc	tcattctatc	ggccaggctg	gagtgacgtg	gtgcaatcac	3300
agctcactgc	agccttgctc	tcccaggctc	aagctatcct	tgacctccag	ccctccaaag	3360
agctgggacc	acaggcatgc	accactacgc	atgactaatt	ttttaaatat	ttgagacggg	3420
gtctccctgt	gttaccccag	ctggtctcaa	actcctgggc	tcaagtgtac	ctcccatctt	3480
ggcctccag	agttatggga	ttacagacat	gagccactgc	acctgccacc	ctccccaact	3540
ccctgcattt	ttttaagaga	cagtttcgct	ccatcgccca	ggcctgggat	gcagtgtgtg	3600
gatcatagct	cactgtaacc	tcaaacctct	gggctoaagg	agttctccca	ccagcctcct	3660
ttttattttt	ttgtacagat	ggggtcttgc	tatgttgccc	aagctggctg	taaaactcctg	3720
gcctccaca	atcctctctg	cttgccccc	caaagtctgt	ggattgtggg	catgagctgc	3780
tgctgccagc	ctccatgttt	taatatcaac	ctcactcct	gaattcagtt	gotttgccca	3840
agataggagt	tctctgatgc	agaaattatt	gggtcttttt	agggtaaaga	gtttgtgtct	3900
ttgtctggcc	acactctgac	taggtattgt	ctactctgaa	gacotttaat	ggcttccctc	3960
tttcatctcc	tgagtattga	acttgcaatg	gcagcgtatc	cagtgtactg	ttctgagtaa	4020
gtgtgtctac	taattgttat	ttagctctga	agcagaagtg	atatactcca	ggaactagaa	4080
tagtgctcaa	agtgctccag	ccaaagacag	agcggaacta	tgaaaagtgt	ctgtggagat	4140
ggcaggagag	cttgctcaatg	agcctggcaa	tttagcaaac	tgatgctgag	gatgatggag	4200
ctgaggtctac	ctcatctctg	aaaattctgg	aggaatgag	ggagctctca	catgtgtttc	4260
tgacacaaga	tcogtgggtt	gtactcaaa	ccagaatacc	caaagtgcct	gcttttgatg	4320
atgtctacag	aaaatgctgt	ctgagctgaa	acattttcgg	caatccagg	tggtcacaga	4380
aaacccgaaa	tattccaaat	tccaaatttt	ttcttaggag	caagaagaaa	atgtggccct	4440
aaagggtggt	agtgaggggg	tagggggtag	tagggatctg	gatttggact	cttttttatt	4500
taaatgtgaa	tttcaacttt	tgacaatcaa	agaaaagact	tttgttgaaa	tagcttttact	4560
gtttctcaag	tggttttgag	aaaaaaatca	acctctgcaat	cactttttgg	aattgtcttg	4620
gtttttcggc	agtttccag	atatogaata	tagttctgtg	tagaagaatg	acagtgtagt	4680
ttgagtgtat	acatgtgtgg	gtgctgataa	ttgtgtattt	tttttggggg	tggaagagga	4740
aaacaattca	agctgagaaa	agttattcca	aagatgcatt	tttataaatt	ttatttaaca	4800
attttgttaa	acccaataaa	aaaaaaa				4828

<210> 37

<211> 882

<212> PRT

<213> Homo sapiens

<400> 37

```

Met Gly Pro Trp Ser Arg Ser Leu Ser Ala Leu Leu Leu Leu Gln
 1          5          10          15
Val Ser Ser Trp Leu Cys Gln Glu Pro Glu Pro Cys His Pro Gly Phe
 20
Asp Ala Glu Ser Tyr Thr Phe Thr Val Pro Arg Arg His Leu Glu Arg
 35          40          45
Gly Arg Val Leu Gly Arg Val Asn Phe Glu Asp Cys Thr Gly Arg Gln
 50          55          60
Arg Thr Ala Tyr Phe Ser Leu Asp Thr Arg Phe Lys Val Gly Thr Asp
 65          70          75          80
Gly Val Ile Thr Val Lys Arg Pro Leu Arg Phe His Asn Pro Gln Ile
 85          90          95
His Phe Leu Val Tyr Ala Trp Asp Ser Thr Tyr Arg Lys Phe Ser Thr
100          105          110
Lys Val Thr Leu Asn Thr Val Gly His His Arg Pro Pro Pro His
115          120          125
Gln Ala Ser Val Ser Gly Ile Gln Ala Glu Leu Leu Thr Phe Pro Asn
130          135          140
Ser Ser Pro Gly Leu Arg Arg Gln Lys Arg Asp Trp Val Ile Pro Pro
145          150          155          160
Ile Ser Cys Pro Glu Asn Glu Lys Gly Pro Phe Pro Lys Asn Leu Val
165          170          175
Gln Ile Lys Ser Asn Lys Asp Lys Glu Lys Val Phe Tyr Ser Ile
180          185          190
Thr Gly Gln Gly Ala Asp Thr Pro Pro Val Gly Val Phe Ile Ile Glu
195          200          205
Arg Glu Thr Gly Trp Leu Lys Val Thr Glu Pro Leu Asp Arg Glu Arg
210          215          220
Ile Ala Thr Tyr Thr Leu Phe Ser His Ala Val Ser Ser Asn Gly Asn
225          230          235          240
Ala Val Glu Asp Pro Met Glu Ile Leu Ile Thr Val Thr Asp Gln Asn
245          250          255
Asp Asn Lys Pro Glu Phe Thr Gln Glu Val Phe Lys Gly Ser Val Met
260          265          270
Glu Gly Ala Leu Pro Gly Thr Ser Val Met Glu Val Thr Ala Thr Asp
275          280          285
Ala Asp Asp Asp Val Asn Thr Tyr Asn Ala Ala Ile Ala Tyr Thr Ile
290          295          300
Leu Ser Gln Asp Pro Glu Leu Pro Asp Lys Asn Met Phe Thr Ile Asn
305          310          315          320
Arg Asn Thr Gly Val Ile Ser Val Val Thr Thr Gly Leu Asp Arg Glu
325          330          335
Ser Phe Pro Thr Tyr Thr Leu Val Val Gln Ala Ala Asp Leu Gln Gly
340          345          350
Glu Gly Leu Ser Thr Thr Ala Thr Ala Val Ile Thr Val Thr Asp Thr
355          360          365
Asn Asp Asn Pro Pro Ile Phe Asn Pro Thr Thr Tyr Lys Gly Gln Val
370          375          380
Pro Glu Asn Glu Ala Asn Val Val Ile Thr Thr Leu Lys Val Thr Asp
385          390          395          400
Ala Asp Ala Pro Asn Thr Pro Ala Trp Glu Ala Val Tyr Thr Ile Leu
405          410          415
Asn Asp Asp Gly Gly Gln Phe Val Val Thr Thr Asn Pro Val Asn Asn
420          425          430
Asp Gly Ile Leu Lys Thr Ala Lys Gly Leu Asp Phe Glu Ala Lys Gln

```

435					440					445					
Gln	Tyr	Ile	Leu	His	Val	Ala	Val	Thr	Asn	Val	Val	Pro	Phe	Glu	Val
450						455					460				
Ser	Leu	Thr	Thr	Ser	Thr	Ala	Thr	Val	Thr	Val	Asp	Val	Leu	Asp	Val
465					470					475					480
Asn	Glu	Ala	Pro	Ile	Phe	Val	Pro	Pro	Glu	Lys	Arg	Val	Glu	Val	Ser
				485					490					495	
Glu	Asp	Phe	Gly	Val	Gly	Gln	Glu	Ile	Thr	Ser	Tyr	Thr	Ala	Gln	Glu
			500					505					510		
Pro	Asp	Thr	Phe	Met	Glu	Gln	Lys	Ile	Thr	Tyr	Arg	Ile	Trp	Arg	Asp
		515					520					525			
Thr	Ala	Asn	Trp	Leu	Glu	Ile	Asn	Pro	Asp	Thr	Gly	Ala	Ile	Ser	Thr
		530				535					540				
Arg	Ala	Glu	Leu	Asp	Arg	Glu	Asp	Phe	Glu	His	Val	Lys	Asn	Ser	Thr
					550				555						560
Tyr	Thr	Ala	Leu	Ile	Ile	Ala	Thr	Asp	Asn	Gly	Ser	Pro	Val	Ala	Thr
				565					570					575	
Gly	Thr	Gly	Thr	Leu	Leu	Leu	Ile	Leu	Ser	Asp	Val	Asn	Asp	Asn	Ala
			580					585					590		
Pro	Ile	Pro	Glu	Pro	Arg	Thr	Ile	Phe	Phe	Cys	Glu	Arg	Asn	Pro	Lys
		595					600					605			
Pro	Gln	Val	Ile	Asn	Ile	Ile	Asp	Ala	Asp	Leu	Pro	Pro	Asn	Thr	Ser
		610				615					620				
Pro	Phe	Thr	Ala	Glu	Leu	Thr	His	Gly	Ala	Ser	Ala	Asn	Trp	Thr	Ile
		625			630				635						640
Gln	Tyr	Asn	Asp	Pro	Thr	Gln	Glu	Ser	Ile	Ile	Leu	Lys	Pro	Lys	Met
				645					650					655	
Ala	Leu	Glu	Val	Gly	Asp	Tyr	Lys	Ile	Asn	Leu	Lys	Leu	Met	Asp	Asn
			660					665					670		
Gln	Asn	Lys	Asp	Gln	Val	Thr	Thr	Leu	Glu	Val	Ser	Val	Cys	Asp	Cys
		675					680					685			
Glu	Gly	Ala	Ala	Gly	Val	Cys	Arg	Lys	Ala	Gln	Pro	Val	Glu	Ala	Gly
		690				695					700				
Leu	Gln	Ile	Pro	Ala	Ile	Leu	Gly	Ile	Leu	Gly	Gly	Ile	Leu	Ala	Leu
				710					715					720	
Leu	Ile	Leu	Ile	Leu	Leu	Leu	Leu	Phe	Leu	Arg	Arg	Arg	Arg	Ala	Val
				725				730						735	
Val	Lys	Glu	Pro	Leu	Leu	Pro	Pro	Glu	Asp	Asp	Thr	Arg	Asp	Asn	Val
			740					745					750		
Tyr	Tyr	Tyr	Asp	Glu	Glu	Gly	Gly	Gly	Glu	Glu	Asp	Gln	Asp	Phe	Asp
		755					760					765			
Leu	Ser	Gln	Leu	His	Arg	Gly	Leu	Asp	Ala	Arg	Pro	Glu	Val	Thr	Arg
		770				775					780				
Asn	Asp	Val	Ala	Pro	Thr	Leu	Met	Ser	Val	Pro	Arg	Tyr	Leu	Pro	Arg
		785			790				795		</				

<210> 38
 <211> 4521
 <212> DNA
 <213> Homo sapiens

<400> 38
 attctcatca cttgcacatc agtgtgtgcc cacaagagcc agctctcccg agcccgtaac 60
 ctctgcattc caagagctgc agtttcagcc gcgacagcaa gaacggcaga gcggcgagcc 120
 gcggcgccgg cggcgccgga ggcaggagca gcctggcggg gtccgagggg ctccgcgggc 180
 gcagggaagg gagcagagat atcctctgag agccaagcaa agaactataa ggaagggaag 240
 aggaatgagg ctggatcagg tgcagtgaaa aaggcaatct caagagtgagg gcaactacta 300
 gcacacagct gcacggtgcc atcagcatga gaacttaccg ctacttcttg ctgctctttt 360
 ggtgtggcca gccctaccga actctctcaa ctccactatc aaagaggact agtggtttcc 420
 cagcaaaaga aaggccctcg gagctctctg gaaacagcaa aaatgagctg aaccgttcaa 480
 aaaggagctg gatgtggaat cagtctcttc tctcgaggga atacacagga tccgattatc 540
 agtatgtggg caagttacat tcagaccagg atagaggaga tggatcactt aaatatatcc 600
 tttcaggaga tggagcagga gatctcttca ttattaatga aaacacaggg gacatcacagg 660
 ccaccaagag gctggacagg gaagaaaaac ccgtttacat ccttcgagct caagctataa 720
 acagaaggac agggagaccc gtggagcccg agtctgaatt catcatcaag atccatgaca 780
 tcaatgacaa tgaaaccaata ttccaccaag aggtttacac agccactgtc cctgaaatgt 840
 ctgatgtcgg tacatttgtt gtccaagtca ctgcgacgga tgcagatgat ccaacatagt 900
 ggaacagtcg taagtgtgtc tacagttatc tacagggaca gccctatttt tcatgtgaat 960
 cagaaaacagg tattatcaag acagctttgc tcaacatgga tcgagaaaaa agggagcagt 1020
 accaagtgtg gatttcaagcc aagatatgag cggcgccagat gggaggatta tctgggacca 1080
 ccaccgtgag catccactcg actgatgtca acgacaaccc tcccgtattc cccagagata 1140
 cataccagtt taaaactcct gaattcttct caccggggac accaattggc agaattcaag 1200
 ccagcgacgc tgaatgtgga gaaaatgctg aaattgagta cagcatcaca gcggtgagg 1260
 ggcgtggatg gtttgaatgc ataccgcagc aggaaaacca ggaaggattt ataatctgtca 1320
 aaaagctctt ggaacttgaa aagaagaagg tgatatccct taaagtggaa gccctcaact 1380
 cttaigtgga gccacgattt ctctacttgg gccctttcaa agatttcagcc acggttagaa 1440
 ttgtggttga ggaatgtgat gagccacctg tcttcagcaa actggcctac atcttcaaaa 1500
 taagagaaga tgcctagata aacaccacaa taggctccgt cacagcccaa gatccagatg 1560
 ctgccaggaa tctgtcgaag tactctgtag atcgacacac agtatggac agaattttca 1620
 acattgattc tggaaatggt tcgattttta catcgaaact tcttgaccga gaaacactgc 1680
 tatggcacia cattacagtg atagcaacag agatcaataa tccaaagcaa agtagtcgag 1740
 tacctctata tattaaagtt ctatagtgca atgacaacgc ccagaattt gctgagttct 1800
 atgaactttt tgtctgtgaa aaagcaaaag cagatcagtt gatccagacc ctgcagctg 1860
 ttgacaagga tgacccttat agtggacacc aattttcgct tctctggccc ctgaagcag 1920
 ccagtggtct aacttttacc attcaagaca acaagaacaa caccggcgga atcttaactc 1980
 ggaataatgg ctataataga caccgatga gccactatct ctgctctgtg gtcatttcag 2040
 caacagcata ccagatgcaa agcagctgac ggaactgtac ctgcgggtc tgtgcatgtg 2100
 accaccaggg gaacatgcaa tctcgccatg cggagggcgt catccacccc acgggactga 2160
 gcacgggggc tctggttgcc atctctctg ctactagtgt cactctagtgt cctggtgtg 2220
 tgatttcagg tctgaggcgg cagcgaaaaa aagagccctt gatcatcttc aaaggaggca 2280
 tcagagataa cattgtcagt tacaacgacg aaggtggtgg agaggaggac accagagctt 2340
 ttgatattcg caccctgagg aatcctgaag ccatagagga caacaaatta cgaaggagca 2400
 ttgtgccgga agcccttttc ctaccocgac agctcgcagc aaactccaaac gactccagct 2460
 tcagagattt cattaaaccaa aggttaaagg aaaatgacac ggaccccaact gccccagcat 2520
 acgactcctt ggccacttac gcctatgaag ccaactggct cgtggcggat tccctgagct 2580
 cgtgtggagtc agtgaccacg gatgcagatc aagactatga ttaccttagt gactggggac 2640
 ctcgatcaaa aagacttgca gatatgtatg gaggagtgga cagtgcacaa gactcctaact 2700
 ctgttgccct ttctattttc caatacgaca ctgaaatagt tgaagtggct atttctttat 2760
 atttatccac tactccgtga aggcttctct gtctctaccg ttccaaaagg caatggctgc 2820
 agtccgtgtg gctccaattgt tagagacttt ttctctagta acttttatga gcttccaaag 2880
 ggcacatttt tattttttag tgcattccagt taaccaagtc agcccaacag cgaggtgcgg 2940
 gaggggagga cagggaacag taatttcact tgttctcagg gacgctgccc ctactccgct 3000
 gtccgtgtgt ttactacac tccatgtcag gtcagccaac tgccttaact gtacatttca 3060

```

caggctaatt ggataaagga ctgtgcttta aagataaaaa tatcatcata gtaaaagaaa 3120
tgaggggcata tcggctcaca aagagataaaa ctacataggg gtgtttattt gtgtcacaaa 3180
gaatttataaa taacactgtgc ccatgtctatt tgtttctcaa gaactttctc tgcacatcac 3240
tactattcaa aacctcaaat ccaccocatat gttaaaattc tcattactct taaggaaatag 3300
aagcaaatata aacggtaaca tccaaaagca accacaaacc tagtagcagt tcattctctc 3360
cactaaactca tagtttgtta tatctctagac tagacatggc aaagtttggc ttgttaccat 3420
ataaaggggg agggaaatag ctaataatgt taaccaagga aatataattt accatacatt 3480
taaaagtttgg gccaccacat gtatcaacggg tcaacttgaaa ttctttcagc tatcagtagg 3540
ctaattgtcaa aattgttttaa aaattcttga aagaattttc ctgagacaaa ttttaacttc 3600
ttgtctatag ttgtcagtat tattctacta tactgtacat gaaagtagca gtgtgaagta 3660
caataattca tattcttcat atccttctta cagcactaag ttgaattagt aaagttagat 3720
taataaaaaa ttaaatctca ctctaggagt tcaaggtaga ggtagagcc agccacattt 3780
gaacctataa cctgcccttt gacatctgga aacctctaca tatttatata acgtatata 3840
tttgataaaa caacattgag attatgatga aaacctacat attccatggt tggagaagcc 3900
ttggaagagg aaaaattggat tcccttaaac aaaaagtgtt aagattgtaa ttaaaatgat 3960
agttgatttt caaaagcatt aatttttttt cattgttttt aactttgctt tcatgacct 4020
ctgcctatcc ttgactttga actaatgata aagtaatgat ctcaaaacta gacagaaaaa 4080
taagttaaaa tccatccaat ctattatttc tctaattatg caattagcct catagttatt 4140
ctccagagga cccaactgaa ctgaactaat cctctcgcca gattcaaaat gtttatttca 4200
cacgtgtgtc taatggcact tatcattaga atcttacott gtgcagtcac cagaaaattcc 4260
agcgctacta aatgaaaaca tccctgtttt gaaaacctaa aagacagcgt ctgtatatat 4320
atatacttaa gaatatgctg acttcaacta ttagtcttag ggatttattt tcaatttaata 4380
ttaattttct acaataaatt ttagtgtcat ttccatttgg ggattattgt atacagcac 4440
atattttctg ttggaaaaca cactgttgtt tagttaagtt ttaaataggt gtattaccca 4500
agaagtaaaag atggaacgt t 4521

```

<210> 39

<211> 790

<212> PRT

<213> Homo sapiens

<400> 39

```

Met Arg Thr Tyr Arg Tyr Phe Leu Leu Leu Phe Trp Val Gly Gln Pro
1 5 10 15
Tyr Pro Thr Leu Ser Thr Pro Leu Ser Lys Arg Thr Ser Gly Phe Pro
20 25 30
Ala Lys Lys Arg Ala Leu Glu Leu Ser Gly Asn Ser Lys Asn Glu Leu
35 40 45
Asn Arg Ser Lys Arg Ser Trp Met Trp Asn Gln Phe Phe Leu Leu Glu
50 55 60
Glu Tyr Thr Gly Ser Asp Tyr Gln Tyr Val Gly Lys Leu His Ser Asp
65 70 75 80
Gln Asp Arg Gly Asp Gly Ser Leu Lys Tyr Ile Leu Ser Gly Asp Gly
85 90 95
Ala Gly Asp Leu Phe Ile Ile Asn Glu Asn Thr Gly Asp Ile Gln Ala
100 105 110
Thr Lys Arg Leu Asp Arg Glu Glu Lys Pro Val Tyr Ile Leu Arg Ala
115 120 125
Gln Ala Ile Asn Arg Arg Thr Gly Arg Pro Val Glu Pro Glu Ser Glu
130 135 140
Phe Ile Ile Lys Ile His Asp Ile Asn Asp Asn Glu Pro Ile Phe Thr
145 150 155 160
Lys Glu Val Tyr Thr Ala Thr Val Pro Glu Met Ser Asp Val Gly Thr
165 170 175
Phe Val Val Gln Val Thr Ala Thr Asp Ala Asp Asp Pro Thr Tyr Gly
180 185 190
Asn Ser Ala Lys Val Val Tyr Ser Ile Leu Gln Gly Gln Pro Tyr Phe
195 200 205

```

Ser Val Glu Ser Glu Thr Gly Ile Ile Lys Thr Ala Leu Leu Asn Met
 210 215 220
 Asp Arg Glu Asn Arg Glu Gln Tyr Gln Val Val Ile Gln Ala Lys Asp
 225 230 235 240
 Met Gly Gly Gln Met Gly Gly Leu Ser Gly Thr Thr Thr Val Asn Ile
 245 250 255
 Thr Leu Thr Asp Val Asn Asp Asn Pro Pro Arg Phe Pro Gln Ser Thr
 260 265 270
 Tyr Gln Phe Lys Thr Pro Glu Ser Ser Pro Pro Gly Thr Pro Ile Gly
 275 280 285
 Arg Ile Lys Ala Ser Asp Ala Asp Val Gly Glu Asn Ala Glu Ile Glu
 290 295 300
 Tyr Ser Ile Thr Asp Gly Glu Gly Leu Asp Met Phe Asp Val Ile Thr
 305 310 315 320
 Asp Gln Glu Thr Gln Glu Gly Ile Ile Thr Val Lys Lys Leu Leu Asp
 325 330 335
 Phe Glu Lys Lys Lys Val Tyr Thr Leu Lys Val Glu Ala Ser Asn Pro
 340 345 350
 Tyr Val Glu Pro Arg Phe Leu Tyr Leu Gly Pro Phe Lys Asp Ser Ala
 355 360 365
 Thr Val Arg Ile Val Val Glu Asp Val Asp Glu Pro Pro Val Phe Ser
 370 375 380
 Lys Leu Ala Tyr Ile Leu Gln Ile Arg Glu Asp Ala Gln Ile Asn Thr
 385 390 395 400
 Thr Ile Gly Ser Val Thr Ala Gln Asp Pro Asp Ala Ala Arg Asn Pro
 405 410 415
 Val Lys Tyr Ser Val Asp Arg His Thr Asp Met Asp Arg Ile Phe Asn
 420 425 430
 Ile Asp Ser Gly Asn Gly Ser Ile Phe Thr Ser Lys Leu Leu Asp Arg
 435 440 445
 Glu Thr Leu Leu Trp His Asn Ile Thr Val Ile Ala Thr Glu Ile Asn
 450 455 460
 Asn Pro Lys Gln Ser Ser Arg Val Pro Leu Tyr Ile Lys Val Leu Asp
 465 470 475 480
 Val Asn Asp Asn Ala Pro Glu Phe Ala Glu Phe Tyr Glu Thr Phe Val
 485 490 495
 Cys Glu Lys Ala Lys Ala Asp Gln Leu Ile Gln Thr Leu His Ala Val
 500 505 510
 Asp Lys Asp Asp Pro Tyr Ser Gly His Gln Phe Ser Phe Ser Leu Ala
 515 520 525
 Pro Glu Ala Ala Ser Gly Ser Asn Phe Thr Ile Gln Asp Asn Lys Asp
 530 535 540
 Asn Thr Ala Gly Ile Leu Thr Arg Lys Asn Gly Tyr Asn Arg His Glu
 545 550 555 560
 Met Ser Thr Tyr Leu Leu Pro Val Val Ile Ser Asp Asn Asp Tyr Pro
 565 570 575
 Val Gln Ser Ser Thr Gly Thr Val Thr Val Arg Val Cys Ala Cys Asp
 580 585 590
 His His Gly Asn Met Gln Ser Cys His Ala Glu Ala Leu Ile His Pro
 595 600 605
 Thr Gly Leu Ser Thr Gly Ala Leu Val Ala Ile Leu Leu Cys Ile Val
 610 615 620
 Ile Leu Leu Val Thr Val Val Leu Phe Ala Ala Leu Arg Arg Gln Arg
 625 630 635 640
 Lys Lys Glu Pro Leu Ile Ile Ser Lys Glu Asp Ile Arg Asp Asn Ile
 645 650 655
 Val Ser Tyr Asn Asp Glu Gly Gly Gly Glu Glu Asp Thr Gln Ala Phe
 660 665 670

```

Asp Ile Gly Thr Leu Arg Asn Pro Glu Ala Ile Glu Asp Asn Lys Leu
    675                      680                      685
Arg Arg Asp Ile Val Pro Glu Ala Leu Phe Leu Pro Arg Arg Thr Pro
    690                      695                      700
Thr Ala Arg Asp Asn Thr Asp Val Arg Asp Phe Ile Asn Gln Arg Leu
    705                      710                      715                      720
Lys Glu Asn Asp Thr Asp Pro Thr Ala Pro Pro Tyr Asp Ser Leu Ala
    725                      730                      735
Thr Tyr Ala Tyr Glu Gly Thr Gly Ser Val Ala Asp Ser Leu Ser Ser
    740                      745                      750
Leu Glu Ser Val Thr Thr Asp Ala Asp Gln Asp Tyr Asp Tyr Leu Ser
    755                      760                      765
Asp Trp Gly Pro Arg Phe Lys Lys Leu Ala Asp Met Tyr Gly Gly Val
    770                      775                      780
Asp Ser Asp Lys Asp Ser
    785                      790

```

```

<210> 40
<211> 987
<212> DNA
<213> Homo sapiens

```

```

<400> 40
cggagagggg gagaacagac aacgggcggc ggggagcagc atggagccgg cggcggggag 60
cagcatggag ccttcggctg actggctggc caaggccggc gcccggggtc gggtagagga 120
ggcgccggcg ctgctggagg cgggggcgct gcccaacgca ccgaatagtt acgctcggag 180
ggcgatccag gtcgatgatg tgggcagcgc ccgagtggcg gacgtcctgc tgctccacgg 240
cgccggagccc aactgcgccg accccgccac tctcaccgga cccgtgcacg acgctgcccg 300
ggagggcttc ctggacacgc tgggtggtgt gcaccgggcc gggcgccggc tggacgtgcg 360
cgatgcctgg ggccgtctgc ccgtggacct ggctgaggag ctggggccatc gcgatgtcgc 420
acggtacctg cgcgcggctg cggggggcgc cagaggcagt aaccatgccc gcatagatgc 480
cgcggaaggt cctcagaca tcgccgattg aaagaaccag agaggctctg agaaacctcg 540
ggaaacttag atcatcagtc accgaaggtc ctacagggcc acaactgccc ccgccacaac 600
ccaccgccct ttgctagttt tcatttagaa aatagagctt ttaaaaatgt cctgcctttt 660
aacgtagata taagccttcc ccactaccg taaatgtcca ttatatcat ttttatata 720
ttctataaaa aatgtaaaaa agaaaaaacac cgcttctgcc ttttcaactgt gttggagttt 780
ctctgagtga gcactcaccg cctaagcgca cattcatgtg ggcatttctt gcgagcctcg 840
cagctccggg aagctgtcga ctcatgaca agcattttgt gaactaggga agctcagggg 900
ggttactggc ttctcttgag tcacactgct agcaaatggc agaaccaaag ctcaaaataa 960
aataaaataa ttttctattc ttcactc

```

```

<210> 41
<211> 156
<212> PRT
<213> Homo sapiens

```

```

<400> 41
Met Glu Pro Ala Ala Gly Ser Ser Met Glu Pro Ser Ala Asp Trp Leu
    1           5           10
Ala Thr Ala Ala Ala Arg Gly Arg Val Glu Glu Val Arg Ala Leu Leu
    20           25           30
Glu Ala Gly Ala Leu Pro Asn Ala Pro Asn Ser Tyr Gly Arg Arg Pro
    35           40           45
Ile Gln Val Met Met Met Gly Ser Ala Arg Val Ala Glu Leu Leu Leu
    50           55           60
Leu His Gly Ala Glu Pro Asn Cys Ala Asp Pro Ala Thr Leu Thr Arg
    65           70           75           80

```

Pro Val His Asp Ala Ala Arg Glu Gly Phe Leu Asp Thr Leu Val Val
 85 90 95
 Leu His Arg Ala Gly Ala Arg Leu Asp Val Arg Asp Ala Trp Gly Arg
 100 105 110
 Leu Pro Val Asp Leu Ala Glu Glu Leu Gly His Arg Asp Val Ala Arg
 115 120 125
 Tyr Leu Arg Ala Ala Ala Gly Gly Thr Arg Gly Ser Asn His Ala Arg
 130 135 140
 Ile Asp Ala Ala Glu Gly Pro Ser Asp Ile Pro Asp
 145 150 155

<210>: 42
 <211> 5142
 <212> DNA
 <213> Homo sapiens

<400> 42
 gaattcggcc gagaggacga gggggaggcc cagagctgct ogtctgtctt tgcccagacc 60
 cgagcccgag ccagagcccg agcccgagcc cgagcccgag cccgaacgca agcctgggag 120
 ccgggagccc ggctagggac tccctcatt tatggagcag gcacccaaca tggctgagcc 180
 ccggggcccc gtgacacatg gagtccagat tgccttcac acagagccca tgggtgggac 240
 agagatgggc actctacgtc gaggtggacg acgcccagct aaggtgcaa gagccagtac 300
 ctacgggggtt gctgtcgctg tgcagggaaat cgtctggcag cctttgtgg tgctcaacag 360
 tggggagaaaa ggcggtgact cctttggggt ccaaatcaag ggggccaatg accaaggggc 420
 ctcaaggagct ctgagctcag atttggaaat ccttgagaac cctactctc aggtcaaggg 480
 atttcctgcc cctcgcgaga gcagccacatc tgatgaggag cctggggcct actggaatgg 540
 aaagctactc cgttccactc cccaggcctc actggcagcc cctggccaga tggatcctaag 600
 taacagaagc aacagcatgc tggagctagc cccgaaagt gcttccccag gtagcaccat 660
 tgacactgct cccctgtctt cagtggactc actcatcaac aagtttgaca gtcaacttgg 720
 aggcacggcc cggggtcgga ctggcccgcc aacacggatg ctaccccctg aacagcgcaa 780
 acgggacgaag agcctggaca gccgcctccc acgggacacc tttgaggaa cggagcgcca 840
 gtccaccaac cactggacct ctgacacaaa atatgacaa cagtggctt agccgtlctc gtcagactca 960
 gccagccagc agccagaacc ttgaggagcc gccggaggat gcacaggacc ccacctgctt 1020
 ggaactgggtc cttaagagtt ttgaggagcc agaccagcag gaggcagccc caccaggcag 1080
 gcagttcaaa tcaactccag acctccttcg actcctgag gagggaagct caaagaatga 1140
 tgtgcaccat atgaaggcca ccatctatgg cctctgagc gctggagaag atgcagcctc tagtgatggt 1200
 aacctctgtg agggagaagg ttagtttgggt gctggagaag cttaccocaa aggaagtgag 1260
 ttcttcttgt tctaactaagg ccgtggcagg gaagggtgag aagctagagc catccaagt 1320
 gctacacgca aagctggatg aagaggtgaa gaagcggcag aagcagactc aggagctgct 1380
 tgggctggag cggcagctgg agggaaaaac agaagatgct agccagactc aggagctgct 1440
 ggagaggagg aagggggagg ccagacagag caacaaggag ctacagaaca tgaacggcct 1500
 ctgggaccaa ggtgaagatt tacgacatgg gctggagacc caggtgatgg agctgcagaa 1560
 caagctgaaa catgtccagg gtccctgagcc tgctaaggag gcttactga aggacctgtt 1620
 agagaccagg gaactctcgg aagaggtctt ggaggggaaa cagcagtagt aggcagcact 1680
 gaggctcgcg gagcggaggat tgacagccct gaagggggcc ctgaagagag aggtagcctc 1740
 ccgtgaccag gaggtggaac atgtccggca gcagtaaccg cgagacacag agcagctccg 1800
 caggagcatg caagatgcga cccaggacca tgcagtgctg gaggccgaga gacagaagt 1860
 gtacagccct gtgcgagggc tgcagaggga gctggaggag actccaagg agacaggtca 1920
 ttggcagagt atgtccaga agaacaagga ggatctttag gccacagagc aggaactcct 1980
 gcagctcgga atggagaagg agggagatgga agaggagctt ggagagaaga tagaggtctt 2040
 ccagaggaaa tttagagcag ccagagctag tgcctggagat actcggcagg ttagaggtct 2100
 caagaaggag cgtctccgga cacaggagga gcttaaggaa cctcaggcag aacggcagag 2160
 ccagagggtg gctggcgac accgggaccg gaagtggag aagcagctgg cgtctcag 2220
 ggcagggct gatcgaggtc gggagctgga ctcaagatca aaaaagacct 2280
 ccagcaactg cgcagaggat gtgaagaggc ttccaaggct aagatggtgg ccgaggcaga 2340
 ggcaacagtg ctggggcgag ggcgggcgcg agtggagacg acgcttcggg agaccacaga 2400
 ggaaaaatgac gaattccgcc ggcgcactct ggtttggag cagcagctga aggagactcg 2460

```

aggctcgtgtg gatgggtggg aagcgggtgga ggcacgacta cgggacaagc tgcagcggct 2460
ggagggcagag aaacacgacg cctggaatgcg tccacggaag aggagggggag 2520
tctggcgacga gccaaagcggg cactggaggcg acgcctagag gaggctcagc gggggctggc 2580
ccgcctgggg ccaggagcagc agacactgaa ccggcccctg gaggaggaag ggaacgagcg 2640
ggaggtgtctc cggcgaggca aggcctgagct ggaggagcag aagcgtttcg tggacaggac 2700
tgtggaccga ctgaacaagg agttggagaa gatcggggag gactctaaag aagccctgca 2760
gcagctccag gccacgctgg aggtattataa ggaaaaggcc cggcggggagg tggcagatgc 2820
ccagcgccag gccaaaggatt gggccagtga ggctgagaag acctctggag gactgagccg 2880
acctcaggat gagatccaga ggctgcggca gccctctgag gcatcccgag ctgagcgagcg 2940
cacagccggg ctggacaaaag agctactggc ccagcgactg cagggggctgc agcaagaggc 3000
agagaacaag aagcgttccc agttagatga ggagaagaac accgtggagc tgctaacaga 3120
tgcgttgaat cgtggccggg accaggttga tcagctgagg acagagctca tgcaggaagag 3180
gtctgtccag caggacctgg agtgtgacaa aatctccttg gagagacaga acaaggacct 3240
gaagacccgg ttggccagct cagaaggctt ccagaagcct agtgccagcc ttctcagct 3300
tgagtccagc aatcagttgt tgcaggagcg gctacaggct gaagagaggg aagaacacagt 3360
tctgcagctc accaatcgaa aactggagcg gaaagttaaa gaactatcca tccagattga 3420
agaagagcgg cagcatgtca atgaccagaa agaccagcta agcctgaggg tgaaggcttt 3480
gaagcgtcag gtgatgaag cagaagagga aattgagcga ctggcagggc taggaagaaga 3540
ggccacagct gagggtggagg agcagcatga ggtcaatgaa cagctccagg ccgggactca 3600
gtctctggag aaggactcct ggcgcaaaag ttcccgctca gctgctgagt cagctctcaa 3660
aaacgaaggc ctgagctcag atgaggaatt gcacagtgtc tacgatccct cgtccattgc 3720
atcactgtct accggagagca acctacagac cagctcctgt tagctcgttg tctcaagga 3780
ctcagaaaac cgttgcaggg cctatcccag caagtctgtc tctgtctctg ccacctggg 3840
ttctgtcatc ctatgggtga cccaattatt cagacctaa acaggggagg tgcagagtga 3900
tgtgtcataa aaaaaaaa tctatcgcaa taagtactga tatggacttt ccaactgagg 3960
agtggacgtt tcaagccaaac tgagcccttt cctcaagtgc cgacacotcc ctactctctc 4020
ttatagtga aggatgttca gctattaggct gatggggagt gagaaggata ggaagggata 4080
gaaattgcc a tgtgtataaa gctttattct tttagcctta accctaaggc tcaaggaaat 4140
acctatggt attgtgctcc ctggattcct gcaactcatt tctctccac tctggagcag 4200
ggcgagggga atgttatggg taacagacat gcaaggcatg tcttcccat ttctttgac 4260
aagtatgggg cccatgtggt agtccocata ccctocagt tcttatattt ttgtctctt 4320
cctttccct cctaccttgc atttttctgt tcttctctgt tcaagtgcct agccaaggca 4380
aggagataag gatgctcttc ttgctttita tatctgcaca ttcatacct tccaaagacc 4440
agcttttccc cagccagggc cctcagcctt cctgtctgcc ccagtgattg attgagagag 4500
ctgttgggtt ttctctgcca atgaccctg ggagagggac tttgttaggg tcatgataaa 4560
gtggcggggg tcttggtcct gctcagggtt ttcatctctc ctctctccc tccctgtgta 4620
ctgttgatat ggttataaagg tggttgcacc tgggagccct gacaactggc tgcacaaatt 4680
ccaaaagtaa aggtgtcagt ccctgtggcc cctctgggg ctctctgac ccatgtgccc 4740
caaactcaat aagagaacca agggaccttc attttctgag gtgcttggc ctgactcagg 4800
tctgttcaag ggggtagaag ctgactgtaa aaatgggaag aggcaacgga agacatttat 4860
ttctcttttg aagtttgggg agaaccaagc cctggtaggg aagaggttaa ggggatgatt 4920
cacctccata ttctctaagg aggttgtata gggagccggt ggcaggagga aggcgtttt 4980
cacaaatgac ttgtaatgtc gtattataaa aaattcctat attctctgc aaatacaaac 5040
ttcttcccca atccaatcca gcttgggttt tattttaaat taaattatta aattacacat 5100
ttatattgaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aa 5142

```

<210> 43

<211> 1203

<212> PRT

<213> Homo sapiens

<400> 43

```

Met Glu Gln Ala Pro Asn Met Ala Glu Pro Arg Gly Pro Val Asp His
1           5           10           15
Gly Val Gln Ile Arg Phe Ile Thr Glu Pro Val Ser Gly Ala Glu Met
20           25           30
Gly Thr Leu Arg Arg Gly Gly Arg Arg Pro Ala Lys Asp Ala Arg Ala

```


	35				40				45						
Ser	Thr	Tyr	Gly	Val	Ala	Val	Arg	Val	Gln	Gly	Ile	Ala	Gly	Gln	Pro
	50					55					60				
Phe	Val	Val	Leu	Asn	Ser	Gly	Glu	Lys	Gly	Gly	Asp	Ser	Phe	Gly	Val
	65					70				75					80
Gln	Ile	Lys	Gly	Ala	Asn	Asp	Gln	Gly	Ala	Ser	Gly	Ala	Leu	Ser	Ser
				85					90						95
Asp	Leu	Glu	Leu	Pro	Glu	Asn	Pro	Tyr	Ser	Gln	Val	Lys	Gly	Phe	Pro
				100					105						110
Ala	Pro	Ser	Gln	Ser	Ser	Thr	Ser	Asp	Glu	Glu	Pro	Gly	Ala	Tyr	Trp
				115					120						125
Asn	Gly	Lys	Leu	Leu	Arg	Ser	His	Ser	Gln	Ala	Ser	Leu	Ala	Gly	Pro
				130			135					140			
Gly	Pro	Val	Asp	Pro	Ser	Asn	Arg	Ser	Asn	Ser	Met	Leu	Glu	Leu	Ala
				145			150								160
Pro	Lys	Val	Ala	Ser	Pro	Gly	Ser	Thr	Ile	Asp	Thr	Ala	Pro	Leu	Ser
				165						170					175
Ser	Val	Asp	Ser	Leu	Ile	Asn	Lys	Phe	Asp	Ser	Gln	Leu	Gly	Gly	Gln
				180					185						190
Ala	Arg	Gly	Arg	Thr	Gly	Arg	Arg	Thr	Arg	Met	Leu	Pro	Pro	Glu	Gln
				195					200						205
Arg	Lys	Arg	Ser	Lys	Ser	Leu	Asp	Ser	Arg	Leu	Pro	Arg	Asp	Thr	Phe
				210			215								220
Glu	Glu	Arg	Glu	Arg	Gln	Ser	Thr	Asn	His	Trp	Thr	Ser	Ser	Thr	Lys
				225			230				235				240
Tyr	Asp	Asn	His	Val	Gly	Thr	Ser	Lys	Gln	Pro	Ala	Gln	Ser	Gln	Asn
				245					250						255
Leu	Ser	Pro	Leu	Ser	Gly	Phe	Ser	Arg	Ser	Arg	Gln	Thr	Gln	Asp	Trp
				260					265						270
Val	Leu	Gln	Ser	Phe	Glu	Glu	Pro	Arg	Arg	Ser	Ala	Gln	Asp	Pro	Thr
				275					280						285
Met	Leu	Gln	Phe	Lys	Ser	Thr	Pro	Asp	Leu	Leu	Arg	Asp	Gln	Gln	Glu
				290			295					300			
Ala	Ala	Pro	Pro	Gly	Ser	Val	Asp	His	Met	Lys	Ala	Thr	Ile	Tyr	Gly
				305			310					315			320
Ile	Leu	Arg	Glu	Gly	Ser	Ser	Glu	Ser	Glu	Thr	Ser	Val	Arg	Arg	Lys
				325					330						335
Val	Ser	Leu	Val	Leu	Glu	Lys	Met	Gln	Pro	Leu	Val	Met	Val	Ser	Ser
				340					345						350
Gly	Ser	Thr	Lys	Ala	Val	Ala	Gly	Gln	Gly	Glu	Leu	Thr	Arg	Lys	Val
				355			360								365
Glu	Glu	Leu	Gln	Arg	Lys	Leu	Asp	Glu	Glu	Val	Lys	Lys	Arg	Gln	Lys
				370			375					380			
Leu	Glu	Pro	Ser	Gln	Val	Gly	Leu	Glu	Arg	Gln	Leu	Glu	Glu	Lys	Thr
				385			390					395			400
Glu	Glu	Cys	Ser	Arg	Leu	Gln	Glu	Leu	Leu	Glu	Arg	Arg	Lys	Gly	Glu
				405					410						415
Ala	Gln	Gln	Ser	Asn	Lys	Glu	Leu	Gln	Asn	Met	Lys	Arg	Leu	Leu	Asp
				420					425						430
Gln	Gly	Glu	Asp	Leu	Arg	His	Gly	Leu	Glu	Thr	Gln	Val	Met	Glu	Leu
				435			440								445
Gln	Asn	Lys	Leu	Lys	His	Val	Gln	Gly	Pro	Glu	Pro	Ala	Lys	Glu	Val
				450			455								460
Leu	Leu	Lys	Asp	Leu	Leu	Glu	Thr	Arg	Glu	Leu	Leu	Glu	Glu	Val	Leu
				465			470					475			480
Glu	Gly	Lys	Gln	Arg	Val	Glu	Glu	Gln	Leu	Arg	Leu	Arg	Glu	Arg	Glu
				485					490						495
Leu	Thr	Ala	Leu	Lys	Gly	Ala	Leu	Lys	Glu	Glu	Val	Ala	Ser	Arg	Asp

										500						505						510			
Gln	Glu	Val	Glu	His	Val	Arg	Gln	Gln	Tyr	Gln	Arg	Asp	Thr	Glu	Gln										
										515			520			525									
Leu	Arg	Arg	Ser	Met	Gln	Asp	Ala	Thr	Gln	Asp	His	Ala	Val	Leu	Glu										
										530			535			540									
Ala	Glu	Arg	Gln	Lys	Met	Ser	Ala	Leu	Val	Arg	Gly	Leu	Gln	Arg	Glu										
										545			550			555									
Leu	Glu	Glu	Thr	Ser	Glu	Glu	Thr	Gly	His	Trp	Gln	Ser	Met	Phe	Gln										
										565			570			575									
Lys	Asn	Lys	Glu	Asp	Leu	Arg	Ala	Thr	Lys	Gln	Glu	Leu	Gln	Leu											
										580			585			590									
Arg	Met	Glu	Lys	Glu	Glu	Met	Glu	Glu	Leu	Gly	Glu	Lys	Ile	Glu											
										595			600			605									
Val	Leu	Gln	Arg	Glu	Leu	Glu	Gln	Ala	Arg	Ala	Ser	Ala	Gly	Asp	Thr										
										610			615			620									
Arg	Gln	Val	Glu	Val	Leu	Lys	Lys	Glu	Leu	Leu	Arg	Thr	Gln	Glu	Glu										
										625			630			635									
Leu	Lys	Glu	Leu	Gln	Ala	Glu	Arg	Gln	Ser	Gln	Glu	Val	Ala	Gly	Arg										
										645			650			655									
His	Arg	Asp	Arg	Glu	Leu	Glu	Lys	Gln	Leu	Ala	Val	Leu	Arg	Val	Glu										
										660			665			670									
Ala	Asp	Arg	Gly	Arg	Glu	Leu	Glu	Gln	Asn	Leu	Gln	Leu	Gln	Lys											
										675			680			685									
Thr	Leu	Gln	Gln	Leu	Arg	Gln	Asp	Cys	Glu	Glu	Ala	Ser	Lys	Ala	Lys										
										690			695			700									
Met	Val	Ala	Glu	Ala	Glu	Ala	Thr	Val	Leu	Gly	Gln	Arg	Arg	Ala	Ala										
										705			710			715									
Val	Glu	Thr	Thr	Leu	Arg	Glu	Thr	Gln	Glu	Glu	Asn	Asp	Glu	Phe	Arg										
										725			730			735									
Arg	Arg	Ile	Leu	Gly	Leu	Glu	Gln	Gln	Leu	Lys	Glu	Thr	Arg	Gly	Leu										
										740			745			750									
Val	Asp	Gly	Gly	Glu	Ala	Val	Glu	Ala	Arg	Leu	Arg	Asp	Lys	Leu	Gln										
										755			760			765									
Arg	Leu	Glu	Ala	Glu	Lys	Gln	Gln	Leu	Glu	Glu	Ala	Leu	Asn	Ala	Ser										
										770			775			780									
Gln	Glu	Glu	Glu	Gly	Ser	Leu	Ala	Ala	Ala	Lys	Arg	Ala	Leu	Glu	Ala										
										785			790			795									
Arg	Leu	Glu	Glu	Ala	Gln	Arg	Gly	Leu	Ala	Arg	Leu	Gly	Gln	Glu	Gln										
										805			810			815									
Gln	Thr	Leu	Asn	Arg	Ala	Leu	Glu	Glu	Glu	Gly	Lys	Gln	Arg	Glu	Val										
										820			825			830									
Leu	Arg	Arg	Gly	Lys	Ala	Glu	Leu	Glu	Gln	Lys	Arg	Leu	Leu	Asp											
										835			840			845									
Arg	Thr	Val	Asp	Arg	Leu	Asn	Lys	Glu	Leu	Glu	Lys	Ile	Gly	Glu	Asp										
										850			855			860									
Ser	Lys	Gln	Ala	Leu	Gln	Gln	Leu	Gln	Ala	Gln	Leu	Glu	Asp	Tyr	Lys										
										865			870			875									
Glu	Lys	Ala	Arg	Arg	Glu	Val	Ala	Asp	Ala	Gln	Arg	Gln	Ala	Lys	Asp										
										885			890			895									
Trp	Ala	Ser	Glu	Ala	Glu	Lys	Thr	Ser	Gly	Gly	Leu	Ser	Arg	Leu	Gln										
										900			905			910									
Asp	Glu	Ile	Gln	Arg	Leu	Arg	Gln	Ala	Leu	Gln	Ala	Ser	Gln	Ala	Glu										
										915			920			925									
Arg	Asp	Thr	Ala	Arg	Leu	Asp	Lys	Glu	Leu	Leu	Ala	Gln	Arg	Leu	Gln										
										930			935			940									
Gly	Leu	Glu	Gln	Glu	Ala	Glu	Asn	Lys	Lys	Arg	Ser	Gln	Asp	Asp	Arg										
										945			950			955									
Ala	Arg	Gln	Leu	Lys	Gly	Leu	Glu	Glu	Lys	Val	Ser	Arg	Leu	Glu	Thr										

										965											970											975
Glu	Leu	Asp	Glu	Glu	Lys	Asn	Thr	Val	Leu	Leu	Thr	Asp	Arg	Val											Glu	Leu	Thr	Asp	Arg	Val		
										980											985											990
Asn	Arg	Gly	Arg	Asp	Gln	Val	Asp	Gln	Leu	Arg	Thr	Glu	Leu	Met	Gln											Glu	Leu	Met	Gln			
										995											1000											1005
Glu	Arg	Ser	Ala	Arg	Gln	Asp	Leu	Glu	Cys	Asp	Lys	Ile	Ser	Leu	Glu											Glu	Leu	Ser	Leu	Glu		
										1010											1015											1020
Arg	Gln	Asn	Lys	Asp	Leu	Lys	Thr	Arg	Leu	Ala	Ser	Ser	Glu	Gly	Phe											Glu	Leu	Ser	Leu	Glu		
										1025											1030											1035
Gln	Lys	Pro	Ser	Ala	Ser	Leu	Ser	Gln	Leu	Glu	Ser	Gln	Asn	Gln	Leu											Glu	Leu	Ser	Leu	Glu		
										1045											1050											1055
Leu	Gln	Glu	Arg	Leu	Gln	Ala	Glu	Glu	Arg	Glu	Lys	Thr	Val	Leu	Gln											Glu	Leu	Ser	Leu	Glu		
										1060											1065											1070
Ser	Thr	Asn	Arg	Lys	Leu	Glu	Arg	Lys	Val	Lys	Glu	Leu	Ser	Ile	Gln											Glu	Leu	Ser	Leu	Glu		
										1075											1080											1085
Ile	Glu	Asp	Glu	Arg	Gln	His	Val	Asn	Asp	Gln	Lys	Asp	Gln	Leu	Ser											Glu	Leu	Ser	Leu	Glu		
										1090											1095											1100
Leu	Arg	Val	Lys	Ala	Leu	Lys	Arg	Gln	Val	Asp	Glu	Ala	Glu	Glu	Glu											Glu	Leu	Ser	Leu	Glu		
										1105											1110											1115
Ile	Glu	Arg	Leu	Asp	Gly	Leu	Arg	Lys	Lys	Ala	Gln	Arg	Glu	Val	Glu											Glu	Leu	Ser	Leu	Glu		
										1125											1130											1135
Glu	Gln	His	Glu	Val	Asn	Glu	Gln	Leu	Gln	Ala	Arg	Ile	Lys	Ser	Leu											Glu	Leu	Ser	Leu	Glu		
										1140											1145											1150
Glu	Lys	Asp	Ser	Trp	Arg	Lys	Ala	Ser	Arg	Ser	Ala	Ala	Glu	Ser	Ala											Glu	Leu	Ser	Leu	Glu		
										1155											1160											1165
Leu	Lys	Asn	Glu	Gly	Leu	Ser	Ser	Asp	Glu	Glu	Phe	Asp	Ser	Val	Tyr											Glu	Leu	Ser	Leu	Glu		
										1170											1175											1180
Asp	Pro	Ser	Ser	Ile	Ala	Ser	Leu	Leu	Thr	Glu	Ser	Asn	Leu	Gln	Thr											Glu	Leu	Ser	Leu	Glu		
										1185											1190											1195
Ser	Ser	Cys																							1200							

<210> 44

<211> 1925

<212> DNA

<213> Homo sapiens

<400> 44

```

agtggagtgg gacaggtata taaaggaagt acaggggctg gggaagaggc cctgtctagg 60
tagctggcac caggagccgt gggcaaggga agaggccaca ccctgcccctg ctctgctgca 120
gccagaatgg ggtgtaaggc gtctcaaaaca ggctttgtgg tcttggtgct gctccagtgc 180
tgtctgtcat acaaaactggt ctgctactac accagctggt cccagtaccg ggaaggcgat 240
gggagctgct tccagatgac ccttgaccgc ttctctgtga cccacatcat ctacagcttt 300
gccaatataa gcaacgatca catcgacacc tgggagtggg atgatgtgac gctctacggc 360
atgctcaaca cactcaagaa caggaaacccc aacctgaaga ctctctgtgc tgcggaggga 420
tggaaacttt ggtctcaaaag atttccaag atagcctcca acaccagag tcgcgcggat 480
ttcatcaagt cagtaaccgc attcctgcgc acccatggct ttgatgggct ggaacctggc 540
tggctctacc ctggacggag agacaaacag cattttacca cctaatacaa ggaatgaag 600
gccgaattta taaaggaagc ccagccaggg aaaaagcagc tctctgctag cgcagcactg 660
ctcgggggga aggtcaccat tgacagcagc tatgacattg ccaagatact ccaacacctg 720
gatttcatta gcatcatgac ctacgatttt catggagcct ggcgtgggag cacagggcat 780
cacagtcccc tgttccaggg tcaggaggat gcaagtcctg acagattcag caacactgac 840
tatgctgtgg ggtacatggt gaggctgggg gctcctgcca gtaagctggt gatgggcatc 900
ccacacctgg ggaggagcgt cactctggct tottctgaga ctggtgttgg agcccaatc 960
tcaggaccgg gaattccagg ccggttcacc aaggaggcag ggacccttgc ctactatgat 1020
atctgtgact tctctccggg agccacagtc catgaaaccc tcggccagca ggtcccatat 1080
gccaccaagg gcaaccagtg gtaggagatc gacgaccagg aaagcgtcaa aagcaaggtg 1140

```

```

cagtacctga aggataggca gctggcaggc gccatggtat gggccctgga cctggatgac 1200
ttccagggtt ccttctgagg ccaggatctg cgcttccctc tcaccaatgc catcaaggat 1260
gcactcgtctg caacgtagacc ctctgttctg cacacagcac gggggccaag gatgcccctg 1320
ccccctctgg ctccagctgg ccgggagcct gatcacctgc cctgctgagt ccaggctga 1380
gcctcagttct cctccctctg gggcctatgc agaggtccac aacacacaga tttagctca 1440
gccttggtgg gcagagaggt agggatgggg ctgtggggat agtgagcat cgcaatgtaa 1500
gactcgggat tagtacacac ttgttgatga ttaatggaaa tgtttacaga tcccacagcc 1560
tggaaggga atttcttcaa ctccctgccc cctagccctc cttatcaag gagaccattt 1620
tggaagcttc tatccacaag gagccaaaca tctacaaga cacagtgaac atactaatta 1680
taccctctgc aaagccagct tgaacacctc acttaggaac gtaatcgtgt cccctatcct 1740
acttcccctt cctaattcca cagctgctca ataaagtaca agagttaa ac agtgtgttg 1800
cgctttgctt tggtctatct ttgagcgccc actagaccac ctggactcac ctccccatc 1860
tctctgggtt tccttctctt gagccttggg acccctgagc ttgcagagat gaagccgccc 1920
atgtt

```

<210> 45

<211> 383

<212> PRT

<213> Homo sapiens

<400> 45

```

Met Gly Val Lys Ala Ser Gln Thr Gly Phe Val Val Leu Val Leu Leu
1 5 10 15
Gln Cys Cys Ser Ala Tyr Lys Leu Val Cys Tyr Tyr Thr Ser Thr Ser
20 25 30
Gln Tyr Arg Glu Gly Asp Gly Ser Cys Phe Pro Asp Ala Leu Asp Arg
35 40 45
Phe Leu Cys Thr His Ile Ile Tyr Ser Phe Ala Asn Ile Ser Asn Asp
50 55 60
His Ile Asp Thr Trp Glu Trp Asn Asp Val Thr Leu Tyr Gly Met Leu
65 70 75 80
Asn Thr Leu Lys Asn Arg Asn Pro Asn Leu Lys Thr Leu Leu Ser Val
85 90 95
Gly Gly Trp Asn Phe Gly Ser Gln Arg Phe Ser Lys Ile Ala Ser Asn
100 105 110
Thr Gln Ser Arg Arg Thr Phe Ile Lys Ser Val Pro Pro Phe Leu Arg
115 120 125
Thr His Gly Phe Asp Gly Leu Asp Leu Ala Trp Leu Tyr Pro Gly Arg
130 135 140
Arg Asp Lys Gln His Phe Thr Thr Leu Ile Lys Glu Met Lys Ala Glu
145 150 155 160
Phe Ile Lys Glu Ala Gln Pro Gly Lys Lys Gln Leu Leu Leu Ser Ala
165 170 175
Ala Leu Ser Ala Gly Lys Val Thr Ile Asp Ser Ser Tyr Asp Ile Ala
180 185 190
Lys Ile Ser Gln His Leu Asp Phe Ile Ser Ile Met Thr Tyr Asp Phe
195 200 205
His Gly Ala Trp Arg Gly Thr Thr Gly His His Ser Pro Leu Phe Arg
210 215 220
Gly Gln Glu Asp Ala Ser Pro Asp Arg Phe Ser Asn Thr Asp Tyr Ala
225 230 235 240
Val Gly Tyr Met Leu Arg Leu Gly Ala Pro Ala Ser Lys Leu Val Met
245 250 255
Gly Ile Pro Thr Phe Gly Arg Ser Phe Thr Leu Ala Ser Ser Glu Thr
260 265 270
Gly Val Gly Ala Pro Ile Ser Gly Pro Gly Ile Pro Gly Arg Phe Thr
275 280 285
Lys Glu Ala Gly Thr Leu Ala Tyr Tyr Glu Ile Cys Asp Phe Leu Arg

```

290		295		300
Gly Ala Thr Val His Arg Thr Leu Gly Gln Gln Val Pro Tyr Ala Thr				
305		310		315
Lys Gly Asn Gln Trp Val Gly Tyr Asp Asp Gln Glu Ser Val Lys Ser				
		325		330
Lys Val Gln Tyr Leu Lys Asp Arg Gln Leu Ala Gly Ala Met Val Trp				
		340		345
Ala Leu Asp Leu Asp Asp Phe Gln Gly Ser Phe Cys Gly Gln Asp Leu				
		355		360
Arg Phe Pro Leu Thr Asn Ala Ile Lys Asp Ala Leu Ala Ala Thr				
		370		375
				380

<210> 46

<211> 1528

<212> DNA

<213> Homo sapiens

<400> 46

```

cgggtcccca ttccgggtcc agcctccaat ccgaccccca ttccgggtgc agcctcggac 60
ctagctccgg cccctcgggtct atccgggttc atccctccct cctgttccgg atcttatctt 120
cgcccaagcg cctactccag atcccgtagc cagacctcaa gccatgggtg gtcccttctc 180
cgtctcgtgc tccgcccgcc cgggactcag gctcctgggt ttggccggag cggggtctct 240
agccgctggg ttctcgtccc gaccggaacc tgtacgagct gccagtgaac gacggaggct 300
gtatcccccg agcgtcgtagt acccagacct ccgaaagcac acaactgca tggccagtca 360
cctgacccca cagctctatg caccggtctg acacaagacc acacccactg gttggacgt 420
agatcagtgt atccagactg cgttggacaa cctcggccac cccttcatca agactgtggg 480
catggtggtc ggagatgagg agacctatga ggtatttgtc gacctgttg accctgtgat 540
ccaagagcga cacaatggat atgacccccc gacaatgaag cacaccacgg atctagatgc 600
cagtataaat cgttctgggt actttgatga gagggtatga ttgtcctcta gactcagaac 660
tgccgaagc atccgaggac tcagtcgtcc tcacgcttgc actcgagcag agcgacgaga 720
ggtggaacgt gttgtggttg atgcactgag tggcctgaag ggtgacctg ctggacgtta 780
ctataggctc agtgagatga cagaggctga acagcagcag ctattgatg accactttct 840
gtttgataag cctgtgtccc cgttgcgtac tgcagcagga atggctcgag actggccaga 900
tgctcgttga atttggcaca acaatgagaa gagcttcctg atctgggtga atgaggagga 960
tcatacacgg gtgatctcca tggagaaggg ttgtaacatg aagagagtg ttgaaagatt 1020
ctgcogagcg ctcaaaaggg tggagagact tatccaagaa cgtggctggg agttcatgtg 1080
gaatgagcgt ttgggatata ctttgacctg tccatctaac ctgggcaact gacttcgggc 1140
aggagtgcac atcaaaactgc ccttgctaag caaagatagc cgcttcccaa agatccttga 1200
gaacctaaag acccaaaaac gtgtactcgg aggagtggag actcgtctga caggcggtgt 1260
ctttgatatt tctaatcttg accgactagg caaatcagag gtggagctgg tgcacttgt 1320
catcagtga gtaactatt tgattgattg tgaacggcgt ctggagagag gccaggatat 1380
cgcgactccc acacactgtc tccacaccaa gcattaactc cccatcgcca gctgatgact 1440
caagattccc aggagttttg ctcatctcaa tgatggccca ttctacttgc tctggacctg 1500
ccccgcctc cctgcctcc atccgtagt

```

<210> 47

<211> 417

<212> PRT

<213> Homo sapiens

<400> 47

Met Ala Gly Pro Phe Ser Arg Leu Leu Ser Ala Arg Pro Gly Leu Arg				
1	5	10	15	
Leu Leu Ala Leu Ala Gly Ala Gly Ser Leu Leu Ala Ala Gly Phe Leu Leu				
	20	25	30	
Arg Pro Glu Pro Val Arg Ala Ala Ser Glu Arg Arg Arg Leu Tyr Pro				
	35	40	45	

Pro Ser Ala Glu Tyr Pro Asp Leu Arg Lys His Asn Asn Cys Met Ala
 50 55 60
 Ser His Leu Thr Pro Ala Val Tyr Ala Arg Leu Cys Asp Lys Thr Thr
 65 70 75 80
 Pro Thr Gly Trp Thr Leu Asp Gln Cys Ile Gln Thr Gly Val Asp Asn
 85 90 95
 Pro Gly His Pro Phe Ile Lys Thr Val Gly Met Val Ala Gly Asp Glu
 100 105 110
 Glu Thr Tyr Glu Val Phe Ala Asp Leu Phe Asp Pro Val Ile Gln Glu
 115 120 125
 Arg His Asn Gly Tyr Asp Pro Arg Thr Met Lys His Thr Thr Asp Leu
 130 135 140
 Asp Ala Ser Lys Ile Arg Ser Gly Tyr Phe Asp Glu Arg Tyr Val Leu
 145 150 155 160
 Ser Ser Arg Val Arg Thr Gly Arg Ser Ile Arg Gly Leu Ser Leu Pro
 165 170 175
 Pro Ala Cys Thr Arg Ala Glu Arg Arg Glu Val Glu Arg Val Val Val
 180 185 190
 Asp Ala Leu Ser Gly Leu Lys Gly Asp Leu Ala Gly Arg Tyr Tyr Arg
 195 200 205
 Leu Ser Glu Met Thr Glu Ala Glu Gln Gln Gln Leu Ile Asp Asp His
 210 215 220
 Phe Leu Phe Asp Lys Pro Val Ser Pro Leu Leu Thr Ala Ala Gly Met
 225 230 235 240
 Ala Arg Asp Trp Pro Asp Ala Arg Gly Ile Trp His Asn Asn Glu Lys
 245 250 255
 Ser Phe Leu Ile Trp Val Asn Glu Glu Asp His Thr Arg Val Ile Ser
 260 265 270
 Met Glu Lys Gly Gly Asn Met Lys Arg Val Phe Glu Arg Phe Cys Arg
 275 280 285
 Gly Leu Lys Glu Val Glu Arg Leu Ile Gln Glu Arg Gly Trp Glu Phe
 290 295 300
 Met Trp Asn Glu Arg Leu Gly Tyr Ile Leu Thr Cys Pro Ser Asn Leu
 305 310 315 320
 Gly Thr Gly Leu Arg Ala Gly Val His Ile Lys Leu Pro Leu Leu Ser
 325 330 335
 Lys Asp Ser Arg Phe Pro Lys Ile Leu Glu Asn Leu Arg Leu Gln Lys
 340 345 350
 Arg Gly Thr Gly Gly Val Asp Thr Ala Ala Thr Gly Gly Val Phe Asp
 355 360 365
 Ile Ser Asn Leu Asp Arg Leu Gly Lys Ser Glu Val Glu Leu Val Gln
 370 375 380
 Leu Val Ile Asp Gly Val Asn Tyr Leu Ile Asp Cys Glu Arg Arg Leu
 385 390 395 400
 Glu Arg Gly Gln Asp Ile Arg Ile Pro Thr Pro Val Ile His Thr Lys
 405 410 415
 His

<210> 48
 <211> 2365
 <212> DNA
 <213> Homo sapiens

<400> 48
 ggagcggag agcgacgcgc gctgcagccg gcggcatggc tagcacggct tcggagatca 60
 tcgccttcac ggtctccatc tcaggtctgg tactggtgtc ctccacgctg cccaccgact 120

```

actggaaggt gtctaccatc gacggcacgg tcatcacaac cgccacctat tggggcaacc 180
tgtggaagcg gtgcgttacc gactccacgg gcgtctccaa ctgcaaggac ttccctcca 240
gtcgtggcggt ggaagcgttat atacaggcat gtgaggagact tatgatcgct gctgcgaagc 300
tgggctctctt tgggtccata ttgogctct ttggaatgaa gtgtaccaaa gtgcaggagct 360
cggataaagc caaagctaaa attgcttgtt tggctggggt tgattatcata ctgtcagggc 420
gtgtctcaat gactggatgt tccctatatg caacacaaat cacaacggaa ttctttgatc 480
ctctctttgt tgagcaaaaag tatgaattag gagcgcgtct gttttattgga tgggcaggag 540
cctcactgtg cataatttgg tgggtccatc ttgtcttttc aatatctgac aacacaaaaa 600
caccagata ccaattcaaac gggggccacat ctgtcatgtc ttctcggaga aagtatcatt 660
gtggagaaga ttttaaaaca acaaaccttt caaaacagtt tgataaaaaa gcttatgtct 720
aaaagagctc gctggcaagc tgctcttga gtttgtata aaagcgaaat gttcacaaaa 780
tgatccctac aaggccctcc tgctcttga cataattaac actcaaaact atttttaaaa tatgcatttg 840
aagcatctgt tgattgtatg gatgtaagtg ttcttacata gttagtata tactaatcat 900
ttctctgttg gctttcttat aaaaaataaa cagtttatit acaggatttg taaaatgttt 960
tctacattta tatagaacat gaaaagcatt tagtaccaaa ggttcaagaa gtattcgtac 1020
tctagccttt ttaactcatt atagatagaa tcttttgatc ccactcctta ttgttctttt 1080
cattcataaa caggtgtata aggaacaatg tcttataaac agcatggggg caatctgaga 1140
atattctcca aaaggtgtcc aggttaaata gacatgttac tggctgcaca caggcaaat 1200
ctagtttgtt ttttttaagt atctacaac attatttaa aaaggttaaa ctttttgttg 1260
aagcagaag ttactctgta gaacttaact tctacaggat cagagaggat cttgctcatt 1320
catggccata tcccatgccc catggccact cagttagatt ttgaaaaagc aaagccacac 1380
cattctcttt gtctgtatga tagcaggga tgctctctga ttcttccac 1440
tgccacattt agtgaattat tagttgtttt cataaacgat gctgtgatga agactcatgt 1500
acatatttga caaattttgg ttctttacat gtgcctgtca tgactgtaat tcattatgac 1560
gtctccagga agggctaagt gggccaatat attattgcct gtcagtggc acatccatgt 1620
taaggggctg aggcgtccct ggcacggaat gcagggccct atcagcayag 1680
gctgagatag agatatttgt catggttgac tgaggagcca attaaaacct gtttatgcct 1740
agtgttccat tattggaaca ctaagcatgt gggagttatt tatatcctac tgcctcaaggt 1800
catcgccaag gtgtgatttg aaaaattcaa aaaattgcaa cctcaggcat aaatgggtta 1860
aggaatccc aagcccaagt ggtacgtgcc tcaactcaga ctgacgggac gagtctcttc 1920
taggtgtgct ttccagaacc tgtttacggc taactggata actgagagac tgctcatttc 1980
taaagcatt taagtgtctc cagggatctc tgaaaaaaga cacaggcttc ttcttagagc 2040
cagcctata taactgccc acaaggcaca cagttatcac agttcataca cactttctat 2100
gtcctgtctc actcactcct cacagccatc ctaggagata catattgttt tcatcctgca 2160
tttacagaaa aagaaatgaa aacagagagc ttaataaatt tgccacagta atgtcgaaa 2220
taggcctttg aaccaaggca gtctagggta aaatatagtt tcaagtatg aataagaatt 2280
ggatatttg ttatctttga gtaagaaact gtccgatatg aatcacaacg tgggtgaagt 2340
tagtattttc ctgaagtgtg aaaga 2365

```

<210> 49

<211> 228

<212> PRT

<213> Homo sapiens

<400> 49

```

Met Ala Ser Thr Ala Ser Glu Ile Ile Ala Phe Met Val Ser Ile Ser
  1           5           10           15
Gly Trp Val Leu Val Ser Ser Thr Leu Pro Thr Asp Tyr Trp Lys Val
  20           25           30
Ser Thr Ile Asp Gly Thr Val Ile Thr Thr Ala Thr Tyr Trp Ala Asn
  35           40           45
Leu Trp Lys Ala Cys Val Thr Asp Ser Thr Gly Val Ser Asn Cys Lys
  50           55           60
Asp Phe Pro Ser Met Leu Ala Leu Asp Gly Tyr Ile Gln Ala Cys Arg
  65           70           75           80
Gly Leu Met Ile Ala Ala Val Ser Leu Gly Phe Phe Gly Ser Ile Phe
  85           90           95
Ala Leu Phe Gly Met Lys Cys Thr Lys Val Gly Gly Ser Asp Lys Ala

```

100 105 110
 Lys Ala Lys Ile Ala Cys Leu Ala Gly Ile Val Phe Ile Leu Ser Gly
 115 120 125
 Leu Cys Ser Met Thr Gly Cys Ser Leu Tyr Ala Asn Lys Ile Thr Thr
 130 135 140
 Glu Phe Phe Asp Pro Leu Phe Val Glu Gln Lys Tyr Glu Leu Gly Ala
 145 150 155 160
 Ala Leu Phe Ile Gly Trp Ala Gly Ala Ser Leu Cys Ile Ile Gly Gly
 165 170 175
 Val Ile Phe Cys Phe Ser Ile Ser Asp Asn Asn Lys Thr Pro Arg Tyr
 180 185 190
 Thr Tyr Asn Gly Ala Thr Ser Val Met Ser Ser Arg Thr Lys Tyr His
 195 200 205
 Gly Gly Glu Asp Phe Lys Thr Thr Asn Pro Ser Lys Gln Phe Asp Lys
 210 215 220
 Asn Ala Tyr Val
 225

 <210> 50
 <211> 1024
 <212> DNA
 <213> Homo sapiens

 <400> 50
 cccacccga aacacactca gcccttgcac tgacctgcct tctgattgga ggctgggtgc 60
 ttccgataat gaacctcagg accccactgt tggttacagc ctgtttgtat tattcttact 120
 gcaactcaag acacctgcag caggggcgta gaaaaagtaa aagaccagta tttcacatt 180
 gccaggtacc agaacaacag aagactgaca ccggccaatt aagtggggcc agggctggty 240
 tctgccatg ttgccatcct gatgggctgc ttgcacaat gagggatctt ctccaatata 300
 tcgtctgctt ctttgccttt ttctctgctg ggtttttgat tgtggccacc tggactgact 360
 gttggtggt gaatgctgat gaactctctg aggtgagcac aaaatgccga ggctctggt 420
 gggaatcggt cacaaatgct ttgatggga ttgcacctg tgatgagtac gattccatatac 480
 ttgcggagca tcccttgaag ctggtggtta ctgcagcgtt gatgattact gcagatatc 540
 tagctgggtt tgattttctc accctgctcc ttggtcttga ctgcgtgaaa ttctccctg 600
 atgagccgta catiaaagtc cgcactctgct ttgttctg agccacgcta ctaatagcac 660
 gtacccagg aatcattggc tctgtgtggt atgctgttga tgtgtatgtg gaaccttcta 720
 ctttggtttt gcacaatata tttcttggtta tccaatataa atttgggttg tctctgtggc 780
 ctggaaatggc tgggtctctg ggttgccttt tggctggagc tgttctcacc tgctgcttat 840
 atctttttaa agatgttggg cctgagagaa actatcctta ttccttgagg aaagcctatt 900
 cagccgcggg tgtttccatg gccaaagtc actcagcccc tcgcacagag acggccaaaa 960
 tgtatgctgt agacacaagg gtgtaaaatg cactgttcag ggtgtgttg catatgattt 1020
 aatc 1024

<210> 51
 <211> 305
 <212> PRT
 <213> Homo sapiens

<400> 51
 Met Thr Ser Arg Thr Pro Leu Leu Val Thr Ala Cys Leu Tyr Tyr Ser
 1 5 10 15
 Tyr Cys Asn Ser Arg His Leu Gln Gln Gly Val Arg Lys Ser Lys Arg
 20 25 30
 Pro Val Phe Ser His Cys Gln Val Pro Glu Thr Gln Lys Thr Asp Thr
 35 40 45
 Arg His Leu Ser Gly Ala Arg Ala Gly Val Cys Pro Cys Cys His Pro
 50 55 60

59

Asp Gly Leu Leu Ala Thr Met Arg Asp Leu Gln Tyr Ile Ala Cys
 65 70 75 80
 Phe Phe Ala Phe Phe Ser Ala Gly Phe Leu Ile Val Ala Thr Trp Thr
 85 90 95
 Asp Cys Trp Met Val Asn Ala Asp Asp Ser Leu Glu Val Ser Thr Lys
 100 105 110
 Cys Arg Gly Leu Trp Trp Glu Cys Val Thr Asn Ala Phe Asp Gly Ile
 115 120 125
 Arg Thr Cys Asp Glu Tyr Asp Ser Ile Leu Ala Glu His Pro Leu Lys
 130 135 140
 Leu Val Val Thr Arg Ala Leu Met Ile Thr Ala Asp Ile Leu Ala Gly
 145 150 155 160
 Phe Gly Phe Leu Thr Leu Leu Leu Gly Leu Asp Cys Val Lys Phe Leu
 165 170 175
 Pro Asp Glu Pro Tyr Ile Lys Val Arg Ile Cys Phe Val Ala Gly Ala
 180 185 190
 Thr Leu Leu Ile Ala Gly Thr Pro Gly Ile Ile Gly Ser Val Trp Tyr
 195 200 205
 Ala Val Asp Val Tyr Val Glu Arg Ser Thr Leu Val Leu His Asn Ile
 210 215 220
 Phe Leu Gly Ile Gln Tyr Lys Phe Gly Trp Ser Cys Trp Leu Gly Met
 225 230 235 240
 Ala Gly Ser Leu Gly Cys Phe Leu Ala Gly Ala Val Leu Thr Cys Cys
 245 250 255
 Leu Tyr Leu Phe Lys Asp Val Gly Pro Glu Arg Asn Tyr Pro Tyr Ser
 260 265 270
 Leu Arg Lys Ala Tyr Ser Ala Ala Gly Val Ser Met Ala Lys Ser Tyr
 275 280 285
 Ser Ala Pro Arg Thr Glu Thr Ala Lys Met Tyr Ala Val Asp Thr Arg
 290 295 300
 Val
 305

<210> 52

<211> 1665

<212> DNA

<213> Homo sapiens

<400> 52

gaaggaaact gttctgctca cacttgctgg ctgogcgc aggaactggct ttatctctctg 60
 actcaagggtg caaagggtgca ctctgcgaac gttaagtccg tcccagcgc ttggaatcct 120
 acggccccca cagccgggac cctcaagcct tccaggtcct caactccctg ggacgtgaa 180
 caatggcttc catggggcta caggtaatgg gcatcgcgct ggccgtctct ggctgggtg 240
 ccgtcatgct gtgctgcgag ctgcccattgt ggccgtgac ggccctcctc ggacgcaaca 300
 ttgtcaacct gcagaccatc tgggaggggc tatggatgaa ctgcgtgggtg cagagcacccg 360
 gcagatagca gtgcaagggtg tacgactcgc tgcctggcact gccagcaggac ctgcaggcgg 420
 ccggcgccct cgtcatcctc agcatcatcg tggctgctct ggccgtgctg ctgtccctgg 480
 tgggggggcaa gtgtaccaac tgcctggagg atgaagcgc caaggcccaag accatgatcg 540
 tggcgggcgt ggtgttctctg ttggccggcc ttatggtgat agtgccgggtg tcctggacgg 600
 cccacaacat caaccaagac ttctacaatc cgtctggtgc ctccggggcag aagcgggaga 660
 tgggtgcctc gctctacgtc ggtcggggcc cctccggcct gctgctcctt ggccggggggc 720
 tgccttctgt caactgtcca cccgcacag acaagcctta ctccggccaag tatctgctg 780
 ccgcctctcg tgcctgccagc aactacgtgt aaggtgccac ggctccaact tgttctctc 840
 tgccttggtc tccctgggac tgagctcagc gcaggctgtg accccaggag gccctgccca 900
 cggggcaactg gctgctgggg actggggact ggcagagac tgagccaggc aggaagagca 960
 cagccttcag cctctctggc ccaactcggc aacttcccaa ggccgcctcc ttgtgcgaag 1020
 aacagagtcc accctcctct ggatattggg gagggacgga agtgacaggg tgtggtggtg 1080

```

gagtgggggag ctggccttctg ctggccagga tagcttaacc ctgacttttg gatctgacctg 1140
catcggcgtt ggccactgtc ccatttaca ttttcccac tctgtctgcc tgcattcct 1200
ctgttccggg taggccttga tatcactct gggactgtgc ctgtctcacc gaaaccccg 1260
cccaggagta tggtcgaggc cttgccacc cactgcctg ggaagtgcag agtggaagg 1320
cggttttaga ggggaggggc gaaggtgctg taaacaggtt tgggcagtg tggggaggg 1380
ggccagagag gcggctcagg ttgccacgt ctgtggcctc aggaactctt gctcaccoc 1440
cttcagccca gggccoctgg agactgatcc cctctgagtc ctctgccct tccaaggaca 1500
ctaattgagcc tgggaggggtg gcaggaggga ggggacagct tcacccttg aagtcctgg 1560
gttttctctc ttcctcttt gtggtttctg ttttgaatt taagaagagc tattcatcac 1620
tgtaattatt attattttct acaataaatg ggacctgtgc acagg 1665

```

<210> 53

<211> 209

<212> PRT

<213> Homo sapiens

<400> 53

```

Met Ala Ser Met Gly Leu Gln Val Met Gly Ile Ala Leu Ala Val Leu
1          5          10          15
Gly Trp Leu Ala Val Met Leu Cys Cys Ala Leu Pro Met Trp Arg Val
20          25          30
Thr Ala Phe Ile Gly Ser Asn Ile Val Thr Ser Gln Thr Ile Trp Glu
35          40          45
Gly Leu Trp Met Asn Cys Val Val Gln Ser Thr Gly Gln Met Gln Cys
50          55          60
Lys Val Tyr Asp Ser Leu Leu Ala Leu Pro Gln Asp Leu Gln Ala Ala
65          70          75          80
Arg Ala Leu Val Ile Ser Ile Ile Val Ala Ala Leu Gly Val Leu
85          90          95
Leu Ser Val Val Gly Gly Lys Cys Thr Asn Cys Leu Glu Asp Glu Ser
100         105         110
Ala Lys Ala Lys Thr Met Ile Val Ala Gly Val Val Phe Leu Leu Ala
115         120         125
Gly Leu Met Val Ile Val Pro Val Ser Trp Thr Ala His Asn Ile Ile
130         135         140
Gln Asp Phe Tyr Asn Pro Leu Val Ala Ser Gly Gln Lys Arg Glu Met
145         150         155         160
Gly Ala Ser Leu Tyr Val Gly Trp Ala Ala Ser Gly Leu Leu Leu Leu
165         170         175
Gly Gly Gly Leu Leu Cys Cys Asn Cys Pro Pro Arg Thr Asp Lys Pro
180         185         190
Tyr Ser Ala Lys Tyr Ser Ala Ala Arg Ser Ala Ala Ala Ser Asn Tyr
195         200         205
Val

```

<210> 54

<211> 3457

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<222> (1)...(3457)

<223> n = A,T,C or G

<400> 54

atgaagattt	tgatacttgg	tattttttctg	ttttttatgta	gtacccccagc	ctgggcgaaa	60
gaaaagcatt	attacattgg	aattattgaa	acgacttggg	attatgcctc	tgaccatggg	120
gaaaaaagac	tattttctgt	tgacacggaa	cattccaata	tctatctcca	aaatggccca	180
gtagaattgt	gggagctata	taagaaggcc	cttatctctc	agtaacacaga	tgaacctttt	240
aggacaacta	tagaaaaaac	ggctctggctt	gggttttttag	gccttattat	caaatgctgaa	300
actggagata	aagttttatgt	acaottaaaa	accttgcct	ctaggccctc	caoccttcac	360
tcacatggaa	taacttacta	taaggaacat	gagggggcca	tctaccttga	taacaccaca	420
gattttcaaa	gagcagatga	caaatgtat	ccaggagagc	agtatacata	catgttgctt	480
gcacatgaag	aaccaagtc	tggggaagga	gtggccaatt	gtgtgactag	gattttaccat	540
tcaccacattg	atgtctccaa	agatattgcc	tcaggactca	tcggacacct	aatatctgt	600
aaaaaagatt	ctctagataa	agaaaaagaa	aaacatttg	accgagaatt	tgtgtgtgat	660
ttttctgtgg	tggatgaaaa	tttcagctgg	tacctagaag	acaacattaa	aacctactgc	720
tcagacccag	agaaagttga	caaagacaac	gaagacttcc	aggagagtaa	cagatgtgat	780
tcgtgtaagt	gatcacacttt	tggaaagtctc	ccaggactct	ccatgtgtgc	tgaagacaga	840
gtaaaaatggt	acctttttgg	tatgggtta	gaagtgtgat	tgcacgcagc	ttcttttccac	900
gggcaacgac	tgactaaccaa	gaactacogt	attgacacaa	tcaacctctt	tctctgtacc	960
ctgttttgatg	ctttataggt	ggccagaaac	ctgggaagt	ggatgctcag	ctgtcagaat	1020
ctaaaccatc	tgaagccgg	tttgcaagcc	tttttccagg	tcaggagtg	taacaagtct	1080
tcatacaagg	ataatattcc	tgggaagcat	gttagacact	actacattgc	cgctgaggaa	1140
atcatcttga	actatgctcc	ctctggtata	gacatctcca	ctaaagaaaa	cttaaacaga	1200
ccctggaaagt	actcagcggt	gttttttgaa	caaggtaacca	caagaattgg	aggctcttat	1260
aaaaaacctgg	tttatcttga	gtacacagat	gcctctctca	caaacctgaa	ggagagagcc	1320
cctgaagaag	agacattctgg	catcctgggt	cctgtcattt	gggacagagt	gggagacact	1380
atcagagtaa	cccttccataa	caaaaggaca	tctccctcca	gtattgagcc	gattggggct	1440
agattcaata	agaaacaaga	gggcacatac	tattccccaa	attacaaccc	ccagagcaga	1500
agtgtgcctc	cttcagcctc	ccatgtggca	cccacagaaa	cattcaacct	tgaattggact	1560
gtccccaag	aagtagaaccc	cactaatgca	gatcctgtgt	gtctagctaa	gattgtattat	1620
tcgtctgtgg	attccactaa	agatatatto	ctggggctta	ttggggccaat	gaaatatgct	1680
aagaagggaa	gtttacatgc	aaatgggaga	cagaaagatg	tagacaagga	attctatttg	1740
tttccctacag	tatttgatga	gaatgagagt	ttactcctgg	aagataatat	tagaatgttt	1800
acaaactgcac	gatctcaggt	ggataaggaa	gatgaagact	ttcaggaaatc	taataaaatg	1860
cactccatga	atggatctcat	gtatgggaat	cagccgggtc	tcacttatgtg	caaaaggagat	1920
tcggtcgtgt	ggtaacttatt	cagccgcgga	aatgagccgc	atgtacatgg	aatcatattt	1980
tcaggaaaaa	cataatctgtg	gagaggagaa	cggagagaca	cagcaaaactc	cttccctcaa	2040
acaagtctta	cgctccacat	gtggcctgac	acagagggga	cttttaagt	tgaatgcctt	2100
acaaactgatc	attacacagg	cggcatgaag	caaaaaata	ctgtgaacca	atgcaggcgg	2160
cagtcctgag	attccacctt	ctacctggga	gagaggacat	actatatcgc	agcgtggag	2220
gtggaaatggg	attattcccc	acaaaaggga	tgggaaaagg	agctgcacat	tttacaagag	2280
cagaatgttt	acaatgcatt	tttagataag	ggagagtttt	acataaggctc	aaagtacaag	2340
aaagtgtgtg	atcggcagta	tactgatagc	acattccgtg	ttccagtgga	gagaaaaagct	2400
gaagaagaac	tttgggaat	tctaggtcca	caactctact	cagatgtgtg	agacaaagtc	2460
aaaaattatct	ataaaaaaat	ggccacaagg	ccactctcaa	tacatgcca	tggggtacaa	2520
acagagaatt	ctacagttac	tcacaacatta	ccagtgtaaa	ctctccacta	cgtatggaaa	2580
atccacagaa	ctctctggagc	tggaaacagag	gattctgctt	gtattccatg	gccttattat	2640
tcacactgtg	atcaagttaa	ggacctctac	agtgatttaa	ttggcccccct	gattgtttgt	2700
cgaagacctt	acttgaaggt	attcaatccc	agaaaggaa	tggaaatttgc	ctctctgttt	2760
ctagtttttg	atgagaatga	atcttggtag	ttagatgaca	acatcaaaaac	atactctgat	2820
caccocgaga	aagtaaaacaa	agatgatgag	atttcatag	aaagcaataa	aatgcactgt	2880
attaatggaa	gaatgttttg	aaacctacaa	ggcctcacaa	tgcacgtggg	agatgaagtc	2940
aactgggtata	tgatgggaat	gggcaatgaa	ataccttcat	acaactgtaca	ttttccacgc	3000
catagcttcc	aatacaagca	caggggagtt	tatagttctg	atgtcttga	cattttccct	3060
ggaaacatacc	aaaccctaga	aatgtttcca	agaacacctg	gaattttggt	actccactgc	3120
catgttgaccg	accacattca	tgttggaaatg	acacacctct	acacgccttc	acaaaattgaa	3180
gggtgaattac	cagacaccaa	atctggctga	atgaaaataa	tgtgtgataa	gtggaaaaaa	3240
gagaaaaaac	aatgattcat	acaaatgata	gtgaaagtgt	aaaaatagaa	gttactttgg	3300
aatgactata	aacattataaa	gnagactgn	agcatacaac	tttgtacatt	tgtgggggaa	3360
aacttaaat	tttgcccaaa	tggaaaagatc	acagactat	ataatgatac	atgactgaca	3420
ctgttacact	aggtataaaa	actgattcat	acagctct			3480

<210> 55
 <211> 1069
 <212> PRT
 <213> Homo sapiens

<400> 55

```

Met Lys Ile Leu Ile Leu Gly Ile Phe Leu Phe Leu Cys Ser Thr Pro
 1           5           10           15
Ala Trp Ala Lys Glu Lys His Tyr Tyr Ile Gly Ile Ile Glu Thr Thr
 20           25           30
Trp Asp Tyr Ala Ser Asp His Gly Glu Lys Lys Leu Ile Ser Val Asp
 35           40           45
Thr Glu His Ser Asn Ile Tyr Leu Gln Asn Gly Pro Asp Arg Ile Gly
 50           55           60
Arg Leu Tyr Lys Lys Ala Leu Tyr Leu Gln Tyr Thr Asp Glu Thr Phe
 65           70           75
Arg Thr Thr Ile Glu Lys Pro Val Trp Leu Gly Phe Leu Gly Pro Ile
 85           90           95
Ile Lys Ala Glu Thr Gly Asp Lys Val Tyr Val His Leu Lys Asn Leu
100           105           110
Ala Ser Arg Pro Tyr Thr Phe His Ser His Gly Ile Thr Tyr Tyr Lys
115           120           125
Glu His Glu Gly Ala Ile Tyr Pro Asp Asn Thr Thr Asp Phe Gln Arg
130           135           140
Ala Asp Asp Lys Val Tyr Pro Gly Glu Gln Tyr Thr Tyr Met Leu Leu
145           150           155
Ala Thr Glu Glu Gln Ser Pro Gly Glu Gly Asp Gly Asn Cys Val Thr
165           170           175
Arg Ile Tyr His Ser His Ile Asp Ala Pro Lys Asp Ile Ala Ser Gly
180           185           190
Leu Ile Gly Pro Leu Ile Ile Cys Lys Lys Asp Ser Leu Asp Lys Glu
195           200           205
Lys Glu Lys His Ile Asp Arg Glu Phe Val Val Met Phe Ser Val Val
210           215           220
Asp Glu Asn Phe Ser Trp Tyr Leu Glu Asp Asn Ile Lys Thr Tyr Cys
225           230           235
Ser Glu Pro Glu Lys Val Asp Lys Asp Asn Glu Asp Phe Gln Glu Ser
245           250           255
Asn Arg Met Tyr Ser Val Asn Gly Tyr Thr Phe Gly Ser Leu Pro Gly
260           265           270
Leu Ser Met Cys Ala Glu Asp Arg Val Lys Trp Tyr Leu Phe Gly Met
275           280           285
Gly Asn Glu Val Asp Val His Ala Ala Phe Phe His Gly Gln Ala Leu
290           295           300
Thr Asn Lys Asn Tyr Arg Ile Asp Thr Ile Asn Leu Phe Pro Ala Thr
305           310           315
Leu Phe Asp Ala Tyr Met Val Ala Gln Asn Pro Gly Glu Trp Met Leu
325           330           335
Ser Cys Gln Asn Leu Asn His Leu Lys Ala Gly Leu Gln Ala Phe Phe
340           345           350
Gln Val Gln Glu Cys Asn Lys Ser Ser Ser Lys Asp Asn Ile Arg Gly
355           360           365
Lys His Val Arg His Tyr Tyr Ile Ala Ala Glu Glu Ile Ile Trp Asn
370           375           380
Tyr Ala Pro Ser Gly Ile Asp Ile Phe Thr Lys Glu Asn Leu Thr Ala
385           390           395
Pro Gly Ser Asp Ser Ala Val Phe Phe Glu Gln Gly Thr Thr Arg Ile
400

```

405										410					415				
Gly	Gly	Ser	Tyr	Lys	Lys	Leu	Val	Tyr	Arg	Glu	Tyr	Thr	Asp	Ala	Ser				
Phe	Thr	Asn	Arg	Lys	Glu	Arg	Gly	Pro	Glu	Glu	Glu	His	Leu	Gly	Ile				
Leu	Gly	Pro	Val	Ile	Trp	Ala	Glu	Val	Gly	Asp	Thr	Ile	Arg	Val	Thr				
Phe	His	Asn	Lys	Gly	Ala	Tyr	Pro	Leu	Ser	Ile	Glu	Pro	Ile	Gly	Val				
Arg	Phe	Asn	Lys	Asn	Asn	Glu	Gly	Thr	Tyr	Tyr	Ser	Pro	Asn	Tyr	Asn				
Pro	Gln	Ser	Arg	Ser	Val	Pro	Pro	Ser	Ala	Ser	His	Val	Ala	Pro	Thr				
Glu	Thr	Phe	Thr	Tyr	Glu	Trp	Thr	Val	Pro	Lys	Glu	Val	Gly	Pro	Thr				
Asn	Ala	Asp	Pro	Val	Cys	Leu	Ala	Lys	Met	Tyr	Tyr	Ser	Ala	Val	Asp				
Pro	Thr	Lys	Asp	Ile	Phe	Thr	Gly	Leu	Ile	Gly	Pro	Met	Lys	Ile	Cys				
Lys	Lys	Gly	Ser	Leu	His	Ala	Asn	Gly	Arg	Gln	Lys	Asp	Val	Asp	Lys				
Glu	Phe	Tyr	Leu	Phe	Pro	Thr	Val	Phe	Asp	Glu	Asn	Glu	Ser	Leu	Leu				
Leu	Glu	Asp	Asn	Ile	Arg	Met	Phe	Thr	Thr	Ala	Pro	Asp	Gln	Val	Asp				
Lys	Glu	Asp	Glu	Asp	Phe	Gln	Glu	Ser	Asn	Lys	Met	His	Ser	Met	Asn				
Gly	Phe	Met	Tyr	Gly	Asn	Gln	Pro	Gly	Leu	Thr	Met	Cys	Lys	Gly	Asp				
Ser	Val	Val	Trp	Tyr	Leu	Phe	Ser	Ala	Gly	Asn	Glu	Ala	Asp	Val	His				
Gly	Ile	Tyr	Phe	Ser	Gly	Asn	Thr	Tyr	Leu	Trp	Arg	Gly	Glu	Arg	Arg				
Asp	Thr	Ala	Asn	Leu	Phe	Pro	Gln	Thr	Ser	Leu	Thr	Leu	His	Met	Trp				
Pro	Asp	Thr	Glu	Gly	Thr	Phe	Asn	Val	Glu	Cys	Leu	Thr	Thr	Asp	His				
Tyr	Thr	Gly	Gly	Met	Lys	Gln	Lys	Tyr	Thr	Val	Asn	Gln	Cys	Arg	Arg				
Gln	Ser	Glu	Asp	Ser	Thr	Phe	Tyr	Leu	Gly	Glu	Arg	Thr	Tyr	Tyr	Ile				
Ala	Ala	Val	Glu	Val	Glu	Trp	Asp	Tyr	Ser	Pro	Gln	Arg	Glu	Trp	Glu				
Lys	Glu	Leu	His	His	Leu	Gln	Glu	Gln	Asn	Val	Ser	Asn	Ala	Phe	Leu				
Asp	Lys	Gly	Glu	Phe	Tyr	Ile	Gly	Ser	Lys	Tyr	Lys	Lys	Val	Val	Tyr				
Arg	Gln	Tyr	Thr	Asp	Ser	Thr	Phe	Arg	Val	Pro	Val	Glu	Arg	Lys	Ala				
Glu	Glu	Glu	His	Leu	Gly	Ile	Leu	Gly	Pro	Gln	Leu	His	Ala	Asp	Val				
Gly	Asp	Lys	Val	Lys	Ile	Ile	Phe	Lys	Asn	Met	Ala	Thr	Arg	Pro	Tyr				
Ser	Ile	His	Ala	His	Gly	Val	Gln	Thr	Glu	Ser	Ser	Thr	Val	Thr	Pro				
Thr	Leu	Pro	Gly	Glu	Thr	Leu	Thr	Tyr	Val	Trp	Lys	Ile	Pro	Glu	Arg				
Ser	Gly	Ala	Gly	Thr	Glu	Asp	Ser	Ala	Cys	Ile	Pro	Trp	Ala	Tyr	Tyr				

865				870				875				880			
Ser	Thr	Val	Asp	Gln	Val	Lys	Asp	Leu	Tyr	Ser	Gly	Leu	Ile	Gly	Pro
				885				890				895			
Leu	Ile	Val	Cys	Arg	Arg	Pro	Tyr	Leu	Lys	Val	Phe	Asn	Pro	Arg	Arg
			900					905				910			
Lys	Leu	Glu	Phe	Ala	Leu	Leu	Phe	Leu	Val	Phe	Asp	Glu	Asn	Glu	Ser
		915					920				925				
Trp	Tyr	Leu	Asp	Asp	Asn	Ile	Lys	Thr	Tyr	Ser	Asp	His	Pro	Glu	Lys
	930				935						940				
Val	Asn	Lys	Asp	Asp	Glu	Glu	Phe	Ile	Glu	Ser	Asn	Lys	Met	His	Ala
	945				950					955					960
Ile	Asn	Gly	Arg	Met	Phe	Gly	Asn	Leu	Gln	Gly	Leu	Thr	Met	His	Val
				965						970				975	
Gly	Asp	Glu	Val	Asn	Trp	Tyr	Leu	Met	Gly	Met	Gly	Asn	Glu	Ile	Asp
			980					985				990			
Leu	His	Thr	Val	His	Phe	His	Gly	His	Ser	Phe	Gln	Tyr	Lys	His	Arg
	995						1000					1005			
Gly	Val	Tyr	Ser	Ser	Asp	Val	Phe	Asp	Ile	Phe	Pro	Gly	Thr	Tyr	Gln
	1010				1015						1020				
Thr	Leu	Glu	Met	Phe	Pro	Arg	Thr	Pro	Gly	Ile	Trp	Leu	Leu	His	Cys
	1025				1030				1035						1040
His	Val	Thr	Asp	His	Ile	His	Ala	Gly	Met	Glu	Thr	Thr	Tyr	Thr	Val
				1045				1050						1055	
Leu	Gln	Asn	Glu	Gly	Glu	Tyr	Pro	Asp	Thr	Lys	Ser	Gly			
			1060				1065								

<210> 56

<211> 2807

<212> DNA

<213> Homo sapiens

<400> 56

cgtgggtgatg	ttttctgtgg	tggatgaaaa	tttcagctgg	tacctagaag	acaacattaa	60
aacctactgc	tcagaaccag	agaaagtga	caaagacaa	gaagacttcc	aggagagtaa	120
cagaatgtat	tctgtgaatg	gataacattt	tggaaagtctc	ccaggactct	ccatgtgtgc	180
tgaagacaga	gtaaaatgg	aoccttttgg	tatgggtaat	gaagttgatg	tgcaagcagc	240
ttttcttccac	gggcaagcac	tgactaacaa	gaactacocgt	attgacacaa	tcaaacctctt	300
tctgtctaac	ctgttttgatg	cttatatgg	ggccacgaac	ctgtgagaat	ggatgtctacg	360
ctgtcagaat	ctaaaccatc	tgaagccgg	tttgcaggcc	tttttccagg	tccaggagtg	420
taacaagtct	tcatacaagg	ataatatccg	tgggaagcat	gttagacact	actacattgc	480
cctgtaggaa	atcatctgga	actatgctcc	ctctgtgata	gacactctca	ctaaagaaaa	540
cttaacagca	ccctggaagt	actcagcgg	gttttttgaa	caaggtacca	caagaattgg	600
aggctcttat	aaaaagctgg	tttatcgtga	gtacacagat	gcctccttca	caaatcgaaa	660
ggagagaggo	oatgaagaag	agcatcttgg	catctcgtgt	cctgtcattt	gggcagaggt	720
gggagacacc	atcagagtta	ccttcacata	caaaggagca	tatccctcca	gtattgagoc	780
gattgggtgt	agattcaata	agaacaaaga	gggcacatac	tattccocaa	attacaaccc	840
ccagagacaga	agtgtgcctc	cttcagcctc	ccatgtggca	ccacacagaa	cattcaccta	900
tgaatggact	gtcccccaga	aagtaggacc	cactaatgca	gatcctgtgt	gtctagotaa	960
gatgtattat	tctgtgttgg	atcccaactaa	aatcatattc	actgggctta	tggggccaat	1020
gaaaatatgc	agaaaaggaa	gtttacatgc	aaatgggaga	cagaaagatg	tgcacaaagg	1080
attctatttt	tttctctacg	tatttgatga	tgatgagagt	ttaactcctg	aagataatat	1140
tagaattgtt	acaactgcac	ctgatcaggt	ggataaggaa	gatgaagact	ttcagggaatc	1200
taataaaatg	cactccatga	atggattcat	gtatgggaat	cagccgggtc	tcactatgtg	1260
caaaaggagat	tgggtcgtgt	ggtaacttatt	cagccgcgga	aatgagggcc	atgtacatgtg	1320
aatatatttt	tcagaaacaa	catatctgtg	gagaggagaa	cggagagaca	cagcaaacct	1380
cttcctccaa	acaagtctta	cgtccacat	gtggcctgac	acagagggga	cttttaattg	1440
tgaatgcctt	acaactgac	attacacagg	cggcatgaag	caaaaaata	ctgtgaacca	1500

```

atgcaggcgg cagtctgagg attccacott ctacctggga gagaggacat actatatcgc 1560
agcagtgagg gtgggaatggg attattcccc acaaaggagg tgggaaaagg agctgcacaa 1620
tttacaagag cagaatgttt caaatgcatt tttagataag ggagagtttt acataggctc 1680
aaagtacaag aaagtgtgtg atcggcagta tactgatagc acattccgtg ttccagtggg 1740
gagaaaagct gaagaagaac atctgggaat tctagggtcca caacttcacg cagatgttgg 1800
agacaaaagt aaaaattatct ttaaaaacat ggccacaagg cactactcaa tacatgccca 1860
tgggggtacaa acagagaggtt ctacagttac tccaacatta ccaggtgaaa ctctcaccta 1920
cgatgagaaa attcccagaaa gatctggagc tggaaacagag gattctgctt gtattccatg 1980
ggcttattat tcaactgtgg atcaagttaa ggacctctac agtggattaa ttggccccct 2040
gattgtttgt cgaagacott acttgaaagt attcaatccc agaaggaac tggaatttgc 2100
cctctgtttt ctagtttttg atgagaatga atcttggtac ttagatgaca acatcaaaac 2160
atactctgat caccocgaga aagtaaacaa agatgatgag gaattcatag aaagcaataa 2220
aatgcattgc attaatggaa gaatgtttgg aaacctacaa ggctcaca tgacgtgggg 2280
agatgaagtc aactggtatc tgatgggaat gggaatgaa atagacttac acactgtaca 2340
ttttcacggc catagcttcc aatacaagca caggggagtt tatagttctg atgtctttga 2400
cattttccct ggaacatacc aaaccctaga aatgtttcca agaacaactg gaatttgggt 2460
actccactgc catgtgacccg accacattca tgctggaagt gaaaccaact acacggctct 2520
acaaaattgaa ggtgaatatc cagacaccaa atctggctga atgaaataaa ttgggtgataa 2580
gtggaaaaaa gagaaaaaac aatgattcat aacaattgat gtgaaagtgt aaaatagaat 2640
gttactttgg aatgactata aacattaaaa gaagactgga agcatacaac ttgtacatt 2700
tgtgggggaa aactatataa tttgcccaaa tgggaagatc aacagactat ataagtatac 2760
atgactgaca cttgtacact aggtataataa actgattcat acagtct 2807

```

<210> 57

<211> 852

<212> PRT

<213> Homo sapiens

<400> 57

```

Val Val Met Phe Ser Val Val Asp Glu Asn Phe Ser Trp Tyr Leu Glu
1 5 10 15
Asp Asn Ile Lys Thr Tyr Cys Ser Glu Pro Glu Lys Val Asp Lys Asp
20 25 30
Asn Glu Asp Phe Gln Glu Ser Asn Arg Met Tyr Ser Val Asn Gly Tyr
35 40 45
Thr Phe Gly Ser Leu Pro Gly Leu Ser Met Cys Ala Glu Asp Arg Val
50 55 60
Lys Trp Tyr Leu Phe Gly Met Gly Asn Glu Val Asp Val His Ala Ala
65 70 75 80
Phe Phe His Gly Gln Ala Leu Thr Asn Lys Asn Tyr Arg Ile Asp Thr
85 90 95
Ile Asn Leu Phe Pro Ala Thr Leu Phe Asp Ala Tyr Met Val Ala Gln
100 105 110
Asn Pro Gly Glu Trp Met Leu Ser Cys Gln Asn Leu Asn His Leu Lys
115 120 125
Ala Gly Leu Gln Ala Phe Phe Gln Val Gln Glu Cys Asn Lys Ser Ser
130 135 140
Ser Lys Asp Asn Ile Arg Gly Lys His Val Arg His Tyr Tyr Ile Ala
145 150 155 160
Ala Glu Glu Ile Ile Trp Asn Tyr Ala Pro Ser Gly Ile Asp Ile Phe
165 170 175
Thr Lys Glu Asn Leu Thr Ala Pro Gly Ser Asp Ser Ala Val Phe Phe
180 185 190
Glu Gln Gly Thr Thr Arg Ile Gly Gly Ser Tyr Lys Lys Leu Val Tyr
195 200 205
Arg Glu Tyr Thr Asp Ala Ser Phe Thr Asn Arg Lys Glu Arg Gly Pro
210 215 220
Glu Glu Glu His Leu Gly Ile Leu Gly Pro Val Ile Trp Ala Glu Val

```

225	230										235										240																																																																																																																																																		
Gly	Asp	Thr	Ile	Arg	Val	Thr	Phe	His	Asn	Lys	Gly	Ala	Tyr	Pro	Leu	245	250	255	260	265	270	275	280	285	290	295	300	305	310	315	320	325	330	335	340	345	350	355	360	365	370	375	380	385	390	395	400	405	410	415	420	425	430	435	440	445	450	455	460	465	470	475	480	485	490	495	500	505	510	515	520	525	530	535	540	545	550	555	560	565	570	575	580	585	590	595	600	605	610	615	620	625	630	635	640	645	650	655	660	665	670	675	680	685	690	695	700	705	710	715	720	725	730	735	740	745	750	755	760	765	770	775	780	785	790	795	800	805	810	815	820	825	830	835	840	845	850	855	860	865	870	875	880	885	890	895	900	905	910	915	920	925	930	935	940	945	950	955	960	965	970	975	980	985	990	995	1000
Ser	Ile	Glu	Pro	Thr	Gly	Val	Arg	Phe	Asn	Lys	Asn	Asn	Glu	Pro	Thr	245	250	255	260	265	270	275	280	285	290	295	300	305	310	315	320	325	330	335	340	345	350	355	360	365	370	375	380	385	390	395	400	405	410	415	420	425	430	435	440	445	450	455	460	465	470	475	480	485	490	495	500	505	510	515	520	525	530	535	540	545	550	555	560	565	570	575	580	585	590	595	600	605	610	615	620	625	630	635	640	645	650	655	660	665	670	675	680	685	690	695	700	705	710	715	720	725	730	735	740	745	750	755	760	765	770	775	780	785	790	795	800	805	810	815	820	825	830	835	840	845	850	855	860	865	870	875	880	885	890	895	900	905	910	915	920	925	930	935	940	945	950	955	960	965	970	975	980	985	990	995	1000
Tyr	Tyr	Ser	Pro	Asn	Tyr	Asn	Pro	Gln	Ser	Arg	Ser	Val	Pro	Pro	Ser	245	250	255	260	265	270	275	280	285	290	295	300	305	310	315	320	325	330	335	340	345	350	355	360	365	370	375	380	385	390	395	400	405	410	415	420	425	430	435	440	445	450	455	460	465	470	475	480	485	490	495	500	505	510	515	520	525	530	535	540	545	550	555	560	565	570	575	580	585	590	595	600	605	610	615	620	625	630	635	640	645	650	655	660	665	670	675	680	685	690	695	700	705	710	715	720	725	730	735	740	745	750	755	760	765	770	775	780	785	790	795	800	805	810	815	820	825	830	835	840	845	850	855	860	865	870	875	880	885	890	895	900	905	910	915	920	925	930	935	940	945	950	955	960	965	970	975	980	985	990	995	1000
Ala	Ser	His	Val	Ala	Pro	Thr	Glu	Thr	Phe	Thr	Tyr	Glu	Trp																																																																																																																																																										

690	695	700
Val Phe Asp Glu Asn Glu Ser Trp Tyr Leu Asp Asn Ile Lys Thr		
705	710	715
Tyr Ser Asp His Pro Glu Lys Val Asn Lys Asp Asp Glu Glu Phe Ile		
	725	730
Glu Ser Asn Lys Met His Ala Ile Asn Gly Arg Met Phe Gly Asn Leu		
	740	745
Gln Gly Leu Thr Met His Val Gly Asp Glu Val Asn Trp Tyr Leu Met		
	755	760
Gly Met Gly Asn Glu Ile Asp Leu His Thr Val His Phe His Gly His		
	770	775
Ser Phe Gln Tyr Lys His Arg Gly Val Tyr Ser Ser Asp Val Phe Asp		
785	790	795
Ile Phe Pro Gly Thr Tyr Gln Thr Leu Glu Met Phe Pro Arg Thr Pro		
	805	810
Gly Ile Trp Leu Leu His Cys His Val Thr Asp His Ile His Ala Gly		
	820	825
Met Glu Thr Thr Tyr Thr Val Leu Gln Asn Glu Gly Glu Tyr Pro Asp		
	835	840
Thr Lys Ser Gly		
850		

<210> 58

<211> 3321

<212> DNA

<213> Homo sapiens

<400> 58

atgaagattt	tgatacttgg	tattttttctg	ttttttatgta	gtacccaccagc	ctgggcgaaa	60
gaaagcattt	attacattgg	aattattgaa	acgactttggg	attatgcttc	tgaccatggg	120
gaaaaaagac	ttattttctgt	tgacacggaa	catttccaata	tctatcttca	aaatggccca	180
gatagaattg	ggagactata	taagaaggcc	ctttatcttc	agtacacaga	tgaacctttt	240
aggacaacta	tagaaaaacc	ggctctggctt	gggttttttag	gccctattat	caaatgctgaa	300
actggagata	aagttttatgt	acacttaaaa	aaccttgcct	ctaggcccta	caacctttcat	360
tcacatggaa	taacttacta	taaggaacat	gagggggcca	tctaccctga	taacaccaca	420
gatttttcaa	gagcagatga	caaagtatal	ccaggagagc	agtatacata	catgttgctt	480
gccactgaag	aacaaagtcc	tggggaaagg	gatggcaatt	gtgtgactag	gattttaccat	540
toccacattg	atgtctccaaa	agatatgtgc	tcaggactca	tcggaccttt	aataatctgt	600
aaaaaagatt	ctctagataa	agaaaaagaa	aaacatatgt	accgagaatt	tggtgtgatg	660
tttttctgtg	tggatgaaaa	tttcagctgg	tacctagaag	acaacattaa	aacctactgc	720
tcagaaccag	agaaagtgtg	caaagacaac	gaagacttcc	aggagagtaa	cagaatgtat	780
ctctgtgaatg	gatacaactt	tgggaagtct	ccaggactct	ccatgtgtgc	tgaagacaga	840
gtaaaatggg	accttttttg	tatgggtaat	gaagttgatg	tgacgcagc	cttctttcac	900
gggcaagcac	tgactaacaa	gaactaccgt	altgacacaa	tcaacctctt	tctgtctacc	960
ctgttttgatg	cttatatggg	ggcccaagaac	cctggagaat	ggatgctcag	cgctgaggaa	1020
ctaaaccatc	tgaagccggg	tttgcaagcc	tttttccagg	tcaggagtg	taacaagtct	1080
tcatacaagg	ataatatccg	tgggaagcat	gttagacact	acacatttgc	cgctgaggaa	1140
atcatctgga	actatgtctc	ctctggtata	gacatcttca	ctaaagaaaa	cttaacagca	1200
ccgtggaagt	actcagcggg	gttttttgaa	caaggatcca	caagaattgg	aggctcttat	1260
aaaaagctgg	tttatctgtga	gtacacagat	gcctctctca	caaatcgaaa	ggagagaggg	1320
ctcgaagaag	agactatttg	catcctgggt	cctgtcattt	gggcagaggt	gggagacacc	1380
atcagagtaa	cccttccataa	caaaggagca	tattcccttc	gtattgagcc	gattgggggt	1440
agatttcaata	agacaacaga	gggcacatac	tattccccaa	attacaacc	ccagagcaga	1500
agtggtcctc	cttcagcctc	ccatgtggca	ccacacagaa	cattcaccta	tgaattggat	1560
ctccccaaag	aagtagagcc	cactaatgca	gatcctgtgt	gtctagctaa	gatgtattat	1620
ctcgtctgtg	atccccataa	agatatattc	actgggctta	ttgggccaat	gaaatatg	1680
aagaaaggaa	gtttacatgc	aaatgggaga	cagaaagatg	tagacaagga	attctatttg	1740

```

tttccacag tatttgatga gaatgagagt ttactoctgg aagataatat tagaatgttt 1800
acaactgcac ctgatcagggt ggataaggaa gatgaagact ttcaggaact taataaaatg 1860
cactccatga atggattccat gtatggggaat cagccgggtc tcactatgtg caaaggagat 1920
tcggtcgtgt ggtacttatt cagcgccgga aatgaggccg atgtacatgg aataactatt 1980
tcaggaaaca catatctgtg gagaggagaa cggagagaca cagcaaacct cttccctcaa 2040
acaagtccta cgtccacat gtggcctgac acagagggga ctttaagtgt tgaatgcctt 2100
acaactgac attacacagg cggcatgaag caaaaatata ctgtgaacca atgcaggcgg 2160
cagtctgggg attcacacct ctacctggga gagaggacat actatatcgc agcagtgagg 2220
gtggaatggg tttaatcccc acaaaaggag tgggaaaagg agctgcatca tttacaagag 2280
cagaatgttt caaatgcatt tttagataag ggagagtttt acataggctc aaagtacaag 2340
aaagtgtgt atcggcagta tactgatagc acattccgtg ttccagtggg gagaaaaagt 2400
gaagaagaac atctgggaat tctaggtcca caactccatg cagatgttgg agacaaagtc 2460
aaaattatct ttaaaaacat ggccacaagg cctactcaa tacatgcccc tggggtacaa 2520
acagagagtt ctacagttac tccaacatta ccaggtgaaa ctctcactta cgtatggaaa 2580
atccccagaa gatctggagc tggaaacagag ggcactctac agtgatttaa ttggccccc 2640
tcaactgtgg atcaagttaa ggacactctac agtgatttaa ttggccccc 2700
cgaagacctt actgaaagt attcaatccc agnagggaagc tggaaattgc cctctgttt 2760
ctagttttt atgagaatga atcttggtac ttagatgaca acatcaaaac atactctgt 2820
caccccgaga aagtaaacaa agatgatgag gaattccatg aaagcaataa aatgcattgt 2880
attaatggaa gaatgtttgg aaacctacaa ggccctacaa tgcacgtggg agatgaagtc 2940
aactgtgata tgatgggaat gggcaatgaa atagacttac acactgtaca ttttcacggc 3000
catagctccc aatcacagca caggggagtt tatagttctg atgtcttga cttttccct 3060
ggaacatacc aaacctaga aatgtttcca agaaccactg gaatttgggt actccaatgc 3120
catgtaccgc accaattcca tgcgtggaat gaaaccactt acaccgttct acaaaatgaa 3180
gacaccaaat ctggtcgaat gaaataaatt ggtgataagt ggaaaaaaga gaaaaacca 3240
tgattcataa caatgtatgt gaaagtgtaa aatagaatgt tactttgaaa tgactataa 3300
cattaaaaa gactggagca t 3321

```

<210> 59

<211> 1065

<212> FRT

<213> Homo sapiens

<400> 59

```

Met Lys Ile Leu Ile Leu Gly Ile Phe Leu Phe Leu Cys Ser Thr Pro
  1             5             10             15
Ala Trp Ala Lys Glu Lys His Tyr Tyr Ile Gly Ile Ile Glu Thr Thr
  20             25             30
Trp Asp Tyr Ala Ser Asp His Gly Glu Lys Lys Leu Ile Ser Val Asp
  35             40             45
Thr Glu His Ser Asn Ile Tyr Leu Gln Asn Gly Pro Asp Arg Ile Gly
  50             55             60
Arg Leu Tyr Lys Lys Ala Leu Tyr Leu Gln Tyr Thr Asp Glu Thr Phe
  65             70             75             80
Arg Thr Thr Ile Glu Lys Pro Val Trp Leu Gly Phe Leu Gly Pro Ile
  85             90             95
Ile Lys Ala Glu Thr Gly Asp Lys Val Tyr Val His Leu Lys Asn Leu
 100             105             110
Ala Ser Arg Pro Tyr Thr Phe His Ser His Gly Ile Thr Tyr Tyr Lys
 115             120             125
Glu His Glu Gly Ala Ile Tyr Pro Asp Asn Thr Thr Asp Phe Gln Arg
 130             135             140
Ala Asp Asp Lys Val Tyr Pro Gly Glu Gln Tyr Thr Tyr Met Leu Leu
 145             150             155             160
Ala Thr Glu Glu Gln Ser Pro Gly Glu Gly Asp Gly Asn Cys Val Thr
 165             170             175
Arg Ile Tyr His Ser His Ile Asp Ala Pro Lys Asp Ile Ala Ser Gly
 180             185             190

```

Leu Ile Gly Pro Leu Ile Ile Cys Lys Lys Asp Ser Leu Asp Lys Glu
 195 200 205
 Lys Glu Lys His Ile Asp Arg Glu Phe Val Val Met Phe Ser Val Val
 210 215 220
 Asp Glu Asn Phe Ser Trp Tyr Leu Glu Asp Asn Ile Lys Thr Tyr Cys
 225 230 235 240
 Ser Glu Pro Glu Lys Val Asp Lys Asp Asn Glu Asp Phe Gln Glu Ser
 245 250 255
 Asn Arg Met Tyr Ser Val Asn Gly Tyr Thr Phe Gly Ser Leu Pro Gly
 260 265 270
 Leu Ser Met Cys Ala Glu Asp Arg Val Lys Trp Tyr Leu Phe Gly Met
 275 280 285
 Gly Asn Glu Val Asp Val His Ala Ala Phe Phe His Gly Gln Ala Leu
 290 295 300
 Thr Asn Lys Asn Tyr Arg Ile Asp Thr Ile Asn Leu Phe Pro Ala Thr
 305 310 315 320
 Leu Phe Asp Ala Tyr Met Val Ala Gln Asn Pro Gly Glu Trp Met Leu
 325 330 335
 Ser Cys Gln Asn Leu Asn His Leu Lys Ala Gly Leu Gln Ala Phe Phe
 340 345 350
 Gln Val Gln Glu Cys Asn Lys Ser Ser Ser Lys Asp Asn Ile Arg Gly
 355 360 365
 Lys His Val Arg His Tyr Tyr Ile Ala Ala Glu Glu Ile Ile Trp Asn
 370 375 380
 Tyr Ala Pro Ser Gly Ile Asp Ile Phe Thr Lys Glu Asn Leu Thr Ala
 385 390 395 400
 Pro Gly Ser Asp Ser Ala Val Phe Phe Glu Gln Gly Thr Thr Arg Ile
 405 410 415
 Gly Gly Ser Tyr Lys Lys Leu Val Tyr Arg Glu Tyr Thr Asp Ala Ser
 420 425 430
 Phe Thr Asn Arg Lys Glu Arg Gly Pro Glu Glu Glu His Leu Gly Ile
 435 440 445
 Leu Gly Pro Val Ile Trp Ala Glu Val Gly Asp Thr Ile Arg Val Thr
 450 455 460
 Phe His Asn Lys Gly Ala Tyr Pro Leu Ser Ile Glu Pro Ile Gly Val
 465 470 475 480
 Arg Phe Asn Lys Asn Asn Glu Gly Thr Tyr Tyr Ser Pro Asn Tyr Asn
 485 490 495
 Pro Gln Ser Arg Ser Val Pro Pro Ser Ala Ser His Val Ala Pro Thr
 500 505 510
 Glu Thr Phe Thr Tyr Glu Trp Thr Val Pro Lys Glu Val Gly Pro Thr
 515 520 525
 Asn Ala Asp Pro Val Cys Leu Ala Lys Met Tyr Tyr Ser Ala Val Asp
 530 535 540
 Pro Thr Lys Asp Ile Phe Thr Gly Leu Ile Gly Pro Met Lys Ile Cys
 545 550 555 560
 Lys Lys Gly Ser Leu His Ala Asn Gly Arg Gln Lys Asp Val Asp Lys
 565 570 575
 Glu Phe Tyr Leu Phe Pro Thr Val Phe Asp Glu Asn Glu Ser Leu Leu
 580 585 590
 Leu Glu Asp Asn Ile Arg Met Phe Thr Thr Ala Pro Asp Gln Val Asp
 595 600 605
 Lys Glu Asp Glu Asp Phe Gln Glu Ser Asn Lys Met His Ser Met Asn
 610 615 620
 Gly Phe Met Tyr Gly Asn Gln Pro Gly Leu Thr Met Cys Lys Gly Asp
 625 630 635 640
 Ser Val Val Trp Tyr Leu Phe Ser Ala Gly Asn Glu Ala Asp Val His
 645 650 655

Gly Ile Tyr Phe Ser Gly Asn Thr Tyr Leu Trp Arg Gly Glu Arg Arg
 660 665 670
 Asp Thr Ala Asn Leu Phe Pro Gln Thr Ser Leu Thr Leu His Met Trp
 675 680 685
 Pro Asp Thr Glu Gly Thr Phe Asn Val Glu Cys Leu Thr Thr Asp His
 690 700
 Tyr Thr Gly Gly Met Lys Gln Lys Tyr Thr Val Asn Gln Cys Arg Arg
 705 710 715 720
 Gln Ser Glu Asp Ser Thr Phe Tyr Leu Gly Glu Arg Thr Tyr Tyr Ile
 725 730 735
 Ala Ala Val Glu Val Glu Trp Asp Tyr Ser Pro Gln Arg Glu Trp Glu
 740 745 750
 Lys Glu Leu His His Leu Gln Glu Gln Asn Val Ser Asn Ala Phe Leu
 755 760 765
 Asp Lys Gly Glu Phe Tyr Ile Gly Ser Lys Tyr Lys Val Val Tyr
 770 775 780
 Arg Gln Tyr Thr Asp Ser Thr Phe Arg Val Pro Val Glu Arg Lys Ala
 785 790 795 800
 Glu Glu Glu His Leu Gly Ile Leu Gly Pro Gln Leu His Ala Asp Val
 805 810 815
 Gly Asp Lys Val Lys Ile Ile Phe Lys Asn Met Ala Thr Arg Pro Tyr
 820 825 830
 Ser Ile His Ala His Gly Val Gln Thr Glu Ser Ser Thr Val Thr Pro
 835 840 845
 Thr Leu Pro Gly Glu Thr Leu Thr Tyr Val Trp Lys Ile Pro Glu Arg
 850 855 860
 Ser Gly Ala Gly Thr Glu Asp Ser Ala Cys Ile Pro Trp Ala Tyr Tyr
 865 870 875 880
 Ser Thr Val Asp Gln Val Lys Asp Leu Tyr Ser Gly Leu Ile Gly Pro
 885 890 895
 Leu Ile Val Cys Arg Arg Pro Tyr Leu Lys Val Phe Asn Pro Arg Arg
 900 905 910
 Lys Leu Glu Phe Ala Leu Leu Phe Leu Val Phe Asp Glu Asn Glu Ser
 915 920 925
 Trp Tyr Leu Asp Asp Asn Ile Lys Thr Tyr Ser Asp His Pro Glu Lys
 930 935 940
 Val Asn Lys Asp Asp Glu Glu Phe Ile Glu Ser Asn Lys Met His Ala
 945 950 955 960
 Ile Asn Gly Arg Met Phe Gly Asn Leu Gln Gly Leu Thr Met His Val
 965 970 975
 Gly Asp Glu Val Asn Trp Tyr Leu Met Gly Met Gly Asn Glu Ile Asp
 980 985 990
 Leu His Thr Val His Phe His Gly His Ser Phe Gln Tyr Lys His Arg
 995 1000 1005
 Gly Val Tyr Ser Ser Asp Val Phe Asp Ile Phe Pro Gly Thr Tyr Gln
 1010 1015 1020
 Thr Leu Glu Met Phe Pro Arg Thr Pro Gly Ile Trp Leu Leu His Cys
 1025 1030 1035 1040
 His Val Thr Asp His Ile His Ala Gly Met Glu Thr Thr Tyr Thr Val
 1045 1050 1055
 Leu Gln Asn Glu Asp Thr Lys Ser Gly
 1060 1065

<210> 60

<211> 3881

<212> DNA

<213> Homo sapiens

<220>
 <221> misc_feature
 <222> (1)...(3881)
 <223> n = A,T,C or G

<400> 60

```

agaaagtctt atttataaaa ttgcotgctc ctgattttac ttcatctctt ctcaggctcc 60
aagaagtgga aaaaaatgaa gatcttgata tctgttattt ttctgttttt atgtagtacc 120
ccagcctggg cgaaagaaaa gcattattac attggaatta ttgaaacgac ttgggattat 180
gcctctgacc atggggaaaa gaaacttatt tctgttgaca cggaacattc caatatctat 240
cttcaaaatg gccacgatag aattgggaga ctatataaga aggcctctta tcttcagtac 300
acagatgaaa cctttaggac aactatagaa aaaccggtct ggcttggggt tttaggccct 360
attatcaaaq ctgaaactgg agataaagtt tatgtacact taaaaaacct tgcctctagg 420
cctcacacct ttcaattaca tggaaatact tactataagg aacatgaggg ggccatctac 480
cctgataaca ccacagattt tcaaaagaca gatgacaaag tatatccagg agagcagtat 540
acatacatgt tgccttgccac tgaagaacaa agtccctggg aaggagatgg caattgtgtg 600
actaggattt accattccca caltgatgct ccaaaagata ttgctcagg actcatcgga 660
ccttlaataa tctgtaaaaa agattctcta gataaagaaa aagaaaaaca tattgaccga 720
gaatttgtag tgatgttttc tgtggtggat gaaaatttca gctggtacct agagagacca 780
attaaaaact actgctcaga accagagaaa gttgacaaag acaacgaaga cttccaggag 840
agtaacagca tglattctgt gaatggatac acttttgtaa gtctcccagg actctccatg 900
tgtgctgaag acagagataa atggtacctt tttggtatgg gtaatgaagt tgatgtgca 960
gcagcttctt ttacggggca agcactgaat acaagaactc accgtattga caacaactac 1020
ctctttctgt ctaccctgtt tgatgcttat atggtggccc agaacctcgt agaatggatg 1080
ctcagctgtc agaattctaa ccatctgaaa gcgcttttgc agcctttttt ccaggctcag 1140
gagtgtaaca agtcttcac c aaagataat atccgtggga agcatgttag acactactac 1200
attgcgcctg aggaatacat ctggaactat gctccctctg gtatagacat ctccaataaa 1260
gaaaacttaa cagcacctgg aagtgaacta gcggtgtttt ttgaacaagg taccacaaga 1320
attggaggct cttataaaaa gctggtttat cgtgagtaca cagatgcctc cttcacaagt 1380
cgaaaggaga gaggccttga agaagagcat ctltggcatcc tgggtcctgt catltgggca 1440
gaggtgggag acaccatcag agtaaccttc catacaaaag gagcatatcc cctcagttat 1500
gagccgatgt ggggtgagatt caataagaac aacgagggca catactatc cccaattac 1560
aaccococaga gcgaagtggt gctccttcca gctcccatc tggcaccacc agaaacattc 1620
accatagaat ggactgtccc caaagaagta ggacccacta atgcagatcc tgtgtgtcta 1680
gctaagatgt attattctgc tgtggatccc actaaagata tattcactgg gcttattggg 1740
coaatgaaaa tatgcaagaa aggaagttaa catgcaaatg ggagacagaa agatgtagac 1800
aaggaattct atttgtttcc tacagtattt gatgagaatg agagtttact cctgggaagt 1860
aatattagaa tghtttacaac tgcaactgat caggtggata aggaagtaga agactttcag 1920
gaatctaata aaatgcactc catgaatgga ttcatgtatg ggaatcagcc ggggtctact 1980
atgtgcaaaq gagattcggg cgtgtggtac ttattcagcg cgggaatga ggcctgatga 2040
catggaatat acttttcagg aaacacatat cttgtggagag gagaacggag agacaagca 2100
aaactcttcc cctaaacaag tcttacgctc cacatgtggc ctgacacaga ggggactctt 2160
aatgttgaa tctctcaaac tgatcattac tgaagcaaaa atatactgtg 2220
aaccatgca gggcgagctc tgaggatttc acctctacc tgggagagag gacatactat 2280
atcgacagca tggagtggtg atgggattat tccccacaaa gggagtggga aaagagctg 2340
catcatctac aagacagaaa tgtttcaaat gcaattttag ataaaggaga gttttacata 2400
ggctcaaagt acaagaagtg tgtgtatcgg cagtatactg atagcacatt ccgtgttcca 2460
gtggagagaa aagctgaaga agaacatctg ggaattctag gtccacaact ctatgcagat 2520
gttgagacaa aagtcaaaat tatctttaaa aaactggcca caaggcccta ctcaatcact 2580
gccatggggg tacaacacaga gagttctaca gttactccaa cattaccagg tgaaactctc 2640
acttaagtat ggaataatccc agaaagatct ggagctggaa cagaggattc tgctgtat 2700
ccatgggctt attattcaac tgtggtcaa gttaaaggacc tctacagtggt ttaattggc 2760
ccctgatgtt tttgtcgaag accttacttg aaagtattca atcccagaag gaagctggaa 2820
tttgcccttc tttgtctagt ttttgatgag aatgaatctt ggtacttga tgacaacatc 2880
aaaaataact ctgatcacc cagaaaagta aacaagatg atgaggaatt catagaagac 2940
aataaaatgc atgctattaa tggagaagtg tttggaacc tacaaggcct cacaatgcac 3000
gtgggagatg aagtcaactg gtatctgatg ggaatgggca atgaaataga cttacacact 3060

```

```

gtacattttc acggccatag cttccaatac aagcacaggg gagtttatag ttctgatgtc 3120
tttgacattt tccctgggaa ataccaaacc ctagaatgt ttccaagaac acctggaatt 3180
tggttactcc actgccatgt gaccgaccac attcatgctg gaatgggaac cacttacacc 3240
gtttctacaaa atgaagcatc ttctgagact cacaggagaa tatggaatgt gatctaccac 3300
atcacagtca gtgtgattat ttatttccaa atatctacca aggaatgacc aggagaataa 3360
gatcctccga tgttcgcaat ggtgtggtgt caggaggctg cctcttagac aatctccaga 3420
tggtactgtg atgtgagttt gaaaagagt tcttgaagta ccacatctgg gagacatgcc 3480
actagctgag cttcccaaaa gtctaccaag agctgaggaa ttgtatcttc atccttgacc 3540
aaagcaoctt aaaaacagta aaaggagcct ctatatccca gataaatata gcactgataa 3600
agcgacagct gggcgtgaat atcacagcat cacatggggc acgcaatcaa taatttacca 3660
ttagcattag cgccatggca aaagcaaat tagacatttt taaaaggaaa cagattctag 3720
gagtgataaaa taaagtgtct acagatatag tcttttaatt agtactttta tcttataaac 3780
tgattatgta tgtctgtttt ttcaataaac aaacaaatga aaaaaaaaaa aaaaaaatgg 3840
cggccgcaag cttattanc cttagtggag gttaatttta a 3881

```

<210> 61

<211> 1090

<212> PRT

<213> Homo sapiens

<400> 61

```

Met Lys Ile Leu Ile Leu Gly Ile Phe Leu Phe Leu Cys Ser Thr Pro
1 5 10 15
Ala Trp Ala Lys Glu Lys His Tyr Tyr Ile Gly Ile Ile Glu Thr Thr
20 25 30
Trp Asp Tyr Ala Ser Asp His Gly Glu Lys Lys Leu Ile Ser Val Asp
35 40 45
Thr Glu His Ser Asn Ile Tyr Leu Gln Asn Gly Pro Asp Arg Ile Gly
50 55 60
Arg Leu Tyr Lys Lys Ala Leu Tyr Leu Gln Tyr Thr Asp Glu Thr Phe
65 70 75 80
Arg Thr Thr Ile Glu Lys Pro Val Trp Leu Gly Phe Leu Gly Pro Ile
85 90 95
Ile Lys Ala Glu Thr Gly Asp Lys Val Tyr Val His Leu Lys Asn Leu
100 105 110
Ala Ser Arg Pro Tyr Thr Phe His Ser His Gly Ile Thr Tyr Tyr Lys
115 120 125
Glu His Glu Gly Ala Ile Tyr Pro Asp Asn Thr Thr Asp Phe Gln Arg
130 135 140
Ala Asp Asp Lys Val Tyr Pro Gly Glu Gln Tyr Thr Tyr Met Leu Leu
145 150 155 160
Ala Thr Glu Glu Gln Ser Pro Gly Glu Gly Asp Gly Asn Cys Val Thr
165 170 175
Arg Ile Tyr His Ser His Ile Asp Ala Pro Lys Asp Ile Ala Ser Gly
180 185 190
Leu Ile Gly Pro Leu Ile Ile Cys Lys Lys Asp Ser Leu Asp Lys Glu
195 200 205
Lys Glu Lys His Ile Asp Arg Glu Phe Val Val Met Phe Ser Val Val
210 215 220
Asp Glu Asn Phe Ser Trp Tyr Leu Glu Asp Asn Ile Lys Thr Tyr Cys
225 230 235 240
Ser Glu Pro Glu Lys Val Asp Lys Asp Asn Glu Asp Phe Gln Glu Ser
245 250 255
Asn Arg Met Tyr Ser Val Asn Gly Tyr Thr Phe Gly Ser Leu Pro Gly
260 265 270
Leu Ser Met Cys Ala Glu Asp Arg Val Lys Trp Tyr Leu Phe Gly Met
275 280 285
Gly Asn Glu Val Asp Val His Ala Ala Phe Phe His Gly Gln Ala Leu

```

290		295		300	
Thr Asn Lys Asn Tyr Arg Ile Asp Thr Ile Asn Leu Phe Pro Ala Thr					
305		310		315	320
Leu Phe Asp Ala Tyr Met Val Ala Gln Asn Pro Gly Glu Trp Met Leu					
	325		330		335
Ser Cys Gln Asn Leu Asn His Leu Lys Ala Gly Leu Gln Ala Phe Phe					
	340		345		350
Gln Val Gln Glu Cys Asn Lys Ser Ser Ser Lys Asp Asn Ile Arg Gly					
	355		360		365
Lys His Val Arg His Tyr Tyr Ile Ala Ala Glu Glu Ile Ile Trp Asn					
	370		375		380
Tyr Ala Pro Ser Gly Ile Asp Ile Phe Thr Lys Glu Asn Leu Thr Ala					
385		390		395	400
Pro Gly Ser Asp Ser Ala Val Phe Phe Glu Gln Gly Thr Thr Arg Ile					
	405		410		415
Gly Gly Ser Tyr Lys Lys Leu Val Tyr Arg Glu Tyr Thr Asp Ala Ser					
	420		425		430
Phe Thr Asn Arg Lys Glu Arg Gly Pro Glu Glu Glu His Leu Gly Ile					
	435		440		445
Leu Gly Pro Val Ile Trp Ala Glu Val Gly Asp Thr Ile Arg Val Thr					
	450		455		460
Phe His Asn Lys Gly Ala Tyr Pro Leu Ser Ile Glu Pro Ile Gly Val					
465		470		475	480
Arg Phe Asn Lys Asn Asn Glu Gly Thr Tyr Ser Pro Asn Tyr Asn					
	485		490		495
Pro Gln Ser Arg Ser Val Pro Pro Ser Ala Ser His Val Ala Pro Thr					
	500		505		510
Glu Thr Phe Thr Tyr Glu Trp Thr Val Pro Lys Glu Val Gly Pro Thr					
	515		520		525
Asn Ala Asp Pro Val Cys Leu Ala Lys Met Tyr Tyr Ser Ala Val Asp					
	530		535		540
Pro Thr Lys Asp Ile Phe Thr Gly Leu Ile Gly Pro Met Lys Ile Cys					
545		550		555	560
Lys Lys Gly Ser Leu His Ala Asn Gly Arg Gln Lys Asp Val Asp Lys					
	565		570		575
Glu Phe Tyr Leu Phe Pro Thr Val Phe Asp Glu Asn Glu Ser Leu Leu					
	580		585		590
Leu Glu Asp Asn Ile Arg Met Phe Thr Thr Ala Pro Asp Gln Val Asp					
	595		600		605
Lys Glu Asp Glu Asp Phe Gln Glu Ser Asn Lys Met His Ser Met Asn					
	610		615		620
Gly Phe Met Tyr Gly Asn Gln Pro Gly Leu Thr Met Cys Lys Gly Asp					
625		630		635	640
Ser Val Val Trp Tyr Leu Phe Ser Ala Gly Asn Glu Ala Asp Val His					
	645		650		655
Gly Ile Tyr Phe Ser Gly Asn Thr Tyr Leu Trp Arg Gly Glu Arg Arg					
	660		665		670
Asp Thr Ala Asn Leu Phe Pro Gln Thr Ser Leu Thr Leu His Met Trp					
	675		680		685
Pro Asp Thr Glu Gly Thr Phe Asn Val Glu Cys Leu Thr Thr Asp His					
	690		695		700
Tyr Thr Gly Gly Met Lys Gln Lys Tyr Thr Val Asn Gln Cys Arg Arg					
705		710		715	720
Gln Ser Glu Asp Ser Thr Phe Tyr Leu Gly Glu Arg Thr Tyr Tyr Ile					
	725		730		735
Ala Ala Val Glu Val Glu Trp Asp Tyr Ser Pro Gln Arg Glu Trp Glu					
	740		745		750
Lys Glu Leu His His Leu Gln Glu Gln Asn Val Ser Asn Ala Phe Leu					

[illegible]

```
<210> 62
<211> 969
<212> DNA
<213> Homo sapiens
```

400> 62							
agcttctgggg	ttgtccctgg	acctgtcttg	gttcagaaac	ctgacgaccc	ggcgacggcg	60	
acgtctctctt	tgactaaaag	acagttgtcca	ctgtcccaagc	ctagagtgctc	acggggcagac	120	
ctctcccgccg	ggccacactt	cccaactctt	ctgcgaactc	gaaataactc	cgatcgggaaa	180	
ctctctgagga	atctctcaaa	gtgctctgggg	tgaattgtgt	ctgaggagaa	attgtctgtg	240	
ctctcagctga	caagcagcga	gtggagatcta	acacaggagga	agacactctt	tactacaaaa	300	
ctctccaccac	cgctgcaccc	acagagatta	acttcaaaatg	tgggggagag	ttttgaggacg	360	
agactctatga	ttggagaccc	ttctaagacac	tactgaatacg	cggagagatcg	aataataatcg	420	


```

tctgtgagca gaagctcctg aagggagagg gccccaaagc ctcgtggacc agagaactga 480
ccaacgatgg ggaactgac ctagcatga cggcggtatga cgttgtgtgc accagggtct 540
acgtcgcaga gtgagtggcc acaggtagaa ccgcggccga agcccaaccac tggccatgct 600
caccgcccgt cttcactgcc cctctcgtcc caccgccctc ttctaggata gcgctccct 660
taccaccagtc actctctggg gtcactggga tgcctcttgc agggctcttg tttctttgac 720
ctctctcttc ctcccctaca ccaacaaaga ggaatggctg caagagccca gatcacccat 780
tccgggttca tccccgcct ccccaagtea gcagtcctag ccccaaaaca gccagagaca 840
gggtctctct aaaggggact tgagggccctg agcaggaagaa actggccctc tagctctctac 900
cctttgtccc tgtagcctat acagtttaga atattttatt gttaatttta ttaaaatgct 960
ttaaaaaaa

```

<210> 63

<211> 138

<212> PRT

<213> Homo sapiens

<400> 63

```

Met Pro Asn Phe Ser Gly Asn Trp Lys Ile Ile Arg Ser Glu Asn Phe
1 5 10 15
Glu Glu Leu Leu Lys Val Leu Gly Val Asn Val Met Leu Arg Lys Ile
20 25 30
Ala Val Ala Ala Ala Ser Lys Pro Ala Val Glu Ile Lys Gln Glu Gly
35 40 45
Asp Thr Phe Tyr Ile Lys Thr Ser Thr Thr Val Arg Thr Thr Glu Ile
50 55 60
Asn Phe Lys Val Gly Glu Glu Phe Glu Glu Gln Thr Val Asp Gly Arg
65 70 75 80
Pro Cys Lys Ser Leu Val Lys Trp Glu Ser Glu Asn Lys Met Val Cys
85 90 95
Glu Gln Lys Leu Leu Lys Gly Glu Gly Pro Lys Thr Ser Thr Thr Arg
100 105 110
Glu Leu Thr Asn Asp Gly Glu Leu Ile Leu Thr Met Thr Ala Asp Asp
115 120 125
Val Val Cys Thr Arg Val Tyr Val Arg Glu
130 135

```

<210> 64

<211> 927

<212> DNA

<213> Homo sapiens

<400> 64

```

ggaagttag gttactgtc ttaattttcc aaagctgtaa tcattatttt cattctcaaa 60
gtgatggcct tgtgttttgc tctctctctc cagggccaga ctgagccacg gttgatttca 120
gcgggacacc aatagaactcc acagcagctc caggagccca gacaccggcg gccagaagca 180
aggctaggag ctgctgcagc catgtcggcc ctacagctcc tcattctggg cctgctcaag 240
cgactgcacc ctgcccagctg tcagcaaggc ctggggaacc ttcagccctg gatcgaggc 300
cttatcgccg ttgcccgtgtt cctgtctctc gttgcaatcg cctttgcagt caaccacttc 360
tggtgcagg aggagccgga gcctgcacac atgactctga ccgtogaaaa caaggcagat 420
ggagtctctg ttggaacaga ttgaaggtac tcttcgatgg cggccagtgt caggtccagt 480
gagcatgaga atgcctatga gaattgtgcc gaggaggaag gcaaggtccg cagcaccctg 540
atgtaacctt ctctgtggct ccaaccccaa gactcccagg caatagggat ggatgtccag 600
tgtactaccac caagccccct cctctcttct gtggaatctg caatagtggt ctgactccct 660
ccagcccat gccggcccta ccgccccttg aagtatagcc agccaaggtt ggagctcaga 720
cgtgtctatg gttggggctc ggctgtggcc ctgggggtct ctgctcagct cagaagagcc 780
ttctggagag gacagtcagc tgagcacctc coactctgct cacacgtcct tcccataaac 840
tatggaatg gccctaattt ctgtgaaata aagacttttt gtatttctgg ggtgaggct 900

```

cagcaacagc cctcaggct tccaaaa

927

<210> 65

<211> 114

<212> PRT

<213> Homo sapiens

<400> 65

```

Met Ser Ala Leu Ser Leu Leu Ile Leu Gly Leu Leu Thr Ala Val Pro
 1          5          10          15
Pro Ala Ser Cys Gln Gln Gly Leu Gly Asn Leu Gln Pro Trp Met Gln
 20          25          30
Gly Leu Ile Ala Val Ala Val Phe Leu Val Ala Ile Ala Phe
 35          40          45
Ala Val Asn His Phe Trp Cys Gln Glu Glu Pro Glu Pro Ala His Met
 50          55          60
Ile Leu Thr Val Gly Asn Lys Ala Asp Gly Val Leu Val Gly Thr Asp
 65          70          75          80
Gly Arg Tyr Ser Ser Met Ala Ala Ser Phe Arg Ser Ser Glu His Glu
 85          90          95
Asn Ala Tyr Glu Asn Val Pro Glu Glu Gly Lys Val Arg Ser Thr
100          105          110
Pro Met

```

<210> 66

<211> 3641

<212> DNA

<213> Homo sapiens

<400> 66

```

ctgcactgaa gagggagagc gagagagaga ctggagacgc acagatcccc ccaaggtctc 60
ccaagcctac cgtccacagc attattgtac agagccccaa aaatcgaaac agaggaaacg 120
aacagcagtt gaacatggac gaaggaattc ctcatattgca agagagacag ttactggaaac 180
atagagattt tataggactg gactattcct ctttgtatat gtgtaaaccc aaaaggagca 240
tgaaacgaga cgacaccaag gatacctaca aattaccgca cagattaata gaaaagaaaa 300
gaagagaccc aattaatgaa tgcattgctc agctgaaaga ttactgcctc gaacatctga 360
aattgacaac tctgggacat ctggagaagc ctgtagtctt ggaattaact ttgaaacact 420
taaaagcttt aacgccttta accgagcaac agcatcagaa gataattgct ttacagaatg 480
gggagcgcat tctgaaatcg ccatttcagt ccgacttgga tgcgtccac tcgggatttc 540
aaacatgcgc caaagaagtc ttgcaatacc tctcccggtt tgagagctgg acaccaggg 600
agccgcgggt gtgccagctg atcaaccact tgcacgcgct ggccaccagc tctttgccca 660
ccccgcagct gttgactcaa caggtccctc tgagcaaaag caccggcgct cctctggccg 720
ccgggtccgc ggccgccccc tgcctggagc gcgcggggca gaagctggag cccctgcctc 780
actcgctgcc cgtcatccag cggactcagc ccagcgccga gctcgccgcc gagaacgaca 840
cggaacccga cagcgctac ggccgggaag ccgagggccc gccggaccgc gagaagagca 900
aaggcgcggg ggcgacgcgc gtcacatca agcaggagcc tcccggggag gactcgccg 960
cgcccaagag gatgaagctg gattcccgcg gcggggcgag cggcgccgag ccggggggcg 1020
gcgcggcgcg ggcgggagcc gcgctctcgg ggcccgaccc tgcgcgcgcg gccgcgctgc 1080
tgagaccoga cgcgcctctg ctacgctcgc tggtagcggt gcggcgagcg ggagggcgcg 1140
ccttcgccga gcccggggcc gcccgggccc cctctcgcct gcctctctgc ttctctcgc 1200
cttctcgagc tgcgcctac gtgcagccct tctggacaaa gagcgccgct gagaagtatc 1260
tgtaccggcg ggcgctgcc gccccgttcc cgctgtcata ccccgccatc cccgcccccg 1320
cggaacgcgc ggcaagcgcc gcccgcgctg ccgcgcgcgc ccccgctctc cctcgtctgt 1380
cctcggtgtt gtgcgccctc ccgagaaagg cggcgcgccc cgcccgagcc ctctcgccg 1440
acgagtgcc gcccttggg gcgcgcacc ccagcacc ccacgcgcg acccactgc 1500
ccttcgcccc ggccccgcag ccggggaacc cggagagctc tgctcaggaa gatccctcgc 1560

```

```

agccaggaaa ggaagctccc tgaatccttg cgtcccgagg gacggagggt caagcagagt 1620
gagaaggttaa aataccctta aggagggttca agcagagtga gaaggttaaaa tacccttaag 1680
gtcctttaagg gaggaagtgt aatagatgca ogacaggcat aaacaagaac aacaaaaaag 1740
gtgttatgttg taacattcggg gtccctgttt tgctcatccc gcaccacccc accctccaca 1800
cactaaacatc cctttctctcc cccaccagc tgtaaaagat cctatgcgaa agacactggc 1860
tccttttttt taatcccaca ataaaatttg ccccttttta ggccatgttc cattatctct 1920
taaaattgga acctaattcg agaggaagta agaaggggtc gtctctgtggc ttagctaggt 1980
gaaccccgagg gttaggggaaa gatgttaaca cctttgagct ctctggagtt gacatggaa 2040
agcaggtagtg tgttatgtag agctagtctt caaagctgcc ctgcctgttt tagggagcgt 2100
tcaccaaaaa gattgaggct ctttttagaa ttgaatttac tcttcagtat ttcttaagt 2160
tcagctttct aaaaggcata tatttttcaa agaagtggag atgcagtttc tcacgttgca 2220
acctattctc aagtgtgtta aatggtatct cttagtaact tgcactcgtt aaagaacac 2280
ggagctgggc catcgtcaga actaagtcag ggaaggagat ggatgagaag gccagaatca 2340
ttcctagtac atttgctaac actttattga gaaattgacc atgaattaat ggactcatct 2400
taatttcttc taagtcacata tatagataga tatctatctg tacagatttc tatttatcca 2460
tagataggta tctatacata cacatctcaa gtgcatttat tcccactctc attaatccat 2520
catgttctcta aatttttgta atcttaactg aaaaaaaagt gcactgaact tcaaaaaaaa 2580
acaaaaaaca acaacaacaa aaaaacagtc caaactgata tatctata tctgttaaaa 2640
ttcaaaagtg aacgaaagca tttaactggc cagttttgat tgcaaatgct gtaagatat 2700
agaatgaagt cctgtgaggc cttcctatct ccaagtcgat gtattttctg gagaccaaac 2760
cagataccag ataatacaca agaaagcttt tttaataagg cttaaaccaa gcccttgtct 2820
agatatcttt agttttgtgc caaggtagca ctgtgagaaa tctcacttgg atgttatgta 2880
aggggtgaga cacacaagtc tgactatgag tgaggaaaa atctgggctt tctcgtcagt 2940
ttggtgcatt tgcgtcgtct gttgctactg ttgctcctca acgctgtgct taaacaagct 3000
taaaactctta gccatacaagg tggctcttat gtacatagtt gttatacat ccaatlaagt 3060
atgtctgaca tgcatttttt gtaggagaaa aatgatgtct aatgatattt tgagttaaaa 3120
caotcttttg ggagatttg ctgaaaagtt gcacttttgt tacaatgctt atgcttgcta 3180
taagcttatg ctgctctaaa ttattttaaa aaattatatt actgtctgtg agaaccacgc 3240
tggtttagaa aagtttagta tgtgacgata aactagaaat tacccttata tcttagtatt 3300
tctcagcact cataaaattct attaccataa tattgcoaca ctatttttgg atttaaaaat 3360
tcttaactaa gaataaaaaac tttaatacac gatattgata tgcctaataa taaaaaaga 3420
cataatggat gctcaattag ttttaagata tctataacta tagggatata aatcactaca 3480
gttctcagat ttacaccttt tttttgcat tggcttgat tcacacattt ccaactctct 3540
gcaagcctcc aggcctctggc tttgtctacc tgctcgttcc caatgtatct taatgaaaag 3600
tgcnaaagaa aaacctacca attaaaaaaa aaaaaaaaaa a 3641

```

<210> 67
 <211> 482
 <212> PRT
 <213> Homo sapiens

<400> 67
 Met Asp Glu Gly Ile Pro His Leu Gln Glu Arg Gln Leu Leu Glu His
 1 5 10 15
 Arg Asp Phe Ile Gly Leu Asp Tyr Ser Ser Leu Tyr Met Cys Lys Pro
 20 25 30
 Lys Arg Ser Met Lys Arg Asp Asp Thr Lys Asp Thr Tyr Lys Leu Pro
 35 40 45
 His Arg Leu Ile Glu Lys Lys Arg Arg Asp Arg Ile Asn Glu Cys Ile
 50 55 60
 Ala Gln Leu Lys Asp Leu Leu Pro Glu His Leu Lys Leu Thr Thr Leu
 65 70 75 80
 Gly His Leu Glu Lys Ala Val Val Leu Glu Leu Thr Leu Lys His Leu
 85 90 95
 Lys Ala Leu Thr Ala Leu Thr Glu Gln Gln His Gln Lys Ile Ile Ala
 100 105 110
 Leu Gln Asn Gly Glu Arg Ser Leu Lys Ser Pro Ile Gln Ser Asp Leu
 115 120 125

Asp Ala Phe His Ser Gly Phe Gln Thr Cys Ala Lys Glu Val Leu Gln
 130 135 140
 Tyr Leu Ser Arg Phe Glu Ser Trp Thr Pro Arg Glu Pro Arg Cys Val
 145 150 155 160
 Gln Leu Ile Asn His Leu His Ala Val Ala Thr Gln Phe Leu Pro Thr
 165 170 175
 Pro Gln Leu Leu Thr Gln Gln Val Pro Leu Ser Lys Gly Thr Gly Ala
 180 185 190
 Pro Ser Ala Ala Gly Ser Ala Ala Ala Pro Cys Leu Glu Arg Ala Gly
 195 200 205
 Gln Lys Leu Glu Pro Leu Ala Tyr Cys Val Pro Val Ile Gln Arg Thr
 210 215 220
 Gln Pro Ser Ala Glu Leu Ala Ala Glu Asn Asp Thr Asp Thr Asp Ser
 225 230 235 240
 Gly Tyr Gly Gly Glu Ala Glu Ala Arg Pro Asp Arg Glu Lys Gly Lys
 245 250 255
 Gly Ala Gly Ala Ser Arg Val Thr Ile Lys Gln Glu Pro Pro Gly Glu
 260 265 270
 Asp Ser Pro Ala Pro Lys Arg Met Lys Leu Asp Ser Arg Gly Gly Gly
 275 280 285
 Ser Gly Gly Gly Pro Gly Gly Gly Ala Ala Ala Ala Ala Ala Leu
 290 295 300
 Leu Gly Pro Asp Pro Ala Ala Ala Ala Leu Leu Arg Pro Asp Ala
 305 310 315 320
 Ala Leu Leu Ser Ser Leu Val Ala Phe Gly Gly Gly Gly Ala Pro
 325 330 335
 Phe Pro Gln Pro Ala Ala Ala Ala Ala Pro Phe Cys Leu Pro Phe Cys
 340 345 350
 Phe Leu Ser Pro Ser Ala Ala Ala Ala Tyr Val Gln Pro Phe Leu Asp
 355 360 365
 Lys Ser Gly Leu Glu Lys Tyr Leu Tyr Pro Ala Ala Ala Ala Pro
 370 375 380
 Phe Pro Leu Leu Tyr Pro Gly Ile Pro Ala Pro Ala Ala Ala Ala
 385 390 395 400
 Ala Ala Ala Ala Ala Ala Ala Ala Ala Phe Pro Cys Leu Ser
 405 410 415
 Ser Val Leu Ser Pro Pro Pro Glu Lys Ala Gly Ala Ala Ala Thr
 420 425 430
 Leu Leu Pro His Glu Val Ala Pro Leu Gly Ala Pro His Pro Gln His
 435 440 445
 Pro His Gly Arg Thr His Leu Pro Phe Ala Gly Pro Arg Glu Pro Gly
 450 455 460
 Asn Pro Glu Ser Ser Ala Gln Glu Asp Pro Ser Gln Pro Gly Lys Glu
 465 470 475 480
 Ala Pro

<210> 68

<211> 3624

<212> DNA

<213> Homo sapiens

<400> 68

tagtctaact cgcggtgtgc acgcccactg cagcggagcc gcccgccggg gcgctgcggg 60
 acgggcccggc ggctgccggc aggaagccgac gagccgggtg actgcccgcc cgggcacagt 120
 ccggggccac agcgcgcgagc ccgggcggga gtggcccccgc gcaggcaggg acggcgccg 180
 cgactccaa cccggcgggc acctcggggg cggcgcgggg gcgcagcctt ctgctcccg 240

cctctgtgac	aagcgcccg	gagcgggag	ccgattgcc	gggctcggg	tgggcggga	300
cgacggcaact	gggctgtgc	ggggcccg	gcgtcgcat	gaacatcgt	gtggagtct	360
ctcgtgtcac	tttcaaaag	ctctggcg	tcgtctggc	cgcgccgc	tggtcgtgc	420
ggcccaagg	gaagagcgt	gcggccag	tgtgctcat	caccgcgc	ggagccgc	480
tgggccgct	cttcgcgt	gagttcgcc	ggcgtcggc	gctgctggt	ctgtgggaca	540
tcaacacga	aagcaacag	gagacgct	gcactgctg	ccacatctac	cgcgacctg	600
agggcgccga	cgccgctgc	ctgcaagct	ggaatggtga	ggaagaaat	ctgcgccact	660
taacttgc	ggtttttacc	tacacctgt	acgtgggaa	gagggagaac	gtctacctga	720
cggtgaaag	agtcocgaag	gaggttggc	aagtctcagt	ctgggtcaat	aatgctggtg	780
tggtctctg	gcataccct	ctggaatgt	ctgatgagct	cattgagaga	accatgatgg	840
tcaattgcc	tgacacatc	tggaacct	aggctttct	tcctacgatg	ctggagatta	900
atcatgttca	tatttgtaca	gttgcaagt	ccttgggatt	gttcagta	gccggagttg	960
aggattact	tgccagtaaa	tttggagttg	tggttttca	tgaatccctg	agccatgaac	1020
taaaagctgc	tgaaaaggat	ggaattaaaa	caaccttgg	ttgcccttat	ctttagagaca	1080
ctggcatgtt	cagagctgc	cgaatcagga	aagaaattga	gccttttctg	ccacctctga	1140
agcctgatta	ctgtgtgaag	caggccatga	aggccatcct	cactgaccag	cccatgatct	1200
cgacacccg	cctcatgtac	atcgtgacct	tcactgaag	catctacca	cttgaagcag	1260
ttgtgtgcat	tcatcggttc	ctaggagcgg	acaagtgtat	gtaccccttt	attgctcaaa	1320
gaaagcaagc	cacaacaact	aatgaagcaa	aaaatggaat	ctaaagaact	ttttgtatgg	1380
aatattactt	ctatcagaag	atgatcaaga	tgittcagtc	cagtgacat	cagcatctgt	1440
gacattttat	ggattctaaa	cttgtgttg	ttctttttta	aatcaactt	ttaaaaaat	1500
aaagtgtaaa	ttaacccag	agagtactg	gaaaattgtga	tcagtacaag	tgnaacttagg	1560
ttgttgccaa	cagggtcctt	ttaggcagaa	cccagaiaacc	agtcacatct	gtagagaagc	1620
agtgtgacat	cttcaggtta	coattatttt	ttaatgagca	ggaagtctag	aaatgataac	1680
tagactgtat	gtttcatgtg	tgtattttt	cagaattccc	agagtttact	catctctgtt	1740
attaaactct	agccactgca	catcttcgca	atttcaagg	ctgatagtct	tgattattct	1800
caagctttct	aagtttccgt	tttgcaaggc	ctaggtgact	ttttcatgtg	gtttgtatgt	1860
ttagctttt	tgaaaaggaa	ttttgaaatc	tcatacaact	gaagttaagt	atgtctgagt	1920
gttacagtwa	aggtgaccaa	gtctctttct	taaaagcaca	atgactaaag	tattagtgtga	1980
ttttttttt	ttttttttga	tgaggtctcg	ctctgtcac	aggctggagt	gcagtagcac	2040
aatcacggct	cactgcaatc	tcgtctccc	rgtttcaagt	gattctgctg	cttcagcctc	2100
ccaagtact	gggactacag	gcactgcga	ccacgcccag	ctaatttttg	tatttttagt	2160
agagacggga	tttccacatg	ttggtcagga	tggtctccat	ctcttgacat	tgtgaacgaa	2220
ctgcctcgcc	ctcccaaggt	gctgggatta	caggcatag	ccactgcacc	cagccttgaa	2280
tttttaattt	tatctctgat	atacttcatt	aagtgctctg	agacctaatt	atccataaag	2340
atcatcattt	ttctactcat	gaattttgct	gcatacagaa	agtgccttt	cctcagggaag	2400
ttgtctgttt	tcattttttt	ggatggactc	ttatctagaa	tacatagcag	cttgcaaaag	2460
aaacagtttt	taaaaaatggg	aactcttaca	ttgaaaagtc	cccatttttg	tgcccaactat	2520
gattagtga	aggaagaaat	ctattctcat	ggcatatgta	tggaagggtg	taaaagtatt	2580
tttgaaggt	ttattccat	tgtagaacag	caaatgacat	ttttacagta	tttttttgta	2640
aagcaacta	ttttgtgct	tgaatttggt	atatgtgtat	tagtgaacaa	ttgtaaaggt	2700
gaactctac	ctctgtatct	aaatgtatac	catccacttg	taaatgacta	taaaactatta	2760
tgtgattgt	ttttttttta	gaatgtcttg	tttaaatagt	ggccaatgtt	taaggotgtt	2820
aaaaaagcc	acacttttact	aattggggag	ttttataaat	gactgattaa	atttaaagaa	2880
ttactctaca	tgcacttggt	tgtattttag	ttatcagcag	tgttgtaagg	aaattatttg	2940
tgttttttt	tatgatctg	atccccactt	aggtaaagaa	aaatatttga	atggaatagt	3000
cttgggaaac	agacattaac	aacctaggg	gcctgcactc	aaatagccga	tggtactgtc	3060
ctagatgat	agacttgatt	aagggttgt	ttgtaccaaa	agtggggaaa	caatgccatg	3120
acctgtgttt	tagttttgct	gcaccacaga	tcaaatctgc	actgtgtcta	catataggaa	3180
aggtctggt	gtgtgctaat	gttcccaatg	caggacttga	ggaagagctc	gtttatattg	3240
ttccatttct	cttlatcaaa	gataaccaaa	ccttatggcc	cttatacaaa	tggaaggcact	3300
ggctgcctct	taattttcaa	tcattggacct	aaagaagtac	cttgaagggt	ctcaacagt	3360
ccagtgggg	acagatatac	tcagagatta	ctcaggtgtg	gcctccagcg	agcctggagt	3420
acaccagacc	ctcttagaga	aatctgttat	aatttaacaa	cccactatct	caacttaaaa	3480
ctgaggaag	tgctctttac	atctaatttt	attctgtgtg	ttataaact	aaacctattt	3540
ctatttttgt	tgtgtattgc	ccttataaag	gtgtccctct	ccaagttcaa	taaaactaatt	3600
catttaaaaa	aaaaaaaaaa	aaaa				3624

<210> 69

<211> 341

<212> PRT

<213> Homo sapiens

<400> 69

```

Met Asn Ile Val Val Glu Phe Phe Val Val Thr Phe Lys Val Leu Trp
 1          5          10          15
Ala Phe Val Leu Ala Ala Ala Arg Trp Leu Val Arg Pro Lys Glu Lys
 20          25          30
Ser Val Ala Gly Gln Val Cys Leu Ile Thr Gly Ala Gly Ser Gly Leu
 35          40          45
Gly Arg Leu Phe Ala Leu Glu Phe Ala Arg Arg Arg Ala Leu Leu Val
 50          55          60
Leu Trp Asp Ile Asn Thr Gln Ser Asn Glu Glu Thr Ala Gly Met Val
 65          70          75          80
Arg His Ile Tyr Arg Asp Leu Glu Ala Ala Asp Ala Ala Leu Gln
 85          90          95
Ala Gly Asn Gly Glu Glu Glu Ile Leu Pro His Cys Asn Leu Gln Val
100          105          110
Phe Thr Tyr Thr Cys Asp Val Gly Lys Arg Glu Asn Val Tyr Leu Thr
115          120          125
Ala Glu Arg Val Arg Lys Glu Val Gly Glu Val Ser Val Leu Val Asn
130          135          140
Asn Ala Gly Val Val Ser Gly His His Leu Leu Glu Cys Pro Asp Glu
145          150          155          160
Leu Ile Glu Arg Thr Met Met Val Asn Cys His Ala His Phe Trp Thr
165          170          175
Thr Lys Ala Phe Leu Pro Thr Met Leu Glu Ile Asn His Gly His Ile
180          185          190
Val Thr Val Ala Ser Ser Leu Gly Leu Phe Ser Thr Ala Gly Val Glu
195          200          205
Asp Tyr Cys Ala Ser Lys Phe Gly Val Val Gly Phe His Glu Ser Leu
210          215          220
Ser His Glu Leu Lys Ala Ala Glu Lys Asp Gly Ile Lys Thr Thr Leu
225          230          235          240
Val Cys Pro Tyr Leu Val Asp Thr Gly Met Phe Arg Gly Cys Arg Ile
245          250          255
Arg Lys Glu Ile Glu Pro Phe Leu Pro Pro Leu Lys Pro Asp Tyr Cys
260          265          270
Val Lys Gln Ala Met Lys Ala Ile Leu Thr Asp Gln Pro Met Ile Cys
275          280          285
Thr Pro Arg Leu Met Tyr Ile Val Thr Phe Met Lys Ser Ile Leu Pro
290          295          300
Phe Glu Ala Val Val Cys Met Tyr Arg Phe Leu Gly Ala Asp Lys Cys
305          310          315          320
Met Tyr Pro Phe Ile Ala Gln Arg Lys Gln Ala Thr Asn Asn Asn Glu
325          330          335
Ala Lys Asn Gly Ile
340

```

<210> 70

<211> 1428

<212> DNA

<213> Homo sapiens

<400> 70

```

ggcagcaggc ggagacagag acttcacgac tcccagtcct ctcctcgccg cggccgcgcg 60
ctctctcttc tctctctctc ctctctcttc tctctctctg ctccacacgc catgtctgct 120
tagaccagag cagccccaca gccaaactagg gcagctgcgc ccgccacaac agcaaggaca 180
gcgcgtgcgc cgcgcgtga gogatgacag gagtgtttga cagaagggtc cccagcatcc 240
gatccgcgca cttccaaagc cgttccaga cgtccgcagc tatgcacocat cgtctcagg 300
aatccccaac tttgcccgag tcttcageta cogatctetga ctactacagc cctacggggg 360
gagcccgcca cggctactgc tctctactct cggcttctta tggcaaaagt ctcaacccct 420
accagtatca gtatcacgag gtgaacggct cgcgcgggag ctaccacagc aagcgtctag 480
ctacgctatg ctacgctagc tctaccaccc agtacggcgc cgcctacaac caggtcccaa 540
gcgcacacaa ccagccagag aaagaagtga cagagcccca ggtgagaagt gtgaatggca 600
aaccaagaa agttcgtaaa cccaggacta tttattccag ctttcagctg cgcgcattac 660
agagaagggt tcagaagact cagtacctcg cttgcgcgga acgcgcgcgag ctggccgcct 720
cgtcgtgatt gacacaaaaca caggtgaaaa tctgttttca gaacaaaaga tccaagatca 780
agaagatcat gaaaaacggg gagatgcccc cggagcacag tcccagctcc agcgacccaa 840
tgcgctgtaa ctgcgcgcag tctccagcgg tgtgggagcc ccagggtctg tcccgctcgc 900
tcagccacca cctcatgcc caccctccga cctccaacca gtcccaagcg tccagctacc 960
tgggaactc tgcactctgg tacacaaagt cagccagctc aatcaattcc cacctgccg 1020
cgcgcggctc cttacagcac ccgcgtggcgc tggcctccgg gacactctat tagatggct 1080
gtctctcttc actctctttt ttgggactac tgtgttttgc tgtctataga aatcataaag 1140
aaaggaaatt atatggggaa gttcggaaaa ctgaaaaaga ttcattgtga aagctttttt 1200
ttgcattgtt gttattgcatt ttcaaaagac cccccctttt ttacagagag actttttttg 1260
cgcaactgtg gacaactttca atggtgcctt gaaatctatg acctcaactt tccaagaagc 1320
tttttcaat gttattttag caatgtaaat aagtgtatag agaggaatta aactgtatat 1380
tctggataaa taaaattatt tcgaccatga aaaaaaaaaa aaaaaaaaaa 1428

```

<210> 71

<211> 289

<212> PRT

<213> Homo sapiens

<400> 71

```

Met Thr Gly Val Phe Asp Arg Arg Val Pro Ser Ile Arg Ser Gly Asp
1 5 10 15
Phe Gln Ala Pro Phe Gln Thr Ser Ala Ala Met His His Pro Ser Gln
20 25 30
Glu Ser Pro Thr Leu Pro Glu Ser Ser Ala Thr Asp Ser Asp Tyr Tyr
35 40 45
Ser Pro Thr Gly Gly Ala Pro His Gly Tyr Cys Ser Pro Thr Ser Ala
50 55 60
Ser Tyr Gly Lys Ala Leu Asn Pro Tyr Gln Tyr Gln Tyr His Gly Val
65 70 75 80
Asn Gly Ser Ala Gly Ser Tyr Pro Ala Lys Ala Tyr Ala Asp Tyr Ser
85 90 95
Tyr Ala Ser Ser Tyr His Gln Tyr Gly Gly Ala Tyr Asn Arg Val Pro
100 105 110
Ser Ala Thr Asn Gln Pro Glu Lys Glu Val Thr Glu Pro Glu Val Arg
115 120 125
Met Val Asn Gly Lys Pro Lys Lys Val Arg Lys Pro Arg Thr Ile Tyr
130 135 140
Ser Ser Phe Gln Leu Ala Ala Leu Gln Arg Arg Phe Gln Lys Thr Gln
145 150 155 160
Tyr Leu Ala Leu Pro Glu Arg Ala Glu Leu Ala Ala Ser Leu Gly Leu
165 170 175
Thr Gln Thr Gln Val Lys Ile Trp Phe Gln Asn Lys Arg Ser Lys Ile
180 185 190
Lys Lys Ile Met Lys Asn Gly Glu Met Pro Pro Glu His Ser Pro Ser
195 200 205
Ser Ser Asp Pro Met Ala Cys Asn Ser Pro Gln Ser Pro Ala Val Trp

```

210	215	220
Glu Pro Gln Gly Ser Ser Arg Ser Leu Ser His	His Pro His Ala His	
225	230	235
Pro Pro Thr Ser Asn Gln Ser Pro Ala Ser Ser Tyr Leu Glu Asn Ser		240
	245	250
Ala Ser Trp Tyr Thr Ser Ala Ala Ser Ser Ile Asn Ser His Leu Pro		255
	260	265
Pro Pro Gly Ser Leu Gln His Pro Leu Ala Leu Ala Ser Gly Thr Leu		270
	275	280
		285

Tyr

<210> 72

<211> 2036

<212> DNA

<213> Homo sapiens

<400> 72

```

gacagcttga accattcccc tggccagagt ggattctcca gctatggctc cagcttcagc 60
acctcaccca ctggcacagag ccatacaccc taccagatgc accgacacac agggttctat 120
caaggaggaa atggactggg caacgcagcc gggttcggga gtgtgcacca ggactatcct 180
tctacccccc gcttccccca gagccagtag ccccgattat accgctcacc ctacacacct 240
ccctaagctcc cggccagcag catctgccct tgcgccctct caccgtccac ctacgtctcc 300
caggagcat ctacaaacgt ccccaaccag agttccagat cactgtgtgt tgaatacaac 360
acacacaatg gacctccac accagcgaaa gaggagaca cagacaggcc gaccgggcc 420
tcgacggga agctccagag ccgtctctaa aggagcagtg acccgcccc ggacggggac 480
aatgagatt agcgtgtgtt cgtgtgggac ttgatgaga caataattat ttttaactcc 540
ttactcacgg ggacatttgc atccagatac gggaaggaca ccaacagctc cgtgcgcatt 600
ggccttatga tggaaagat gatcttcaac ctgcagata cacatctgtt ctccaatgac 660
ctggaggatt gtgaccagat caacgttgat gacgtctcat cagatgacaa tggccaagat 720
ttaagcacat acaactcttc cgtgcagcgg ttccacagtt cgccccaggg agccaacctg 780
tgctctgggt ctggcggtgca cggcgcggtg gactggatga ggaagctggc ctccgctac 840
cgcggggtga aggagatgta caatacctac aagaacaacg ttggtgggtt gataggcact 900
cccaaaaggg agacctggct acagctcoga gctgagctgg aagctctcac agacctctgg 960
ctgacccact cctctgaagg actaaacctc atcaactccc ggcccaactg tgtcaatgtg 1020
ctgtgccaca ccactcaact aattcctgcc ctggccaaag tctgtctata tggctggggg 1080
tctgtgtttc ctattgagaa catctacagt gcaaccaaga cagggaagga gactgtcttc 1140
gagaggataa tgcagagatt cggcagaaaa gctgtctcac tgggtgatcg tggatgtgtg 1200
gaagaggagc aaggacgaa aaagcacaaac atgcctttct ggcggatact ctgcacgaca 1260
gacctggagg cactgaggca cgccttgagg ctggagatt tatagcaga tcagcagcat 1320
ctccactcgc catctcaccc tcagaccccc tgcctctccc cactcccca agcagaactc 1380
cagagaccca gatgttgagc accaggaagg ggcacacag ccgagacgac gtgtccagt 1440
acctactcag aagccgtcca tcagtccaaa tgggggttct gagaaggaaa gtaccaaca 1500
ttggctcagg agtatttgac tttggggaaa agggctggct cgagctctag actctctctg 1560
aagactcaca gaacaaaagc aaggaattgc caattgggg ggtgcctggg gatgaggagg 1620
ggatgggttt gtctgtgttt ctttttaatt tatggaactg tctcaatctc ccagaattat 1680
gctctgttac cgtgtggctt ggttttctta agcactctct tctccaagtt gtgctttgtg 1740
gtatgtaggg tggttcacag ttctaatgta agcactctct tctccaagtt gtgctttgtg 1800
gggacaatca tcttttgaac attagagagg aaggcagctc aagctgttga aaagactatt 1860
gctatttttt gcttttaagc acctacttga acctgtggg acagtgcacg tgccttaagc 1920
tacatcttgt tttctaggaa gagggggtag ctgggaagga atgggtgctt tgtgatggat 1980
aaaaggcatt aataaaaacc acgttttcat ttggaaaaaa aaaaaaaa 2036

```

<210> 73

<211> 434

<212> PRT

<213> Homo sapiens

<400> 73

Asp Ser Leu Asn His Ser Pro Gly Gln Ser Gly Phe Leu Ser Tyr Gly
 1 5 10 15
 Ser Ser Phe Ser Thr Ser Pro Thr Gly Gln Ser Pro Tyr Thr Tyr Gln
 20 25 30
 Met His Gly Thr Thr Gly Phe Tyr Gln Gly Gly Asn Gly Leu Gly Asn
 35 40 45
 Ala Ala Gly Phe Gly Ser Val His Gln Asp Tyr Pro Ser Tyr Pro Gly
 50 55 60
 Phe Pro Gln Ser Gln Tyr Pro Gln Tyr Tyr Gly Ser Ser Tyr Asn Pro
 65 70 75 80
 Pro Tyr Val Pro Ala Ser Ser Ile Cys Pro Ser Pro Leu Ser Thr Ser
 85 90 95
 Thr Tyr Val Leu Gln Glu Ala Ser His Asn Val Pro Asn Gln Ser Ser
 100 105 110
 Glu Ser Leu Ala Gly Glu Tyr Asn Thr His Asn Gly Pro Ser Thr Pro
 115 120 125
 Ala Lys Glu Gly Asp Thr Asp Arg Pro His Arg Ala Ser Asp Gly Lys
 130 135 140
 Leu Arg Gly Arg Ser Lys Arg Ser Ser Asp Pro Ser Pro Ala Gly Asp
 145 150 155 160
 Asn Glu Ile Glu Arg Val Phe Val Trp Asp Leu Asp Glu Thr Ile Ile
 165 170 175
 Ile Phe His Ser Leu Leu Thr Gly Thr Phe Ala Ser Arg Tyr Gly Lys
 180 185 190
 Asp Thr Thr Thr Ser Val Arg Ile Gly Leu Met Met Glu Glu Met Ile
 195 200 205
 Phe Asn Leu Ala Asp Thr His Leu Phe Phe Asn Asp Leu Glu Asp Cys
 210 215 220
 Asp Gln Ile His Val Asp Asp Val Ser Ser Asp Asp Asn Gly Gln Asp
 225 230 235 240
 Leu Ser Thr Tyr Asn Phe Ser Ala Asp Gly Phe His Ser Ser Ala Pro
 245 250 255
 Gly Ala Asn Leu Cys Leu Gly Ser Gly Val His Gly Gly Val Asp Trp
 260 265 270
 Met Arg Lys Leu Ala Phe Arg Tyr Arg Arg Val Lys Glu Met Tyr Asn
 275 280 285
 Thr Tyr Lys Asn Asn Val Gly Gly Leu Ile Gly Thr Pro Lys Arg Glu
 290 295 300
 Thr Trp Leu Gln Leu Arg Ala Glu Leu Glu Ala Leu Thr Asp Leu Trp
 305 310 315 320
 Leu Thr His Ser Leu Lys Ala Leu Asn Leu Ile Asn Ser Arg Fro Asn
 325 330 335
 Cys Val Asn Val Leu Val Thr Thr Thr Gln Leu Ile Pro Ala Leu Ala
 340 345 350
 Lys Val Leu Leu Tyr Gly Leu Gly Ser Val Phe Pro Ile Glu Asn Ile
 355 360 365
 Tyr Ser Ala Thr Lys Thr Gly Lys Glu Ser Cys Phe Glu Arg Ile Met
 370 375 380
 Gln Arg Phe Gly Arg Lys Ala Val Tyr Val Val Ile Gly Asp Gly Val
 385 390 395 400
 Glu Glu Glu Gln Gly Ala Lys Lys His Asn Met Pro Phe Trp Arg Ile
 405 410 415
 Ser Cys His Ala Asp Leu Glu Ala Leu Arg His Ala Leu Glu Leu Glu
 420 425 430
 Tyr Leu

<210> 74
 <211> 1907
 <212> DNA
 <213> Homo sapiens

<400> 74
 cggccagata cctcagcgct acctggcgga actggatttc tctccgcct gccggcctgc 60
 ctgccacagc cggactccgc cactccggta gcctcatggc tgcaacctgt gagattagca 120
 acatttttag caactacttc agtgcgatgt acagctcgga ggaactccacc ctggcctctgt 180
 ttccccctgc tgccacottt ggggcccgatg acttggtact gacctgagc aacccccaga 240
 tgtcatttga gggtacagag aagccagct ggttggggga acagccccag ttctggtcga 300
 agacgcaggt tctggactgg atcagctacc aagtggagaa gaacaagtac gacgcaagcg 360
 ccattgactt ctcacgatgt gacatggatg gcgccacct ctgcaattgt gcccttgagg 420
 agctgcgctt ggtcttggg cctctggggg accaactcca tgcaccagct cgagacctca 480
 cttccagctc tctctgatg ctcagtttga tcattgaact gctggagaag gatggcatgg 540
 ccttcacaga gccctagac ccaggccctt ttgaccaggg cagcccttt gccccaggagc 600
 tgcgtgacga cggctacgaa gccagccctt accaccccg cagctgtggc gcaggagccc 660
 cctccctggc cagctctgac gtctccaccg cagggactgg tcttctcgg agctcccact 720
 cctcagactc cgggtgaagt gacgtggacc tggatccacc tgatggcaag cctttcccca 780
 gcgatggttt tctgactct aagaaggggg atcccaagca cgggaagcgg aaacgagccc 840
 ggccccgaaa cctgagcaaa gactactggg actgtctcga gggcaagaa agcaagcacg 900
 cgccccaggg gccccacctg tgggagtcca tccggacat cctcatccac ccaggagctc 960
 acgagggcct catgaagtgg gagaatcgcg atgaaggcgt ctcaagttc ctgcctccg 1020
 aggtgtgggc ccaactatgg gcccaaaaaga aaaaagaacag caacatgacc taccgtaagg 1080
 tgagccggcg catgaggtac tactacaac gggagatcct ggaacgggtg gatggccggc 1140
 gactcgtcta caagtttggc aaaaactcaa gcggctggaa ggaggaagag gtctccaaga 1200
 gtcggaactg agggttggaa ctatacccg gaccaaact acgggaccat cgaggcctgc 1260
 aaacctctct gggaggacag gcagggcaga tggcccctcc actggggaat gctccagct 1320
 gtgctgtgga gagaagctga tgttttggg tatgtcagc catgtcctt ggaactggag 1380
 actatggcct cgctcccca cctcctctt ggaattacaa gccctggggt ttgaagctga 1440
 ctttatagct gcaagtgtat ctcttttat ctggtgcctc ctcaaccaca gtctcagaca 1500
 cttaaatgca gacaacacct tcttctgca gacacttga ctgagccaag gaggcttggg 1560
 aggccttagg gagcaccgtg atggagagga cagagcagg gctccagcac ttctttctgg 1620
 actcggcttc actccctgc tcagtcttg ggtccacgg gcaggggtca gagcactccc 1680
 taatttatgt cctatataaa tatgtcagat gtacataag atctattttt tctaaaacat 1740
 tccccctccc actccttccc cacagagtgc tggactgttc caggccctcc agtgggctga 1800
 tgcctggacc cttagtaggg ggtccccagc tcttttctc tgtgaatgga ggcagagacc 1860
 tocaataaag tgccttctgg gctttttcta aaaaaaaaa aaaaaaa 1907

<210> 75
 <211> 371
 <212> PRT
 <213> Homo sapiens

<400> 75
 Met Ala Ala Thr Cys Glu Ile Ser Asn Ile Phe Ser Asn Tyr Phe Ser
 1 5 10 15
 Ala Met Tyr Ser Ser Glu Asp Ser Thr Leu Ala Ser Val Pro Pro Ala
 20 25 30
 Ala Thr Phe Gly Ala Asp Asp Leu Val Leu Thr Leu Ser Asn Pro Gln
 35 40 45
 Met Ser Leu Glu Gly Thr Glu Lys Ala Ser Trp Leu Gly Glu Gln Pro
 50 55 60
 Gln Phe Trp Ser Lys Thr Gln Val Leu Asp Trp Ile Ser Tyr Gln Val
 65 70 75 80
 Glu Lys Asn Lys Tyr Asp Ala Ser Ala Ile Asp Phe Ser Arg Cys Asp

[illegible]

```
<210> 76
<211> 3951
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<222> (1)...(3951)
<223> n = A,T,C or G
```

#400> 76							
nnagaccacg	cgtcgcgggtg	cgtgtctggac	acgtctccgg	ggtgggtcgt	ccggcctctcg	60	
atcttagacg	aatttttacaa	tgtgaagttc	tgcatagatg	ccagtcaccc	agatgttgga	120	
cagctggctg	atgacatattg	atctctggcg	tgttatgac	agacacacct	tgttgcattcg	180	
agatagatac	atcagatatt	ctatagatga	tgtcgaaca	ttcgcgcgga	agaggagctt	240	
ctcgtgttca	tgaagagcga	agactatccc	catgaactc	tggcgcgcga	tatccacgaa	300	
gaagcgcgat	atcgtctgca	agactgtgac	cagctctttg	aatactagaa	tgaactagcga	360	
gtatcacaaa	agttttcatg	cagactcctc	cactcagcat	tttcaatggt	tgaagagcag	420	
tttcaacaaa	aactcgaaga	cgcaaatgat	ctccagaqa	tatccacgat	ccaggatggt	480	

aaggaatgtg	accaaagttt	tcttgatttg	caaagcctgg	agaaacacat	gctgtcacat	540
actgaagaga	gggaatatcaa	gtgtgatcag	tgtcccaagg	catttaactg	gaagtgcaat	600
ttaattcgcc	acacagatgtc	acatgacagt	ggaaagcact	atgaatgtga	aaactgtgcc	660
aaggttttca	cggacctag	caaccttcag	cggcaccattc	gctctcagca	tgctcggtgcc	720
cgggcccatc	gtctgccgga	gtgtggccaaa	acgttttgcca	cttcgtcggg	cctcacaacaa	780
cacaagcaca	tcacagcag	tgtgaagccg	tttatctcat	ttctctaact	aatgtaccoca	840
tttctgtata	gagacttgag	atcggtacct	ttgaaaatgg	aaaccccaatc	accaggtgaa	900
gtaaagaaac	tgcgaagaagg	cagctctcag	tccccctttg	atctcacacc	taagcgaaag	960
gatgagaagg	ccttgactcc	agtccccctc	aagcctccag	tgacacctcg	cacaagctcaa	1020
gaccagcccc	tggatctaa	tatgggcagt	aggagttagag	ccagtgggac	aaagctgact	1080
gagcctcgaa	aaaaccacgt	gtttggggga	aaaaaaggaa	gcaacgtcga	atcaagacct	1140
gcttcagatg	gttccttgca	gcattgcaaga	cccaactcctt	tctttatgga	ccctattttac	1200
agagttagaa	aaagaaaaact	aactgcaccca	cttgaagctt	taaaagagaa	atactgtgagg	1260
ccttctccag	gatttctgtt	tcaoccccaa	atgtcagcta	ttgaaaacat	ggcagaaaag	1320
ctagagagat	tcagtgccct	gaaacctgag	gccagtgagc	tcttacagtc	agtgccctct	1380
atgttcaact	tcaggggccc	tcccaatgoc	ctgcacagaga	acctctctgcg	gaagggaag	1440
gagcgtcata	cctgcagata	ctgtggcaag	atittttcca	ggctctgcaa	cctacacagc	1500
cacttgagaa	cccaacacagg	agagcagcct	tacagatgca	aatactgtga	cngatcaatt	1560
agcatattct	ctaaacttgca	aaggcatggt	cgcaacatcc	acaataaaga	gaagccattt	1620
aagtgtcaact	tatgtgttag	gtgttttgg	caacaaaccca	atttagacag	acacctaaag	1680
aaacatgaga	atgggaaact	gtccggtaca	gcaacatcgt	cgctctatc	tgaactggaa	1740
agtcacagtg	cgattcttgg	tgacaagaag	gatgcttact	tcacagaagt	tcgaattctt	1800
atttgggaaca	gcaacatgg	cagccaatct	ccaggaatg	tgaggagag	aatgaatggc	1860
agtcatttta	aaagtgaaaa	ggcttttggt	accagtcaat	attcagact	gctgtgaggt	1920
gaagaagttg	aaagttaggt	gttgttagat	gaggaggatg	aaagacaatga	tattacttga	1980
aaaaacaggaa	aggaacacgt	gacaagtaat	ttacatgaag	gaaacccctga	ggatgactat	2040
gaagaaaacca	gtgccctgga	gatgagttgc	aagacatccc	cagtgaggta	taagaggaa	2100
gaatataaaa	gtggaacttc	tgctctagat	catataaggg	acctcacaga	tagcctcaaa	2160
atgaggaaaa	tggaaagata	tcaattattc	gaagctgagc	tgctctctt	tagtactctc	2220
catgtgcgag	aggaacottaa	cgagccgttc	cacagaaggt	ccaaattcgca	ggcatatgct	2280
atgatgtctga	cactgtctga	caaggagctc	ctccatttcta	catccocagc	ttcttccaa	2340
gtgtgcccaca	gtatggccag	ggctgcggcg	gaatccagtg	ctatccagtc	cataagccac	2400
gtatgacgtt	atcaaggttg	accagagttg	gaccaagtcc	aacagtagca	tgctctcttc	2460
atataggact	atttacaaga	ctgctgagca	gaatgcotta	taaacctgca	gggtcactca	2520
tctaaagtct	agtgacctta	aactgaaatga	tttaaaaaag	aaaagaaaga	aaaagaaac	2580
tattttattct	cgatattttg	ttttgcacag	caaaggcagc	tgctgacttc	tgggaagatca	2640
atcaattcgca	cttaaaagtga	ttcagtgaaa	acaaaaaact	tggtggcgctg	aaagcatctt	2700
ccagtttacc	ccaccttagg	gtatgggtgg	gtgagaagg	cagttgagat	ggcagacttg	2760
ctatgaatga	acactccata	gaaactgaat	tctcttttgt	acaagatcac	ctgacatgat	2820
tgggaacag	tgcttttaat	tacagattta	attttttctg	tcgttaaa	ttattgtaat	2880
tttaacccctt	gaagacagaa	gtagtggat	gaaatgcaca	gtcaattatt	atagaaactg	2940
ataaacggga	gtaacttaag	ccocctttgc	cttctatttga	acattgtttg	aaactaggga	3000
aaaaaggat	gtgttatattg	taaaactatg	atgttaaac	ttcaaaagg	tttaagtcagt	3060
gargtaacot	agtaaccctg	agtaaccctg	taccactaat	aaattgtttg	ccaaactcctt	3120
gtaatacaat	cttaatttta	gacaatcatg	toactgtttt	taattgttat	tttttttgtg	3180
gtgtgtcggt	ttattttgtg	ttattttgtg	gcaaaactatt	gtttgtgtat	taaaatagca	3240
ctgtttccagt	cagcacttac	tttatgacgt	ctgaggcaca	ccocctttccg	aatttcaagg	3300
accaaagtgga	cccgactcgt	gtatgagagt	gocaaaatggt	gtttgtgctt	ttttaaactt	3360
cctttttgtt	tgtttgtttt	gttttctctc	ttaatgaact	aaatcagat	agatgcaact	3420
tagtttttgt	aatactgaaa	tnocattcaa	ttgtataaac	gattataatt	tctttctcag	3480
aagcatgatt	ctctctgatta	aaaaactgtac	tcocatattt	atgtcggtgt	tcgtcgaagt	3540
tgtgogagt	tatgtctaat	ttaatccat	ttgtaaaaatg	aagtgctccc	aaacttatng	3600
ttaaaaagaga	gaangtaaaa	aaacgactgt	attcagttat	tttgcccttt	attgaggaac	3660
cgagtttgtt	tctcttttgt	ttgtaatctc	attttngaaa	taatcagcaa	gttgaggtao	3720
tttctccaaa	tgctttgtac	aatataaaat	gttatgcctt	tcagtgcatt	actatgggag	3780
gagcaactaa	aaaaataaga	ctacaaaaaa	ggagtatttt	taagaacaaa	aaactgtgag	3840
ggggccccgg	taccocaattc	gccctatagt	gagtagtatt	acaattcaact	ggcgtcggt	3900
ttacaacgtc	gtgactggga	aaacccctgg	ttaccacact	taatgctctt	n	3951

<210> 77
 <211> 718
 <212> PRT
 <213> Homo sapiens

<400> 77

```

Met Lys Ser Glu Asp Tyr Pro His Glu Thr Met Ala Pro Asp Ile His
 1          5          10          15
Glu Glu Arg Gln Tyr Arg Cys Glu Asp Cys Gln Leu Phe Glu Ser
 20          25          30
Lys Ala Glu Leu Ala Asp His Gln Lys Phe Pro Cys Ser Thr Pro His
 35          40          45
Ser Ala Phe Ser Met Val Glu Glu Asp Phe Gln Gln Lys Leu Glu Ser
 50          55          60
Glu Asn Asp Leu Gln Glu Ile His Thr Ile Gln Glu Cys Lys Glu Cys
 65          70          75          80
Asp Gln Val Phe Pro Asp Leu Gln Ser Leu Glu Lys His Met Leu Ser
 85          90          95
His Thr Glu Glu Arg Glu Tyr Lys Cys Asp Gln Cys Pro Lys Ala Phe
100          105          110
Asn Trp Lys Ser Asn Leu Ile Arg His Gln Met Ser His Asp Ser Gly
115          120          125
Lys His Tyr Glu Cys Glu Asn Cys Ala Lys Val Phe Thr Asp Pro Ser
130          135          140
Asn Leu Gln Arg His Ile Arg Ser Gln His Val Gly Ala Arg Ala His
145          150          155          160
Ala Cys Pro Glu Cys Gly Lys Thr Phe Ala Thr Ser Ser Gly Leu Lys
165          170          175
Gln His Lys His Ile His Ser Ser Val Lys Pro Phe Ile Ser Phe Ser
180          185          190
Gln Ser Met Tyr Pro Phe Pro Asp Arg Asp Leu Arg Ser Leu Pro Leu
195          200          205
Lys Met Glu Pro Gln Ser Pro Gly Glu Val Lys Lys Leu Gln Lys Gly
210          215          220
Ser Ser Glu Ser Pro Phe Asp Leu Thr Thr Lys Arg Lys Asp Glu Lys
225          230          235          240
Pro Leu Thr Pro Val Pro Ser Lys Pro Pro Val Thr Pro Ala Thr Ser
245          250          255
Gln Asp Gln Pro Leu Asp Leu Ser Met Gly Ser Arg Ser Arg Ala Ser
260          265          270
Gly Thr Lys Leu Thr Glu Pro Arg Lys Asn His Val Phe Gly Gly Lys
275          280          285
Lys Gly Ser Asn Val Glu Ser Arg Pro Ala Ser Asp Gly Ser Leu Gln
290          295          300
His Ala Arg Pro Thr Pro Phe Phe Met Asp Pro Ile Tyr Arg Val Glu
305          310          315          320
Lys Arg Lys Leu Thr Asp Pro Leu Glu Ala Leu Lys Glu Lys Tyr Leu
325          330          335
Arg Pro Ser Pro Gly Phe Leu Phe His Pro Gln Met Ser Ala Ile Glu
340          345          350
Asn Met Ala Glu Lys Leu Glu Ser Phe Ser Ala Leu Lys Pro Glu Ala
355          360          365
Ser Glu Leu Leu Gln Ser Val Pro Ser Met Phe Asn Phe Arg Ala Pro
370          375          380
Pro Asn Ala Leu Pro Glu Asn Leu Leu Arg Lys Gly Lys Glu Arg Tyr
385          390          395          400
Thr Cys Arg Tyr Cys Gly Lys Ile Phe Pro Arg Ser Ala Asn Leu Thr

```

				405					410					415			
Arg	His	Leu	Arg	Thr	His	Thr	Gly	Glu	Gln	Pro	Tyr	Arg	Cys	Lys	Tyr		
			420						425				430				
Cys	Asp	Arg	Ser	Phe	Ser	Ile	Ser	Ser	Asn	Leu	Gln	Arg	His	Val	Arg		
		435					440					445					
Asn	Ile	His	Asn	Lys	Glu	Lys	Pro	Phe	Lys	Cys	His	Leu	Cys	Asp	Arg		
	450					455					460						
Cys	Phe	Gly	Gln	Gln	Thr	Asn	Leu	Asp	Arg	His	Leu	Lys	Lys	His	Glu		
465					470					475					480		
Asn	Gly	Asn	Met	Ser	Gly	Thr	Ala	Thr	Ser	Ser	Pro	His	Ser	Glu	Leu		
			485						490					495			
Glu	Ser	Thr	Gly	Ala	Ile	Leu	Asp	Asp	Lys	Glu	Asp	Ala	Tyr	Phe	Thr		
		500						505					510				
Glu	Ile	Arg	Asn	Phe	Ile	Gly	Asn	Ser	Asn	His	Gly	Ser	Gln	Ser	Pro		
	515					520						525					
Arg	Asn	Val	Glu	Glu	Arg	Met	Asn	Gly	Ser	His	Phe	Lys	Asp	Glu	Lys		
530					535					540							
Ala	Leu	Val	Thr	Ser	Gln	Asn	Ser	Asp	Leu	Leu	Asp	Asp	Glu	Glu	Val		
545					550				555					560			
Glu	Asp	Glu	Val	Leu	Leu	Asp	Glu	Glu	Asp	Glu	Asp	Asn	Asp	Ile	Thr		
			565					570					575				
Gly	Lys	Thr	Gly	Lys	Glu	Pro	Val	Thr	Ser	Asn	Leu	His	Glu	Gly	Asn		
	580							585				590					
Pro	Glu	Asp	Asp	Tyr	Glu	Glu	Thr	Ser	Ala	Leu	Glu	Met	Ser	Cys	Lys		
	595						600					605					
Thr	Ser	Pro	Val	Arg	Tyr	Lys	Glu	Glu	Glu	Tyr	Lys	Ser	Gly	Leu	Ser		
	610				615						620						
Ala	Leu	Asp	His	Ile	Arg	His	Phe	Thr	Asp	Ser	Leu	Lys	Met	Arg	Lys		
625					630				635					640			
Met	Glu	Asp	Asn	Gln	Tyr	Ser	Glu	Ala	Glu	Leu	Ser	Ser	Phe	Ser	Thr		
		645						650					655				
Ser	His	Val	Pro	Glu	Glu	Leu	Lys	Gln	Pro	Leu	His	Arg	Lys	Ser	Lys		
	660						665					670					
Ser	Gln	Ala	Tyr	Ala	Met	Met	Leu	Ser	Leu	Ser	Asp	Lys	Glu	Ser	Leu		
	675						680				685						
His	Ser	Thr	Ser	His	Ser	Ser	Ser	Asn	Val	Trp	His	Ser	Met	Ala	Arg		
	690				695						700						
Ala	Ala	Ala	Glu	Ser	Ser	Ala	Ile	Gln	Ser	Ile	Ser	His	Val				
705					710				715								

```
<210> 78
<211> 4950
<212> DNA
<213> Homo sapiens
```

```
<220>
<221> misc_feature
<222> (1)...(4950)
<223> n = A,T,C or G
```

[illegible]

gataccacaaa	agtttccatg	cagtactcct	caotcagcat	tttcaatggt	tgaagaggac	420
tttcagcaaa	aactcgaaa	cgagaatgat	ctccagaga	tacacacgat	ccaggagtgt	480
agggaatgtg	acccaagttt	tctgtatttg	caaaagcttg	agaaacacat	gctgtccaat	540
actgaagaga	gggaatacaa	gtgtgatcag	tgctccaaag	catttaactg	gaagtccaat	600
ttaatttcgc	acccaagatg	acatgacagt	ggaagacact	atgaatgtga	aaactgtgcc	660
aaggttttca	cggagccctg	caaccttcag	cgccacattc	gctctcagca	gtgcggtgcc	720
cgggccccat	catgcgccga	gtgtggccaa	acgtttgcga	cttcgtcggt	ccctaaacaa	780
caacaagaca	tccacagcag	tgtgaagccc	tttatctgtg	aggcttcgca	taaatctcat	840
actcagtttt	caaacctttg	cogtcataag	cgcagtcagt	ctgattgcag	aaocccaaat	900
aagtgcacaa	actgtggaca	aatgttcacg	actacgtctt	ccttaaataa	acacaggagg	960
ttttgtgagg	gcaggaacca	ttttgcggca	ggtggatttt	ttggccaagg	catttcaact	1020
cctggaaacc	cagctatgga	taaaaacgtc	atggttaata	tgagtcatcg	caaccocggc	1080
cttgtgactc	actttggcgc	caataggcat	cctgctggct	ttacotttcc	aacagctcct	1140
ggattttctt	ttagcttccc	tggtctgttt	ccttcoggtc	tgtaccacag	gcctcctttg	1200
ataccctgta	gtttctctgt	taaaagacta	tcaagtaact	aacagacaaa	caaaagtcga	1260
agtcacctca	ctacacatcc	tcagataactg	ccagctacac	aggatatttt	gaaggccact	1320
ctcaaaaccc	gcagcaataag	ggacaataag	cagctggagc	tcacgcccga	gaggtctctt	1380
gaagagagcg	cctttgagaa	aatcagtgac	cagtcagaga	gtagtgaact	tgatgatgtc	1440
agtcacccaa	gtggcagtg	cctggaaaca	acotcgggct	ctgatctgga	aaagtacatt	1500
gaagatgata	aagagaaatt	taaaagaaat	ggtaaaatgt	aagagacaaa	agtaagccct	1560
cttcagaatc	tggtctcaat	aaataataag	aaagaataca	gcaatcattc	caattttctc	1620
coactctttg	aggagccagc	tgoggtgtca	ggagctgtga	atgattctat	aaaggtctat	1680
gcttctattg	ctgaaaaata	ctttggttca	acaggactgg	tggggctgca	agacaaaaaa	1740
gttgagcttt	taocttacc	ttccatgttt	ccctcccatc	atttctccaa	attctctcaa	1800
lcaatgtacc	catttctgta	tagagacttg	agatcgttac	ctttgaaaa	ggaaccccaa	1860
tcaccaggtg	aagcaagaaa	actgcagaag	ggcagctctg	agtcocctct	tgactcacc	1920
actaagcgaa	aggatgagaa	gcccttgact	cagtcocctc	ccaagctctc	agtcgacact	1980
gccacaagcc	aagaccagcc	cctggatcta	agtatgggca	gtaggagtga	agccagtggt	2040
acaaagctga	ctgagccctg	aaaaaacacc	gtgtttgggg	gaaaaaaaag	aagcaagctc	2100
gaatcaagac	ctgcttcaga	tggttccctg	cagcatgcga	gacccactcc	tttctttatg	2160
gaacctattt	acagagtaga	gaaaaagaaa	ctaaactgac	ctactgaagc	tttaaaaagag	2220
aaataactga	ggccttctcc	aggattcttg	tttaccacc	aattccaaat	gcctgatcac	2280
agaacttgga	tgccagagaa	tgaaaaacatg	gcagaaaaag	tagagagctt	cagtgcocctg	2340
aaaaactgag	ccagtgaagt	cttacagtca	gtgcctctca	tggtcaactt	cagggcgccct	2400
cccaactgcc	tgccagagaa	ccttctgcgg	aagggaagag	agcgctatcc	ctgcagatca	2460
tgtggcaga	tttttccaag	gtctgcaaac	ctaacacagc	acttgagaac	ccacacagga	2520
gagcagcttc	acagatgcga	atactgtgac	agatcattta	goatatcttc	taacttgcaa	2580
aggcgtatct	gcaacatcca	caataaagag	aagccattta	agtgctcaat	gtgatgatgg	2640
tggtttgttc	aaocaaacaa	tttagacaga	caoctaaaga	aacatgagaa	tgggaaatcg	2700
tcocgtacag	caacatcgct	gcctcattct	gaactggaaa	gtacaggtgc	gattctggat	2760
gacaaaaga	atgcttactt	cacagaaatt	cgaattttca	ttgggaacag	caacctatgc	2820
agccactccc	ccaggaatgt	ggaggagaga	atgaattgca	gtcattttaa	gtatgaaaag	2880
gctttgggtga	ccagtcacaa	ttcagacttg	ctggatgatg	aagaagtgtg	agatgaggtg	2940
ttgttatgtg	aggaggatga	agacaaatgat	attactggaa	aaacagggaa	ggaacccagt	3000
acaagtattt	tacatgaag	aaacccctgag	gatgactatg	aagaaaccag	tgccctggag	3060
atgagttgca	agacatcccc	agtgagggtat	aaagaggaa	aatataaaa	ttaaactttc	3120
gctctagatc	atataaggca	cttcacagat	agcctcaaaa	tgaggaaaa	ggaagataat	3180
caattattctg	aagctgagct	gtotttcttt	agtacttccc	atgtgccaga	ggaacttaag	3240
cagccgttacc	acagaaaagt	caaatcgacg	gcataatgta	tgatgtctgc	actgtctgac	3300
aaggagctcc	tgcgactctc	atcccaacgt	tcttccaaag	tgtggcacag	tatggccagc	3360
gctgcggcgg	aatccagctc	tatccagctc	ataagccag	tatgacgtta	tcaaggttga	3420
ccagagtggc	acccaagttc	acagtagcat	ggctctttca	tataggacta	tttcaaacag	3480
gtctgagcag	aaacctgcag	gtgcactcat	ctaaagtcta	ctaaagtcta	gtgaccttaa	3540
actgcatgat	ttaaaaaaga	aaagaaagaa	aaaagaaact	atttattctc	gatattttgt	3600
tttgacagcg	aaagcgactc	gctgacttct	ggaaagatcaa	tcaatgcgac	ttaaagtgat	3660
tcagtgaaaa	caaaaaactt	ggtgggctga	aggaactctc	cagttttacc	caaccttagg	3720
tatgggtggg	tgagaaggcg	agttgagatg	cgagcattga	ctgaatgaac	caactccatg	3780
aaaactgaatt	ctcttttgta	caagatcacc	tgacatgatt	gggaacagtt	gcttttaatt	3840

```

acagatttaa tttttttt cgttaaagtt ttatgtaatt taaccctttg aagacagaag 3900
tagttggatg aaatgcacag tcaattatta tagaaactga taacagggag tacttgttcc 3960
cccttttgcc ttcttaagta cattgtttaa aactagggaa aaagggtagt tgtattattgt 4020
aaactatgga tgttaacact tcaaagaggt taagtcatgt argtaacctta ttcataacca 4080
tcagocgtgt accactaata aattgttttg caaatcctgt taataacatc ttaatttttag 4140
acaatcatgt cactgttttt aatgtttatt tttttgtgtg tgttgocgtg atcatgtatt 4200
tatttgttgg caaacatttg ttgtttgatt aaaatagcac tgttccagtc agccactact 4260
ttatgacgtc tgaggcacac ccctttccga atttcaagga ccaaggtgac ccgacctgtg 4320
tatgagatgt ccaaatgtgt ttgtgctttt cttaacattc cttttgtttg gtttgttttg 4380
ttttccttct taatgaacta aatcacgaata gatgcaactt agtttttgta atactgaaat 4440
nogatccaat tgtataaacg attataattt ctttcatgga agcatgattc ttctgtattaa 4500
aaactgtact ccatatttta tgcgtggtgt ctgcaagcct gtgcgattgt atgttcatgt 4560
taactctatt tgtaaaaatga agtgttccca accttatngt taaaagagag aangtaaata 4620
acagactgta ttacgttatt ttgcccctta ttgagggaacc agattttggtt tctttttgtt 4680
tgtaactcca ttttngaana aatcacgaag ttgaggtact ttcttcaaat gctttgtaca 4740
atataaactg ttatgccttt cagtgcatta ctatgggagg agcaactaaa aaataaagac 4800
ttacaaaaag gagtattttt aaagaacaaa aacttgaggg gggggccoggt acccaattcg 4860
ccctatagtg agtagtatta caattcactg gcgctgcttt tacaaactgc tgactgggaa 4920
aacctgggtt taccacaact aatcgtcttn

```

<210> 79

<211> 1051

<212> PRT

<213> Homo sapiens

<400> 79

```

Met Lys Ser Glu Asp Tyr Pro His Glu Thr Met Ala Pro Asp Ile His
1 5 10 15
Glu Glu Arg Gln Tyr Arg Cys Glu Asp Cys Asp Gln Leu Phe Glu Ser
20 25 30
Lys Ala Glu Leu Ala Asp His Gln Lys Phe Pro Cys Ser Thr Pro His
35 40 45
Ser Ala Phe Ser Met Val Glu Glu Asp Phe Gln Gln Lys Leu Glu Ser
50 55 60
Glu Asn Asp Leu Gln Glu Ile His Thr Ile Gln Glu Cys Lys Glu Cys
65 70 75 80
Asp Gln Val Phe Pro Asp Leu Gln Ser Leu Glu Lys His Met Leu Ser
85 90 95
His Thr Glu Glu Arg Glu Tyr Lys Cys Asp Gln Cys Pro Lys Ala Phe
100 105 110
Asn Trp Lys Ser Asn Leu Ile Arg His Gln Met Ser His Asp Ser Gly
115 120 125
Lys His Tyr Glu Cys Glu Asn Cys Ala Lys Val Phe Thr Asp Pro Ser
130 135 140
Asn Leu Gln Arg His Ile Arg Ser Gln His Val Gly Ala Arg Ala His
145 150 155 160
Ala Cys Pro Glu Cys Gly Lys Thr Phe Ala Thr Ser Ser Gly Leu Lys
165 170 175
Gln His Lys His Ile His Ser Ser Val Lys Pro Phe Ile Cys Glu Val
180 185 190
Cys His Lys Ser Tyr Thr Gln Phe Ser Asn Leu Cys Arg His Lys Arg
195 200 205
Met His Ala Asp Cys Arg Thr Gln Ile Lys Cys Lys Asp Cys Gly Gln
210 215 220
Met Phe Ser Thr Thr Ser Ser Leu Asn Lys His Arg Arg Phe Cys Glu
225 230 235 240
Gly Lys Asn His Phe Ala Ala Gly Gly Phe Gly Gln Gly Ile Ser
245 250 255

```